# ALL DAY PAIN RELIEF- naproxen sodium tablet, coated VALU MERCHANDISERS COMPANY

-----

1145-BST-2021-1117

**Drug Facts** 

### **Active ingredient (in each caplet)**

Naproxen sodium 220 mg (naproxen 200 mg) (NSAID <sup>1</sup>)

1 nonsteroidal anti-inflammatory drug

#### **Purpose**

Pain reliever/fever reducer

#### Uses

- temporarily relieves minor aches and pains due to:
  - minor pain of arthritis
  - muscular aches
  - backache
  - menstrual cramps
  - headache
  - toothache
  - the common cold
- temporarily reduces fever

### Warnings

### Allergy alert

Naproxen sodium may cause a severe allergic reaction, especially in people allergic to aspirin. Symptoms may include:

- hives
- facial swelling
- asthma (wheezing)
- shock
- skin reddening
- rash
- blisters

If an allergic reaction occurs, stop use and seek medical help right away.

### Stomach bleeding warning

This product contains an NSAID, which may cause severe stomach bleeding. The chance is higher if you:

- are age 60 or older
- have had stomach ulcers or bleeding problems
- take a blood thinning (anticoagulant) or steroid drug
- take other drugs containing prescription or nonprescription NSAIDs (aspirin, ibuprofen, naproxen, or others)
- have 3 or more alcoholic drinks every day while using this product
- take more or for a longer time than directed

#### Do not use

- if you have ever had an allergic reaction to any other pain reliever/fever reducer
- right before or after heart surgery

#### Ask a doctor before use if

- stomach bleeding warning applies to you
- you have a history of stomach problems, such as heartburn
- you have high blood pressure, heart disease, liver cirrhosis, or kidney disease
- you are taking a diuretic
- you have asthma
- you have problems or serious side effects from taking pain relievers or fever reducers

### Ask a doctor or pharmacist before use if you are

- under a doctor's care for any serious condition
- taking any other drug

### When using this product

- take with food or milk if stomach upset occurs
- long term continuous use may increase the risk of heart attack or stroke

### Stop use and ask a doctor if

- side effects occur. You may report side effects to FDA at 1-800-FDA-1088.
- you experience any of the following signs of stomach bleeding:
  - feel faint
  - vomit blood
  - have bloody or black stools
  - have stomach pain that does not get better
- pain gets worse or lasts more than 10 days
- fever gets worse or lasts more than 3 days
- you have difficulty swallowing
- it feels like the pill is stuck in your throat
- you develop heartburn
- redness or swelling is present in the painful area
- any new symptoms appear

**If pregnant or breast-feeding,** ask a health professional before use. It is especially important not to use naproxen sodium during the last 3 months of pregnancy unless definitely directed to do so by a doctor because it may cause problems in the unborn

child or complications during delivery.

**Keep out of reach of children.** In case of overdose, get medical help or contact a Poison Control Center right away.

#### **Directions**

- do not take more than directed
- the smallest effective dose should be used
- do not take longer than 10 days, unless directed by a doctor (see Warnings)
- drink a full glass of water with each dose

adults and children 12 years and older	<ul> <li>take 1 caplet every 8 to 12 hours while symptoms last</li> <li>for the first dose you may take 2 caplets within the first hour</li> <li>do not exceed 2 caplets in any 8- to 12-hour period</li> <li>do not exceed 3 caplets in a 24-hour period</li> </ul>
children under 12 years	<ul><li>ask a doctor</li></ul>

#### Other information

- each caplet contains: sodium 20 mg
- store at 20°-25°C (68°-77°F). Avoid high humidity and excessive heat above 40°C (104°F).
- retain carton for complete product information

### **Inactive ingredients**

croscarmellose sodium, FD&C blue #2, macrogol, magnesium stearate, polyvinyl alcohol, povidone, pregelatinized starch, talc, titanium dioxide

#### PRINCIPAL DISPLAY PANEL

COMPARE TO THE ACTIVE INGREDIENT IN ALEVE® CAPLETS†

Best Choice®

**ALL DAY** 


Pain Relief

Naproxen Sodium Tablets, 220mg

Pain Reliever / Fever Reducer (NSAID)

24 CAPLETS\*\*

#### \*\*CAPSULE-SHAPED TABLETS



#### **ALL DAY PAIN RELIEF**

naproxen sodium tablet, coated

Product Information			
Product Type	HUMAN OTC DRUG	Item Code (Source)	NDC:63941-245
Route of Administration	ORAL		

Active Ingredient/Active Moiety			
Ingredient Name	<b>Basis of Strength</b>	Strength	
NAPROXEN SODIUM (UNII: 9TN87S3A3C) (NAPROXEN - UNII:57Y76R9ATQ)	NAPROXEN SODIUM	220 mg	

Inactive Ingredients		
Ingredient Name	Strength	
CROSCARMELLOSE SODIUM (UNII: M280L1HH48)		
FD&C BLUE NO. 2 (UNII: L06K8R7DQK)		
POLYETHYLENE GLYCOL, UNSPECIFIED (UNII: 3WJQ0SDW1A)		
MAGNESIUM STEARATE (UNII: 70097M6I30)		
POLYVINYL ALCOHOL (UNII: 532B59J990)		
POVIDONE (UNII: FZ 989GH94E)		
STARCH, CORN (UNII: O8232NY3SJ)		
TALC (UNII: 7SEV7J4R1U)		
TITANIUM DIOXIDE (UNII: 15FIX9V2JP)		

Product Characteristics				
Color	blue	Score	no score	
Shape	OVAL (Biconvex)	Size	12mm	
Flavor		Imprint Code	17	
Contains				

Packaging				
#	Item Code	Package Description	Marketing Start Date	Marketing End Date
1	NDC:63941- 245-01	1 in 1 CARTON	06/10/2010	
1		24 in 1 BOTTLE, PLASTIC; Type 0: Not a Combination Product		
2	NDC:63941- 245-02	1 in 1 CARTON	05/19/2010	
2		50 in 1 BOTTLE, PLASTIC; Type 0: Not a Combination Product		
3	NDC:63941- 245-03	1 in 1 CARTON	05/19/2010	
3		100 in 1 BOTTLE, PLASTIC; Type 0: Not a Combination Product		

Marketing Information			
Marketing Category	Application Number or Monograph Citation	Marketing Start Date	Marketing End Date
ANDA	ANDA079096	05/19/2010	

## Labeler - VALU MERCHANDISERS COMPANY (868703513)

Revised: 11/2021 VALU MERCHANDISERS COMPANY