NITE TIME- acetaminophen, dextromethorphan hydrobromide and doxylamine succinate solution Kroger Company

Disclaimer: Most OTC drugs are not reviewed and approved by FDA, however they may be marketed if they comply with applicable regulations and policies. FDA has not evaluated whether this product complies.

Kroger Co. Nite Time Drug Facts

Active ingredients (in each 30 mL dose cup)

Acetaminophen 650 mg

Dextromethorphan HBr 30 mg

Doxylamine succinate 12.5 mg

Purpose

Pain reliever/fever reducer Cough suppressant Antihistamine

Uses

temporarily relieves common cold/flu symptoms:

- fever
- cough due to minor throat and bronchial irritation
- sore throat
- minor aches and pains
- runny nose and sneezing
- headache

Warnings

Liver warning: This product contains acetaminophen. The maximum daily dose of this product is 4 doses (2,600 mg acetaminophen) in 24 hours. Severe liver damage may occur if you take

- more than 4,000 mg of acetaminophen in 24 hours
- with other drugs containing acetaminophen
- 3 or more alcoholic drinks every day while using this product

Sore throat warning: If sore throat is severe, lasts for more than 2 days, occurs with or is followed by fever, headache, rash, nausea, or vomiting, consult a doctor promptly.

Do not use

- with any other drug containing acetaminophen (prescription or nonprescription). If you are not sure whether a drug contains acetaminophen, ask a doctor or pharmacist.
- if you are now taking a prescription monoamine oxidase inhibitor (MAOI) (certain drugs for

depression, psychiatric, or emotional conditions, or Parkinson's disease), or for 2 weeks after stopping the MAOI drug. If you do not know if your prescription drug contains an MAOI, ask a doctor or pharmacist before taking this product.

- if you have ever had an allergic reaction to this product or any of its ingredients
- to make a child sleepy

Ask a doctor before use if you have

- liver disease
- glaucoma
- cough that occurs with too much phlegm (mucus)
- a breathing problem such as emphysema or chronic bronchitis
- trouble urinating due to an enlarged prostate gland
- persistent or chronic cough as occurs with smoking, asthma, or emphysema
- a sodium-restricted diet

Ask a doctor or pharmacist before use if you are

- taking sedatives or tranquilizers
- taking the blood thinning drug warfarin

When using this product

- excitability may occur, especially in children
- marked drowsiness may occur
- avoid alcoholic drinks
- be careful when driving a motor vehicle or operating machinery
- alcohol, sedatives, and tranquilizers may increase drowsiness

Stop use and ask a doctor if

- pain or cough gets worse or lasts more than 7 days
- fever gets worse or lasts more than 3 days
- redness or swelling is present
- new symptoms occur
- cough comes back or occurs with rash or headache that lasts. These could be signs of a serious condition.

If pregnant or breast-feeding,

ask a health professional before use.

Keep out of reach of children.

In case of overdose, get medical help or contact a Poison Control Center right away. (1-800-222-1222) Quick medical attention is critical for adults as well as for children even if you do not notice any signs or symptoms.

Directions

take only as directed - see Liver warning

- use dose cup
- do not exceed 4 doses per 24 hrs

adults & children 12 yrs & over	30 mL (2 TBSP) every 6 hrs
children 4 to under 12 yrs	ask a doctor
children under 4 yrs	do not use

 when using other Daytime or Nighttime products, carefully read each label to insure correct dosing

Other information

- each 30 mL dose cup contains: sodium 39 mg
- store at 20-25°C (68-77°F)

Inactive ingredients

alcohol, anhydrous citric acid, FD&C blue no. 1, FD&C red no. 40, flavor, high fructose corn syrup, polyethylene glycol, propylene glycol, purified water, saccharin sodium, sodium citrate

Questions or comments?

1-800-632-6900

Package/Label Principal Display Panel

COMPARE TO the active ingredients of VICKS® NYQUIL® COLD & FLU

See back panel

Nite Time

COLD & FLU

Acetaminophen, Pain Reliever/Fever Reducer

Dextromethorphan HBr, Cough Suppressant

Doxylamine Succinate, Antihistamine

Relieves:

Aches, Fever, Sore Throat

Sneezing, Runny Nose

Cough

ALCOHOL 10%

Cherry Flavor

12 FL OZ (354 mL)



DO NOT USE IF PRINTED NECKBAND IS BROKEN OR MISSING

Drug Facts

Uses temporarily relieves common cold/flu symptoms: ■ fever ■ cough due to minor throat and bronchial irritation ■ sore throat ■ minor aches and pains ■ runny nose and sneezing ■ headache

Warnings

Liver warning: This product contains acetaminophen. The maximum daily dose of this product is 4 doses (2,600 mg acetaminophen) in 24 hours. Severe liver damage may occur if you take

- more than 4,000 mg of acetaminophen in 24 hours
- with other drugs containing acetaminophen
- 3 or more alcoholic drinks every day while using this product Sore throat warning: If sore throat is severe, lasts for more than 2 days, occurs with or is followed by fever, headache, rash, nausea, or vomiting, consult a doctor promptly.

Do not use with any other drug containing acetaminophen (prescription or nonprescription). If you are not sure whether a drug contains acetaminophen, ask a doctor or pharmacist.

- if you are now taking a prescription monoamine oxidase inhibitor (MAOI) (certain drugs for depression, psychiatric, or emotional conditions, or Parkinson's disease), or for 2 weeks after stopping the MAOI drug, If you do not know if your prescription drug contains an MAOI, ask a doctor or pharmacist before taking this product.
- if you have ever had an allergic reaction to this product or any of its ingredients to make a child sleepy
- Ask a doctor before use if you have liver disease glaucome cough that occurs with too much phlegm (mucus) persistent or chronic cough as occurs with smoking, asthma, or emphysema
- chronic cough as occurs with smoking, asthma, or emphysema
 a breathing problem such as emphysema or chronic bronchitis
 trouble urinating due to an enlarged
- prostate gland a sodium-restricted diet

 Ask a doctor or pharmacist before use if
 you are taking sedatives or tranquilizers
 taking the blood thinning drug warfarin

: 45940 45 F3

NITE TIME

acetaminophen, dextromethorphan hbr, doxylamine succinate solution

Product Information			
Product Type	HUMAN OTC DRUG LABEL	Item Code (Source)	NDC:30142-459
Route of Administration	ORAL	DEA Schedule	

Active Ingredient/Active Moiety			
Ingredient Name	Basis of Strength	Strength	
ACETAMINO PHEN (ACETAMINO PHEN)	ACETAMINOPHEN	650 mg in 30 mL	
DEXTRO METHO RPHAN HYDRO BRO MIDE (DEXTRO METHO RPHAN)	DEXTROMETHORPHAN HYDROBROMIDE	30 mg in 30 mL	
DO XYLAMINE SUCCINATE (DO XYLAMINE)	DOXYLAMINE SUCCINATE	12.5 mg in 30 mL	

Inactive Ingredients		
Ingredient Name	Strength	
ALCOHOL		
ANHYDRO US CITRIC ACID		
FD&C BLUE NO. 1		
FD&C RED NO. 40		
HIGH FRUCTO SE CORN SYRUP		
POLYETHYLENE GLYCOLS		
PROPYLENE GLYCOL		
WATER		
SACCHARIN SO DIUM		

SODIUM CITRATE

Product Characteristics			
Color	RED (Clear/Dark Red)	Score	
Shape		Size	
Flavor	CHERRY (Menthol Aroma)	Imprint Code	
Contains			

P	Packaging			
#	Item Code	Package Description	Marketing Start Date	Marketing End Date
1	NDC:30142-459-38	295 mL in 1 BOTTLE		
2	NDC:30142-459-30	177 mL in 1 BOTTLE		
3	NDC:30142-459-34	237 mL in 1 BOTTLE		
4	NDC:30142-459-40	354 mL in 1 BOTTLE		

Marketing Information			
Marketing Category	Application Number or Monograph Citation	Marketing Start Date	Marketing End Date
OTC monograph final	part341	08/26/2011	

Labeler - Kroger Company (006999528)

Revised: 11/2014 Kroger Company