LIDOCAINE 2% - lidocaine injection Covetrus North America

Disclaimer: This drug has not been found by FDA to be safe and effective, and this labeling has not been approved by FDA. For further information about unapproved drugs, click here.

Lidocaine 2%

Injectable

Local anesthetic

CAUTION: Federal law restricts this drug to use by or on the order of a licensed veterinarian.

For animal use only

Keep out of the reach of children

Indications

Lidocaine is a potent local anesthetic for producing epidural, nerve conduction, and infiltration anesthesia.

Contraindications

Lidocaine is contraindicated in animals with a known hypersensitivity to the drug.

Precautions

Lidocaine is usually well tolerated. Nevertheless, as with all local anesthetics, untoward effects may occur due to hypersensitivity, faulty technique, overdosage and inadvertent intravascular or subarachnoid injection. In case of respiratory arrest, immediate resuscitation with oxygen is indicated.

Dosage and administration

Epidural:Cattle and Horses - 5 to 15 mL Dogs and Cats - 1 mL per 10 pounds of body weight. **Nerve Block:**Cattle and Horses - 5 to 20 mL

Infiltration:Dilute to 0.5% concentration (1 mL of 2% solution diluted with 3 mL of sterile water = 4 mL of a 0.5% solution)

Composition

Each mL of sterile aqueous solution contains: lidocaine hydrochloride.......2.0% propylene glycol.......5.2% sodium chloride.......0.5% sodium lactate......0.5% with methylparaben 0.15%, sodium metabisulfite 0.10%, propylparaben 0.03%, and disodium edetate 0.001% as preservatives.

Store at controlled room temperature between 15° and 30°C (59°-86°F)

TAKE TIME OBSERVE LABEL DIRECTIONS



Indications: Lidocaine is a potent local anesthetic for producing epidural, nerve conduction, and infiltration anesthesia.

Contraindications: Lidocaine is contraindicated in animals with a known hypersensitivity to the drug.

Precautions: Lidocaine is usually well tolerated. Nevertheless, as with all local anesthetics, untoward effects may occur due to hypersensitivity, faulty technique, overdosage, and inadvertent intravascular or subarachnoid injection. In case of respiratory arrest, immediate resuscitation with oxygen is indicated.

Dosage and administration:

Cattle and horses- 5 to 15 mL.

Dogs and cats—1 mL per 10 pounds of body weight. Nerve block:

Cattle and horses- 5 to 20 mL.

Infiltration:

Dilute to 0.5% concentration (1 mL of 2% solution diluted with 3 mL of sterile water = 4 mL of a 0.5% solution).



NDC: 11695-4149-2

Lidocaine 2% Injectable

Local anesthetic

CAUTION: Federal law restricts this drug to use by or on the order of a licensed veterinarian.

For animal use only Keep out of the reach of children

Volume: 250 mL (8.454 fl oz)

Composition: Each mL of sterile aqueous solution contains:	
lidocaine hydrochloride	2.0%
propylene glycol	5.2%
sodium chloride	0.5%
sodium lactate	
with methylparaben 0.15%, sodium metabisulfite 0.10%,	
propylparaben 0.03% and disodium edetate 0.001	% as
preservatives.	

Store at controlled room temperature between 15° and 30°C (59°-86°F).



Questions? (855) 724-3461

AH-014583-02 ISS: 0120 REV: 0120 L-2000-05

Lot no:

Distributed by: Covetrus North America 400 Metro Place North

Reorder #014583

Dublin, 0H 43017 covetrus.com

Exp. date:

LIDOCAINE 2%

lidocaine injection

Product Information

Product Type PRESCRIPTION ANIMAL DRUG Item Code (Source) NDC:11695-4149

EPIDURAL, INFILTRATION Route of Administration

Active Ingredient/Active Moiety

Basis of Strength Ingredient Name Strength LIDO CAINE HYDRO CHLO RIDE (UNII: V13007Z41A) (LIDO CAINE -LIDOCAINE HYDROCHLORIDE 0.02 gUNII:98PI200987) **ANHYDROUS** in 1 mL

Packaging Marketing Start Date Marketing End Date Item Code Package Description 1 NDC:11695-4149-1 100 mL in 1 BOTTLE 2 NDC:11695-4149-2 250 mL in 1 BOTTLE

Marketing Information Marketing End Date Marketing Category Application Number or Monograph Citation Marketing Start Date unapproved drug other 01/20/2020

Revised: 1/2021 Covetrus North America