VICKS DAYQUIL SEVERE COLD AND FLU- acetaminophen, dextromethorphan hydrobromide, guaifenesin, and phenylephrine hydrochloride solution
The Procter & Gamble Manufacturing Company

Vicks ®DayQuil ® Severe Cold & Flu Liquid

Drug Facts

Active ingredients (in each 15 mL)

Acetaminophen 325 mg
Dextromethorphan HBr 10 mg
Guaifenesin 200 mg
Phenylephrine HCl 5 mg

Purpose

Pain reliever/fever reducer
Cough suppressant
Expectorant
Nasal decongestant

Uses

- temporarily relieves common cold/flu symptoms:
  - nasal congestion
  - sinus congestion & pressure
  - cough due to minor throat & bronchial irritation
  - minor aches & pains
  - headache
  - fever
  - sore throat
  - reduces swelling of nasal passages
  - temporarily restores freer breathing through the nose
  - promotes nasal and/or sinus drainage
  - helps loosen phlegm (mucus) and thin bronchial secretions to rid the bronchial passageways of bothersome mucus and make coughs more productive.

Warnings

Liver warning: This product contains acetaminophen. Severe liver damage may occur if

• adult takes more than 4 doses (30 mL each) in 24 hrs, which is the maximum daily amount for this product

• child takes more than 4 doses (15 mL each) in 24 hrs, which is the maximum daily amount for this product

• taken with other drugs containing acetaminophen

• adult has 3 or more alcoholic drinks every day while using this product

Allergy Alert: Acetaminophen may cause severe skin reactions.

Symptoms may include:

• skin reddening

• blisters

• rash

If a skin reaction occurs, stop use and seek medical help right away.

Sore throat warning: If sore throat is severe, persists for more than 2 days, is accompanied or followed by fever, headache, rash, nausea, or vomiting, consult a doctor promptly.

Do not use

• with any drug containing acetaminophen (prescription or nonprescription). If you are not sure whether a drug contains acetaminophen, ask a doctor or pharmacist.

• if you are now taking a prescription monoamine oxidase inhibitor (MAOI) (certain drugs for depression, psychiatric or emotional conditions, or Parkinson's disease), or for 2 weeks after stopping the MAOI drug. If you do not know if your prescription drug contains an MAOI, ask a doctor or pharmacist before taking this product.

Ask a doctor before use if you have

- liver disease
- heart disease
- high blood pressure
- thyroid disease
- diabetes
- trouble urinating due to enlarged prostate gland
- cough that occurs with too much phlegm (mucus)
- persistent or chronic cough such as occurs with smoking, asthma, chronic bronchitis, or emphysema
- a sodium-restricted diet

Ask a doctor or pharmacist before use if you are

taking the blood thinning drug warfarin.

When using this product,

do not use more than directed.

Stop use and ask a doctor if

- you get nervous, dizzy or sleepless

- pain, nasal congestion, or cough gets worse or lasts more than 5 days (children) or 7 days (adults)
- fever gets worse or lasts more than 3 days
- redness or swelling is present
- new symptoms occur
- cough comes back or occurs with rash or headache that lasts.

  These could be signs of a serious condition.

If pregnant or breast-feeding,

ask a health professional before use.

Keep out of reach of children.

In case of overdose, get medical help or contact a Poison Control Center right away. Quick medical attention is critical for adults as well as for children even if you do not notice any signs or symptoms.

Directions

- take only as directed
- only use the dose cup provided
- do not exceed 4 doses per 24 hrs

adults & children 12 yrs & over	30 mL every 4 hrs
children 6 to under 12 yrs	15 mL every 4 hrs
children 4 to under 6 yrs	ask a doctor
children under 4 yrs	do not use

Other information

- **each 15 mL contains:** sodium 47 mg
- store at no greater than 25ºC and do not refrigerate

Inactive ingredients

citric acid, FD&C Yellow No. 6, flavor, glycerin, propylene glycol, purified water, saccharin sodium, sodium benzoate, sodium chloride, sodium citrate, sorbitol, sucralose, xanthan gum

Questions?

1-800-362-1683

TAMPER EVIDENT: Do not use if printed shrinkband is broken or rmissing.

DIST. BY PROCTER & GAMBLE, CINCINNATI, OH 45202

PRINCIPAL DISPLAY PANEL - 354 ml Bottle Label

VICKS®

DayQuil™

SEVERE

COLD & FLU

Acetaminophen, Guaifenesin, Phenylephrine HCl, Dextromethorphan HBr

Headache, Fever, Sore Throat, Minor Aches & Pains

Chest Congestion, Thins & Loosens Mucus

Nasal Congestion, Sinus Pressure

Cough

Non-Drowsy

Alcohol Free

12 FL OZ (354 ml)



VICKS DAYQUIL SEVERE COLD AND FLU

acetaminophen, dextromethorphan hydrobromide, guaifenesin, and phenylephrine hydrochloride solution

Product Information			
Product Type	HUMAN OTC DRUG	**Item Code (Source)**	NDC:37000-810
Route of Administration	ORAL		

Active Ingredient/Active Moiety

Ingredient Name	Basis of Strength	Strength
ACETAMINOPHEN (UNII: 362O9ITL9D) (ACETAMINOPHEN - UNII:362O9ITL9D)	ACETAMINOPHEN	325 mg in 15 mL
DEXTROMETHORPHAN HYDROBROMIDE (UNII: 9D2RTI9KYH) (DEXTROMETHORPHAN - UNII:7355X3ROTS)	DEXTROMETHORPHAN HYDROBROMIDE	10 mg in 15 mL
GUAIFENESIN (UNII: 495W7451VQ) (GUAIFENESIN - UNII:495W7451VQ)	GUAIFENESIN	200 mg in 15 mL
PHENYLEPHRINE HYDROCHLORIDE (UNII: 04JA59TNSJ) (PHENYLEPHRINE - UNII:1WS297W6MV)	PHENYLEPHRINE HYDROCHLORIDE	5 mg in 15 mL

Inactive Ingredients

Ingredient Name	Strength
CITRIC ACID MONOHYDRATE (UNII: 2968PHW8QP)	
FD&C YELLOW NO. 6 (UNII: H77VEI93A8)	
GLYCERIN (UNII: PDC6A3C0OX)	
PROPYLENE GLYCOL (UNII: 6DC9Q167V3)	
WATER (UNII: 059QF0KO0R)	
SACCHARIN SODIUM (UNII: SB8ZUX40TY)	
SODIUM BENZOATE (UNII: OJ245FE5EU)	
SODIUM CHLORIDE (UNII: 451W47IQ8X)	
SODIUM CITRATE (UNII: 1Q73Q2JULR)	
SORBITOL (UNII: 506T60A25R)	
SUCRALOSE (UNII: 96K6UQ3ZD4)	
XANTHAN GUM (UNII: TTV12P4NEE)	

Product Characteristics

Color	orange	**Score**	
Shape		**Size**	
Flavor	BERRY	**Imprint Code**	
Contains			

Packaging

#	Item Code	Package Description	Marketing Start Date	Marketing End Date
1	NDC:37000-810-08	236 mL in 1 BOTTLE, PLASTIC; Type 0: Not a Combination Product	07/22/2013	
2	NDC:37000-810-12	354 mL in 1 BOTTLE, PLASTIC; Type 0: Not a Combination Product	07/22/2013	
3	NDC:37000-810-01	30 mL in 1 BOTTLE, PLASTIC; Type 0: Not a Combination Product	07/13/2021	
4	NDC:37000-810-04	4 in 1 PACKAGE	07/13/2021	
4		30 mL in 1 BOTTLE, PLASTIC; Type 0: Not a Combination Product		

Marketing Information

Marketing Category	Application Number or Monograph Citation	Marketing Start Date	Marketing End Date
OTC Monograph Drug	M012	07/22/2013	

Labeler - The Procter & Gamble Manufacturing Company (004238200)

Revised: 10/2023 The Procter & Gamble Manufacturing Company