

CLEAR LAX ORIGINAL PRESCRIPTION STRENGTH- polyethylene glycol 3350 powder, for solution

HyVee Inc

Hy-Vee, Inc. ClearLax® Drug Facts

Active ingredient (in each dose)

Polyethylene Glycol 3350, 17 g (cap filled to line)

Purpose

Osmotic Laxative

Use

- relieves occasional constipation (irregularity)
- generally produces a bowel movement in 1 to 3 days

Warnings

Allergy alert: Do not use if you are allergic to polyethylene glycol

Do not use

if you have kidney disease, except under the advice and supervision of a doctor

Ask a doctor before use if you have

- nausea, vomiting or abdominal pain
- a sudden change in bowel habits that lasts over 2 weeks
- irritable bowel syndrome

Ask a doctor or pharmacist before use if you are

taking a prescription drug

When using this product

you may have loose, watery, more frequent stools

Stop use and ask a doctor if

- you have rectal bleeding or your nausea, bloating, cramping or abdominal pain gets worse. These may be signs of a serious condition.
- you get diarrhea
- you need to use a laxative for longer than 1 week

If pregnant or breast-feeding,

ask a health professional before use.

Keep out of reach of children.

In case of overdose, get medical help or contact a Poison Control Center right away. (1-800-222-1222)

Directions

- do not take more than directed unless advised by your doctor
- the bottle top is a measuring cap marked to contain 17 grams of powder when filled to the indicated line (white section in cap)
- adults and children 17 years of age and older:
- fill to top of white section in cap which is marked to indicate the correct dose (17 g)
- stir and dissolve in any 4 to 8 ounces of beverage (cold, hot or room temperature) then drink
- use once a day
- use no more than 7 days
- children 16 years of age or under: ask a doctor

Other information

- store at 20°-25°C (68°-77°F)
- tamper-evident: do not use if printed foil seal under cap is missing, open or broken

Inactive ingredients

none

Questions or comments?

1-800-719-9260

Principal Display Panel

Compare to the Active Ingredient of MiraLAX®

ClearLax®

POLYETHYLENE GLYCOL 3350

POWDER FOR SOLUTION

Osmotic Laxative

Relieves Occasional Constipation / Irregularity

Softens Stool

Unflavored Powder – Grit Free

NET WT 8.3 OZ (238 g)

14 ONCE-DAILY DOSES

HyVee[®]

*Compare to
the Active Ingredient of
MiraLAX[®]

NDC 42507-306-02

ClearLax[®]

POLYETHYLENE GLYCOL 3350
POWDER FOR SOLUTION

Osmotic Laxative

- Relieves Occasional Constipation / Irregularity
- Softens Stool
- Unflavored Powder • Grit Free

NET WT 8.3 OZ (238 g)

14 ONCE-DAILY DOSES

: 306E1 2H F4

Peel Here

Polyethylene Glycol 3350 increases frequency of bowel movements and softens the stool.
Dissolves in Any Beverage

GLUTEN FREE

Drug Facts

Active ingredient (in each dose) Purpose

Polyethylene Glycol 3350, 17 g (cap filled to line).....Osmotic Laxative

Use

- relieves occasional constipation (irregularity)
- generally produces a bowel movement in 1 to 3 days

Warnings

Allergy alert: Do not use if you are allergic to polyethylene glycol

Do not use if you have kidney disease, except under the advice and supervision of a doctor

Ask a doctor before use if you have

- nausea, vomiting or abdominal pain
- a sudden change in bowel habits that lasts over 2 weeks
- irritable bowel syndrome

Ask a doctor or pharmacist before use if you are taking a prescription drug

When using this product you may have loose, watery, more frequent stools

Stop use and ask a doctor if

- you have rectal bleeding or your nausea, bloating, cramping or abdominal pain gets worse. These may be signs of a serious condition.
- you get diarrhea
- you need to use a laxative for longer than 1 week



: 306E1 2H B3

Drug Facts (continued)

If pregnant or breast-feeding, ask a health professional before use.

Keep out of reach of children. In case of overdose, get medical help or contact a Poison Control Center right away. (1-800-222-1222)

Directions

do not take more than directed unless advised by your doctor

the bottle top is a measuring cap marked to contain 17 grams of powder when filled to the indicated line (white section in cap)

adults and children 17 years of age and older:

- fill to top of white section in cap which is marked to indicate the correct dose (17 g)
- stir and dissolve in any 4 to 8 ounces of beverage (cold, hot or room temperature) then drink
- use once a day
- use no more than 7 days
- children 16 years of age or under: ask a doctor

Other information

- store at 20°-25°C (68°-77°F)
- tamper-evident: do not use if printed foil seal under cap is missing, open or broken

Inactive ingredients none

Questions or comments? 1-800-719-9260

• Sugar Free

*This product is not manufactured or distributed by MSD Consumer Care, Inc., distributor of MiraLAX®.

DISTRIBUTED BY HY-VEE, INC. WEST DES MOINES, IA 50266

SATISFACTION GUARANTEED CALL 1-800-289-8343 www.hy-vee.com

CLEAR LAX ORIGINAL PRESCRIPTION STRENGTH

polyethylene glycol 3350 powder, for solution

Product Information

Product Type	HUMAN OTC DRUG LABEL	Item Code (Source)	NDC:42507-306
Route of Administration	ORAL	DEA Schedule	

Active Ingredient/Active Moiety

Ingredient Name	Basis of Strength	Strength
POLYETHYLENE GLYCOL 3350 (POLYETHYLENE GLYCOL 3350)	POLYETHYLENE GLYCOL 3350	17 g in 17 g

Product Characteristics

Color	WHITE	Score	
Shape		Size	
Flavor		Imprint Code	
Contains			

Packaging

#	Item Code	Package Description	Marketing Start Date	Marketing End Date
1	NDC:42507-306-02	238 g in 1 BOTTLE		

2	NDC:42507-306-03	510 g in 1 BOTTLE		
3	NDC:42507-306-19	595 g in 1 BOTTLE		

Marketing Information

Marketing Category	Application Number or Monograph Citation	Marketing Start Date	Marketing End Date
ANDA	ANDA090685	10/11/2009	

Labeler - HyVee Inc (006925671)

Revised: 8/2014

HyVee Inc