

SOUND BODY DAYTIME COLD AND FLU RELIEF NIGHTTIME COLD AND FLU RELIEF- acetaminophen, dextromethorphan hbr, doxylamine succinate, guaifenesin, phenylephrine hcl
Big Lots Stores, Inc.

Disclaimer: Most OTC drugs are not reviewed and approved by FDA, however they may be marketed if they comply with applicable regulations and policies. FDA has not evaluated whether this product complies.

Big Lots, Inc. DayTime Cold & Flu NightTime Cold & Flu Drug Facts

Nighttime Severe Cold & Flu
Active ingredients (in each 30 mL)

Acetaminophen 650 mg

Dextromethorphan HBr 20 mg

Doxylamine succinate 12.5 mg

Phenylephrine HCl 10 mg

Purpose

Pain reliever/fever reducer

Cough suppressant

Antihistamine

Nasal decongestant

Uses

temporarily relieves common cold/flu symptoms:

- nasal congestion
- sinus congestion and pressure
- minor aches and pains
- headache
- fever
- sore throat
- runny nose and sneezing
- cough due to minor throat and bronchial irritation
- cough to help you sleep
- reduces swelling of nasal passages
- promotes nasal and/or sinus drainage
- temporarily restores freer breathing through the nose

Warnings

Liver warning: This product contains acetaminophen. Severe liver damage may occur if you take

- more than 4,000 mg of acetaminophen in 24 hours
- with other drugs containing acetaminophen
- 3 or more alcoholic drinks every day while using this product

Allergy alert: Acetaminophen may cause severe skin reactions. Symptoms may include:

- skin reddening
- blisters
- rash

If a skin reaction occurs, stop use and seek medical help right away.

Sore throat warning: If sore throat is severe, persists for more than 2 days, is accompanied or followed by fever, headache, rash, nausea, or vomiting, consult a doctor promptly.

Do not use

- with any other drug containing acetaminophen (prescription or nonprescription). If you are not sure whether a drug contains acetaminophen, ask a doctor or pharmacist.
- if you are now taking a prescription monoamine oxidase inhibitor (MAOI) (certain drugs for depression, psychiatric, or emotional conditions, or Parkinson's disease), or for 2 weeks after stopping the MAOI drug. If you do not know if your prescription drug contains an MAOI, ask a doctor or pharmacist before taking this product.
- if you have ever had an allergic reaction to this product or any of its ingredients

Ask a doctor before use if you have

- liver disease
- heart disease
- high blood pressure
- thyroid disease
- diabetes
- glaucoma
- cough that occurs with too much phlegm (mucus)
- a breathing problem such as emphysema or chronic bronchitis
- trouble urinating due to an enlarged prostate gland
- persistent or chronic cough as occurs with smoking, asthma, or emphysema
- a sodium-restricted diet

Ask a doctor or pharmacist before use if you are

- taking sedatives or tranquilizers
- taking the blood thinning drug warfarin

When using this product

- **do not use more than directed**
- excitability may occur, especially in children
- marked drowsiness may occur
- avoid alcoholic drinks
- be careful when driving a motor vehicle or operating machinery
- alcohol, sedatives, and tranquilizers may increase drowsiness

Stop use and ask a doctor if

- you get nervous, dizzy or sleepless
- pain, nasal congestion, or cough gets worse or lasts more than 7 days
- fever gets worse or lasts more than 3 days
- redness or swelling is present
- new symptoms occur
- cough comes back or occurs with rash or headache that lasts. These could be signs of a serious condition.

If pregnant or breast-feeding,

ask a health professional before use.

Keep out of reach of children.

Overdose warning: In case of overdose, get medical help or contact a Poison Control Center right away (1-800-222-1222). Quick medical attention is critical for adults as well as for children even if you do not notice any signs or symptoms.

Directions

- take only as directed – see Overdose warning
- only use the dose cup provided
- do not exceed 4 doses per 24 hrs

adults & children 12 yrs & over	30 mL every 4 hrs
children 4 to under 12 yrs	ask a doctor
children under 4 yrs	do not use

Other information

- **each 30 mL contains:** sodium 44 mg
- store at 20-25°C (68-77°F)

Inactive ingredients

alcohol, anhydrous citric acid, D&C yellow #10, edetate disodium, FD&C green #3, FD&C

yellow #6, flavor, glycerin, propylene glycol, purified water, saccharin sodium, sodium benzoate, sodium chloride, sodium citrate, sorbitol solution, sucralose

Questions or comments?

1-800-719-9260

Daytime Severe Cold & Flu

Active ingredients - (in each 15 mL)

Acetaminophen 325 mg

Dextromethorphan HBr 10 mg

Guaifenesin 200 mg

Phenylephrine HCl 5 mg

Purpose

Pain reliever/fever reducer

Cough suppressant

Expectorant

Nasal decongestant

Uses

temporarily relieves common cold/flu symptoms:

- nasal congestion
- sinus congestion and pressure
- cough due to minor throat and bronchial irritation
- minor aches and pains
- headache
- fever
- sore throat
- reduces swelling of nasal passages
- temporarily restores freer breathing through the nose
- promotes nasal and/or sinus drainage
- helps loosen phlegm (mucus) and thin bronchial secretions to rid the bronchial passageways of bothersome mucus and make coughs more productive

Warnings

Liver warning: This product contains acetaminophen. Severe liver damage may occur if

- adult takes more than 4,000 mg of acetaminophen in 24 hours
- child takes more than 5 doses in 24 hours

- taken with other drugs containing acetaminophen
- adult has 3 or more alcoholic drinks every day while using this product

Allergy alert: Acetaminophen may cause severe skin reactions. Symptoms may include:

- skin reddening
- blisters
- rash

If a skin reaction occurs, stop use and seek medical help right away.

Sore throat warning: If sore throat is severe, persists for more than 2 days, is accompanied or followed by fever, headache, rash, nausea, or vomiting, consult a doctor promptly.

Do not use

- with any other drug containing acetaminophen (prescription or nonprescription). If you are not sure whether a drug contains acetaminophen, ask a doctor or pharmacist.
- if you are now taking a prescription monoamine oxidase inhibitor (MAOI) (certain drugs for depression, psychiatric, or emotional conditions, or Parkinson's disease), or for 2 weeks after stopping the MAOI drug. If you do not know if your prescription drug contains an MAOI, ask a doctor or pharmacist before taking this product.
- if you have ever had an allergic reaction to this product or any of its ingredients

Ask a doctor before use if you have

- liver disease
- heart disease
- thyroid disease
- diabetes
- high blood pressure
- trouble urinating due to an enlarged prostate gland
- cough that occurs with too much phlegm (mucus)
- persistent or chronic cough such as occurs with smoking, asthma, chronic bronchitis, or emphysema

Ask a doctor or pharmacist before use if you are

taking the blood thinning drug warfarin

When using this product

do not use more than directed

Stop use and ask a doctor if

- you get nervous, dizzy or sleepless
- pain, nasal congestion or cough gets worse or lasts more than 5 days (children) or

7 days (adults)

- fever gets worse or lasts more than 3 days
- redness or swelling is present
- new symptoms occur
- cough comes back or occurs with rash or headache that lasts. These could be signs of a serious condition.

If pregnant or breast-feeding,

ask a health professional before use.

Keep out of reach of children.

Overdose warning: In case of overdose, get medical help or contact a Poison Control Center right away (1-800-222-1222). Quick medical attention is critical for adults as well as for children even if you do not notice any signs or symptoms.

Directions

- take only as directed – see Overdose warning
- only use the dose cup provided
- do not exceed 4 doses per 24 hrs

adults & children 12 yrs & over	30 mL every 4 hrs
children 6 to under 12 yrs	15 mL every 4 hrs
children 4 to under 6 yrs	ask a doctor
children under 4 yrs	do not use

Other information

- **each 15 mL contains:** sodium 6 mg
- store at 20-25°C (68-77°F). Do not refrigerate.

Inactive ingredients

butylated hydroxyanisole, edetate disodium, FD&C yellow #6, flavor, glycerin, menthol, monobasic sodium phosphate, polyethylene glycol, propylene glycol, purified water, saccharin sodium, sucrose, xanthan gum

Questions or comments?

1-800-719-9260

Package/Label Principal Display Panel

COMBO PACK

SOUND BODY™

Compare to the Active Ingredients in Vicks® DayQuil® Severe Cold & Flu
Non-Drowsy

DAYTIME

Cold & Flu Relief Severe

Pain Reliever / Fever Reducer – Acetaminophen

Cough Suppressant – Dextromethorphan HBr

Expectorant – Guaifenesin

Nasal Decongestant – Phenylephrine HCl

ALCOHOL FREE

MAX STRENGTH

12 FL OZ (355 mL)

SOUND BODY™

Compare to the Active Ingredients in Vicks® NyQuil® Severe Cold & Flu
NIGHTTIME

Cold & Flu Relief Severe

Pain Reliever / Fever Reducer – Acetaminophen

Cough Suppressant – Dextromethorphan HBr

Antihistamine – Doxylamine Succinate

Nasal Decongestant – Phenylephrine HCl

10% ALCOHOL

MAX STRENGTH

12 FL OZ (355 mL)

COMBO PACK



Compare to the Active
Ingredients in Vicks®
DayQuil® Severe Cold & Flu*

Compare to the Active
Ingredients in Vicks®
NyQuil® Severe Cold & Flu*

Non-Drowsy
DAYTIME

**Cold & Flu
Relief Severe**

Pain Reliever / Fever Reducer -
Acetaminophen
Cough Suppressant -
Dextromethorphan HBr
Expectorant - **Guafenesin**
Nasal Decongestant -
Phenylephrine HCl

ALCOHOL FREE

MAX STRENGTH

12 FL OZ
(355 mL)

NIGHTTIME

**Cold & Flu
Relief Severe**

Pain Reliever / Fever Reducer -
Acetaminophen
Cough Suppressant -
Dextromethorphan HBr
Antihistamine -
Doxylamine Succinate
Nasal Decongestant -
Phenylephrine HCl

10% ALCOHOL

MAX STRENGTH

12 FL OZ
(355 mL)

DO NOT USE IF PRINTED NECKBAND IS
BROKEN OR MISSING

PARENTS:

Learn about teen medicine abuse


www.StopMedicineAbuse.org

GLUTEN FREE

*These products are not manufactured or distributed by
Procter & Gamble, distributor of Vicks® DayQuil® Severe
Cold & Flu and Vicks® NyQuil® Severe Cold & Flu.

Packaged by Perrigo Company Inc.
Big Lots Stores, Inc.
P.O. Box 28522
Columbus, OH 43228-0522
VW5003185 ITEM#0002





Nighttime Severe Cold & Flu

Drug Facts	
Active ingredients (in each 30 mL)	Purpose
Acetaminophen 650 mg	Pain reliever/fever reducer
Pseudoephedrine HBr 20 mg	Cough suppressant
Dextromethorphan HBr 12.5 mg	Expectorant
Phenylephrine HCl 10 mg	Nasal decongestant
Uses	
Temporarily relieves common cold/flu symptoms:	
■ nasal congestion	■ sinus congestion and pressure
■ minor aches and pains	■ headache
■ fever	■ sore throat
■ runny nose and sneezing	■ cough due to minor throat and bronchial irritation
■ cough to help you sleep	■ reduces swelling of nasal passages
■ promotes nasal and/or sinus drainage	■ temporarily restores free breathing through the nose
Warnings	
Use warning: This product contains acetaminophen. Severe liver damage may occur if you take:	
■ more than 4,000 mg of acetaminophen in 24 hours	
■ with other drugs containing acetaminophen	
■ 3 or more alcoholic drinks every day while using this product	
Allergy alert: Acetaminophen may cause severe skin reactions. Symptoms may include:	
■ skin redness	
■ blisters	
■ rash	
If a skin reaction occurs, stop use and seek medical help right away.	
Sore throat warning: If sore throat is severe, persists for more than 2 days, is accompanied or followed by fever, headache, rash, or swelling, consult a doctor promptly.	
Do not use	
■ with any other drug containing acetaminophen (prescription or nonprescription), if you are not sure whether a drug contains acetaminophen, ask a doctor or pharmacist.	
■ if you are now taking a prescription monoamine oxidase inhibitor (MAOI) (certain drugs for depression, psychiatric, or emotional conditions, or Parkinson's disease), or for 2 weeks after stopping the MAOI drug. If you do not know if your prescription drug contains an MAOI, ask a doctor or pharmacist before taking this product.	
■ if you have ever had an allergic reaction to this product or any of its ingredients	
Ask a doctor before use if you have	
■ liver disease	
■ heart disease	
■ high blood pressure	
■ thyroid disease	
■ diabetes	
■ glaucoma	
■ cough that occurs with too much phlegm (mucus)	
■ a breathing problem such as emphysema or chronic bronchitis	
■ trouble urinating due to an enlarged prostate gland	
■ persistent or chronic cough such as occurs with smoking, asthma, or emphysema	
■ a sodium-restricted diet	
Ask a doctor or pharmacist before use if you are	
■ taking sedatives or tranquilizers	
■ taking the blood thinning drug warfarin	
When using this product	
■ do not use more than directed	
■ marked drowsiness may occur	
■ avoid alcoholic drinks	
■ be careful when driving a motor vehicle or operating machinery	
■ alcohol, sedatives, and tranquilizers may increase drowsiness	
Stop use and ask a doctor if	
■ you get nervous, dizzy or dizzy	
■ pain, nasal congestion, or cough gets worse or lasts more than 7 days	
■ fever gets worse or lasts more than 3 days	
■ redness or swelling is present	
■ new symptoms occur	
■ cough comes back or occurs with rash or headache that lasts.	
These could be signs of a serious condition.	
If pregnant or breast-feeding, ask a health professional before use.	
Keep out of reach of children. Overdose warning: In case of overdose, get medical help or contact a Poison Control Center right away (1-800-222-1222). Quick medical attention is critical for adults as well as for children even if you do not notice any signs or symptoms.	
Directions	
■ take only as directed - see Overdose warning	
■ only use the dose cup provided	
■ do not exceed 4 doses per 24 hrs	
adults & children 12 yrs & over	30 mL every 4 hrs
children 4 to under 12 yrs	ask a doctor
children under 4 yrs	do not use
Other information	
■ each 30 mL contains: sodium 44 mg	
■ store at 20-25°C (68-77°F)	
Inactive ingredients	
alcohol, anhydrous citric acid, D&C yellow #10, edetate disodium, FD&C green #3, FD&C yellow #6, flavors, glycerin, propylene glycol, purified water, saccharin sodium, sodium benzoate, sodium chloride, sodium citrate, sorbitol solution, sucralose	
Questions or comments? 1-800-718-0200	

Daytime Severe Cold & Flu

Drug Facts	
Active ingredients (in each 15 mL)	Purpose
Acetaminophen 325 mg	Pain reliever/fever reducer
Dextromethorphan HBr 10 mg	Cough suppressant
Guaifenesin 200 mg	Expectorant
Phenylephrine HCl 5 mg	Nasal decongestant
Uses	
Temporarily relieves common cold/flu symptoms:	
■ nasal congestion	■ sinus congestion and pressure
■ cough due to minor throat and bronchial irritation	■ headache
■ minor aches and pains	■ fever
■ sore throat	■ reduces swelling of nasal passages
Temporarily restores free breathing through the nose	
■ promotes nasal and/or sinus drainage	
■ helps loosen phlegm (mucus) and thin bronchial secretions to aid the bronchial passages way of to the nose, mucus and make coughs more productive	
Warnings	
Use warning: This product contains acetaminophen. Severe liver damage may occur if:	
■ adult takes more than 4,000 mg of acetaminophen in 24 hours	
■ child takes more than 5 doses in 24 hours	
■ takes with other drugs containing acetaminophen	
■ adult has 3 or more alcoholic drinks every day while using this product	
Allergy alert: Acetaminophen may cause severe skin reactions. Symptoms may include:	
■ skin redness	
■ blisters	
■ rash	
If a skin reaction occurs, stop use and seek medical help right away.	
Sore throat warning: If sore throat is severe, persists for more than 2 days, is accompanied or followed by fever, headache, rash, or swelling, consult a doctor promptly.	
Do not use	
■ with any other drug containing acetaminophen (prescription or nonprescription), if you are not sure whether a drug contains acetaminophen, ask a doctor or pharmacist.	
■ if you are now taking a prescription monoamine oxidase inhibitor (MAOI) (certain drugs for depression, psychiatric, or emotional conditions, or Parkinson's disease), or for 2 weeks after stopping the MAOI drug. If you do not know if your prescription drug contains an MAOI, ask a doctor or pharmacist before taking this product.	
■ if you have ever had an allergic reaction to this product or any of its ingredients	
Ask a doctor before use if you have	
■ liver disease	
■ heart disease	
■ thyroid disease	
■ diabetes	
■ high blood pressure	
■ trouble urinating due to an enlarged prostate gland	
■ cough that occurs with too much phlegm (mucus)	
■ persistent or chronic cough such as occurs with smoking, asthma, chronic bronchitis, or emphysema	
Ask a doctor or pharmacist before use if you are taking the blood thinning drug warfarin	
When using this product	
do not use more than directed	
Stop use and ask a doctor if	
■ you get nervous, dizzy or dizzy	
■ pain, nasal congestion or cough gets worse or lasts more than 5 days (children) or 7 days (adults)	
■ fever gets worse or lasts more than 3 days	
■ redness or swelling is present	
■ new symptoms occur	
■ cough comes back or occurs with rash or headache that lasts.	
These could be signs of a serious condition.	
If pregnant or breast-feeding, ask a health professional before use.	
Keep out of reach of children. Overdose warning: In case of overdose, get medical help or contact a Poison Control Center right away (1-800-222-1222). Quick medical attention is critical for adults as well as for children even if you do not notice any signs or symptoms.	
Directions	
■ take only as directed - see Overdose warning	
■ only use the dose cup provided	
■ do not exceed 4 doses per 24 hrs	
adults & children 12 yrs & over	20 mL every 4 hrs
children 6 to under 12 yrs	15 mL every 4 hrs
children 4 to under 6 yrs	ask a doctor
children under 4 yrs	do not use
Other information	
■ each 15 mL contains: sodium 6 mg	
■ store at 20-25°C (68-77°F). Do not refrigerate.	
Inactive ingredients	
butylated hydroxyanisole, edetate disodium, FD&C yellow #6, flavors, glycerin, menthol, monobasic sodium phosphate, polyethylene glycol, propylene glycol, purified water, saccharin sodium, sucrose, xanthan gum	
Questions or comments? 1-800-718-0200	

SOUND BODY DAYTIME COLD AND FLU RELIEF NIGHTTIME COLD AND FLU RELIEF

acetaminophen, dextromethorphan hbr, doxylamine succinate, guaifenesin, phenylephrine hcl kit

Product Information

Product Type	HUMAN OTC DRUG	Item Code (Source)	NDC:50594-200
--------------	----------------	--------------------	---------------

Packaging

#	Item Code	Package Description	Marketing Start Date	Marketing End Date
1	NDC:50594-200-02	1 in 1 CARTON; Type 0: Not a Combination Product	07/08/2021	

Quantity of Parts

Part #	Package Quantity	Total Product Quantity
Part 1	1 BOTTLE	355 mL
Part 2	1 BOTTLE	355 mL

Part 1 of 2

SOUND BODY COLD AND FLU RELIEF

acetaminophen, dextromethorphan hbr, doxylamine succinate, phenylephrine hcl solution

Product Information

Item Code (Source)	NDC:50594-123
--------------------	---------------

Route of Administration	ORAL
-------------------------	------

Active Ingredient/Active Moiety

Ingredient Name	Basis of Strength	Strength
ACETAMINOPHEN (UNII: 362O9ITL9D) (ACETAMINOPHEN - UNII:362O9ITL9D)	ACETAMINOPHEN	650 mg in 30 mL
DEXTROMETHORPHAN HYDROBROMIDE (UNII: 9D2RTI9KYH) (DEXTROMETHORPHAN - UNII:7355X3ROTS)	DEXTROMETHORPHAN HYDROBROMIDE	20 mg in 30 mL
DOXYLAMINE SUCCINATE (UNII: V9BI9B5YI2) (DOXYLAMINE - UNII:95QB77JKPL)	DOXYLAMINE SUCCINATE	12.5 mg in 30 mL
PHENYLEPHRINE HYDROCHLORIDE (UNII: 04JA59TNSJ) (PHENYLEPHRINE - UNII:1WS297W6MV)	PHENYLEPHRINE HYDROCHLORIDE	10 mg in 30 mL

Inactive Ingredients

Ingredient Name			Strength	
ALCOHOL (UNII: 3K9958V90M)				
ANHYDROUS CITRIC ACID (UNII: XF417D3PSL)				
D&C YELLOW NO. 10 (UNII: 35SW5USQ3G)				
EDETATE DISODIUM (UNII: 7FLD91C86K)				
FD&C GREEN NO. 3 (UNII: 3P3ONR6O1S)				
FD&C YELLOW NO. 6 (UNII: H77VEI93A8)				
GLYCERIN (UNII: PDC6A3C0OX)				
PROPYLENE GLYCOL (UNII: 6DC9Q167V3)				
WATER (UNII: 059QF0KO0R)				
SACCHARIN SODIUM (UNII: SB8ZUX40TY)				
SODIUM BENZOATE (UNII: OJ245FE5EU)				
SODIUM CHLORIDE (UNII: 451W47IQ8X)				
SODIUM CITRATE, UNSPECIFIED FORM (UNII: 1Q73Q2JULR)				
SORBITOL (UNII: 506T60A25R)				
SUCRALOSE (UNII: 96K6UQ3ZD4)				
Product Characteristics				
Color	GREEN	Score		
Shape		Size		
Flavor	MINT	Imprint Code		
Contains				
Packaging				
#	Item Code	Package Description	Marketing Start Date	Marketing End Date
1	NDC:50594-123-40	355 mL in 1 BOTTLE; Type 0: Not a Combination Product		
Marketing Information				
Marketing Category		Application Number or Monograph Citation	Marketing Start Date	Marketing End Date
OTC monograph final		part341		
Part 2 of 2				
SOUND BODY COLD AND FLU RELIEF				
acetaminophen, dextromethorphan hbr, guaifenesin, phenylephrine hcl solution				
Product Information				
Item Code (Source)		NDC:50594-016		
Route of Administration		ORAL		

Active Ingredient/Active Moiety		
Ingredient Name	Basis of Strength	Strength
ACETAMINOPHEN (UNII: 362O9ITL9D) (ACETAMINOPHEN - UNII:362O9ITL9D)	ACETAMINOPHEN	325 mg in 15 mL
DEXTROMETHORPHAN HYDROBROMIDE (UNII: 9D2RTI9KYH) (DEXTROMETHORPHAN - UNII:7355X3ROTS)	DEXTROMETHORPHAN HYDROBROMIDE	10 mg in 15 mL
GUAIFENESIN (UNII: 495W7451VQ) (GUAIFENESIN - UNII:495W7451VQ)	GUAIFENESIN	200 mg in 15 mL
PHENYLEPHRINE HYDROCHLORIDE (UNII: 04JA59TNSJ) (PHENYLEPHRINE - UNII:1WS297W6MV)	PHENYLEPHRINE HYDROCHLORIDE	5 mg in 15 mL

Inactive Ingredients	
Ingredient Name	Strength
BUTYLATED HYDROXYANISOLE (UNII: REK4960K2U)	
EDETATE DISODIUM (UNII: 7FLD91C86K)	
FD&C YELLOW NO. 6 (UNII: H77VEI93A8)	
GLYCERIN (UNII: PDC6A3C0OX)	
MENTHOL, UNSPECIFIED FORM (UNII: L7T10EIP3A)	
SODIUM PHOSPHATE, MONOBASIC, UNSPECIFIED FORM (UNII: 3980JIH2SW)	
POLYETHYLENE GLYCOL, UNSPECIFIED (UNII: 3WJQ0SDW1A)	
PROPYLENE GLYCOL (UNII: 6DC9Q167V3)	
WATER (UNII: 059QF0KO0R)	
SACCHARIN SODIUM (UNII: SB8ZUX40TY)	
SUCROSE (UNII: C151H8M554)	
XANTHAN GUM (UNII: TTV12P4NEE)	

Product Characteristics			
Color	ORANGE (clear)	Score	
Shape		Size	
Flavor	FRUIT, MENTHOL	Imprint Code	
Contains			

Packaging				
#	Item Code	Package Description	Marketing Start Date	Marketing End Date
1	NDC:50594-016-40	355 mL in 1 BOTTLE; Type 0: Not a Combination Product		

Marketing Information			
Marketing Category	Application Number or Monograph Citation	Marketing Start Date	Marketing End Date
OTC monograph final	part341		

Marketing Information

Marketing Category	Application Number or Monograph Citation	Marketing Start Date	Marketing End Date
OTC monograph final	part341	07/08/2021	

Labeler - Big Lots Stores, Inc. (017885351)

Revised: 11/2022

Big Lots Stores, Inc.