

MAGNESIUM SULFATE- magnesium sulfate heptahydrate injection, solution

Cardinal Health

Disclaimer: This drug has not been found by FDA to be safe and effective, and this labeling has not been approved by FDA. For further information about unapproved drugs, [click here](#).

Magnesium Sulfate Injection, USP

50 %

DESCRIPTION

Magnesium Sulfate Injection, USP is a sterile, concentrated solution of magnesium sulfate heptahydrate for use as an electrolyte replenisher or anticonvulsant by intramuscular (IM) or intravenous (IV) routes of administration. Must be diluted before IV use. The product is available in a 50% concentration.

Each mL contains: Magnesium sulfate heptahydrate 500 mg; Water for Injection q.s. Sulfuric acid and/or sodium hydroxide may have been added for pH adjustment (5.5-7.0). Osmolarity: 4060 mOsmol/L (calc.); 2.03 mM/mL magnesium sulfate anhydrous; 4.06 mEq/mL magnesium sulfate anhydrous. **This product can be used for IM or IV use.**

The solution contains no bacteriostat, antimicrobial agent or added buffer (except for pH adjustment) and is intended only for use as a single dose injection. When smaller doses are required the unused portion should be discarded with the entire unit.

Magnesium sulfate heptahydrate is chemically designated $\text{MgSO}_4 \cdot 7\text{H}_2\text{O}$, with a molecular weight of 246.47 and occurs as colorless crystals or white powder freely soluble in water.

CLINICAL PHARMACOLOGY

Magnesium is an important cofactor for enzymatic reactions and plays an important role in neurochemical transmission and muscular excitability.

As a nutritional adjunct in hyperalimentation, the precise mechanism of action for magnesium is uncertain. Early symptoms of hypomagnesemia (less than 1.5 mEq/L) may develop as early as three to four days or within weeks.

Predominant deficiency effects are neurological, e.g., muscle irritability, clonic twitching and tremors.

Hypocalcemia and hypokalemia often follow low serum levels of magnesium. While there are large stores of magnesium present intracellularly and in the bones of adults, these stores often are not mobilized sufficiently to maintain plasma levels. Parenteral magnesium therapy repairs the plasma deficit and causes deficiency symptoms and signs to cease.

Magnesium prevents or controls convulsions by blocking neuromuscular transmission and decreasing the amount of acetylcholine liberated at the end-plate by the motor nerve impulse. Magnesium is said to have a depressant effect on the central nervous system (CNS), but it does not adversely affect the mother, fetus or neonate when used as directed in eclampsia or pre-eclampsia. Normal plasma magnesium levels range from 1.5 to 2.5 mEq/L.

As plasma magnesium rises above 4 mEq/L, the deep tendon reflexes are first decreased and then disappear as the plasma level approaches 10 mEq/L. At this level respiratory paralysis may occur.

Heart block also may occur at this or lower plasma levels of magnesium. Serum magnesium concentrations in excess of 12 mEq/L may be fatal.

Magnesium acts peripherally to produce vasodilation. With low doses only flushing and sweating occur, but larger doses cause lowering of blood pressure. The central and peripheral effects of magnesium poisoning are antagonized to some extent by IV administration of calcium.

Pharmacokinetics

With IV administration the onset of anticonvulsant action is immediate and lasts about 30 minutes.

Following IM administration, the onset of action occurs in about one hour and persists for three to four hours. Effective anticonvulsant serum levels range from 2.5 to 7.5 mEq/L. Magnesium is excreted solely by the kidney at a rate proportional to the plasma concentration and glomerular filtration.

INDICATIONS AND USAGE

Magnesium Sulfate Injection, USP is suitable for replacement therapy in magnesium deficiency, especially in acute hypomagnesemia accompanied by signs of tetany similar to those observed in hypocalcemia. In such cases, the serum magnesium level is usually below the lower limit of normal (1.5 to 2.5 mEq/L) and the serum calcium level is normal (4.3 to 5.3 mEq/L) or elevated.

In total parenteral nutrition (TPN), magnesium sulfate may be added to the nutrient admixture to correct or prevent hypomagnesemia which can arise during the course of therapy.

Magnesium Sulfate Injection is also indicated as a parenteral anticonvulsant for the prevention and control of seizures (convulsions) in severe toxemia of pregnancy. It effectively prevents and controls the convulsions of eclampsia without producing deleterious depression of the CNS of the mother or infant. However, other effective drugs are available for this purpose.

CONTRAINDICATIONS

Parenteral administration of the drug is contraindicated in patients with heart block or myocardial damage.

WARNINGS

WARNING: This product contains aluminum that may be toxic. Aluminum may reach toxic levels with prolonged parenteral administration if kidney function is impaired. Premature neonates are particularly at risk because their kidneys are immature, and they require large amounts of calcium and phosphate solutions, which contain aluminum.

Research indicates that patients with impaired kidney function, including premature neonates, who receive parenteral levels of aluminum at greater than 4 to 5 mcg/kg/day accumulate aluminum at levels associated with central nervous system and bone toxicity. Tissue loading may occur at even lower rates of administration.

IV use in eclampsia should be reserved for immediate control of life-threatening convulsions.

Parenteral use in the presence of renal insufficiency may lead to magnesium intoxication.

PRECAUTIONS

General

Administer with caution if flushing and sweating occurs. When barbiturates, narcotics or other hypnotics (or systemic anesthetics) are to be given in conjunction with magnesium, their dosage should

be adjusted with caution because of additive CNS depressant effects of magnesium.

Because magnesium is removed from the body solely by the kidneys, the drug should be used with caution in patients with renal impairment. Urine output should be maintained at a level of 100 mL or more during the four hours preceding each dose. Monitoring serum magnesium levels and the patient's clinical status is essential to avoid the consequences of overdosage in toxemia. Clinical indications of a safe dosage regimen include the presence of the patellar reflex (knee jerk) and absence of respiratory depression (approximately 16 breaths or more/min). When repeated doses of the drug are given parenterally, knee jerk reflexes should be tested before each dose and if they are absent, no additional magnesium should be given until they return. Serum magnesium levels usually sufficient to control convulsions range from 3 to 6 mg/100 mL (2.5 to 5 mEq/L). The strength of the deep tendon reflexes begins to diminish when magnesium levels exceed 4 mEq/L. Reflexes may be absent at 10 mEq magnesium/L, where respiratory paralysis is a potential hazard. An injectable calcium salt should be immediately available to counteract the potential hazards of magnesium intoxication in eclampsia.

Magnesium Sulfate Injection (50%) must be diluted to a concentration of 20% or less prior to IV infusion. Rate of administration should be slow and cautious, to avoid producing hypermagnesemia. The 50% solution also should be diluted to 20% or less for IM injection in infants and children.

Laboratory Tests

Magnesium Sulfate Injection should not be given unless hypomagnesemia has been confirmed and the serum concentration of magnesium is monitored. The normal serum level is 1.5 to 2.5 mEq/L.

Drug Interactions

CNS Depressants—When barbiturates, narcotics or other hypnotics (or systemic anesthetics), or other CNS depressants are to be given in conjunction with magnesium, their dosage should be adjusted with caution because of additive CNS depressant effects of magnesium. CNS depression and peripheral transmission defects produced by magnesium may be antagonized by calcium.

Neuromuscular Blocking Agents—Excessive neuromuscular block has occurred in patients receiving parenteral magnesium sulfate and a neuromuscular blocking agent; these drugs should be administered concomitantly with caution.

Cardiac Glycosides—Magnesium sulfate should be administered with extreme caution in digitalized patients, because serious changes in cardiac conduction which can result in heart block may occur if administration of calcium is required to treat magnesium toxicity.

Pregnancy Category A

Studies in pregnant women have not shown that Magnesium Sulfate Injection increases the risk of fetal abnormalities if administered during all trimesters of pregnancy. If this drug is used during pregnancy, the possibility of fetal harm appears remote. However, because studies cannot rule out the possibility of harm, Magnesium Sulfate Injection should be used during pregnancy only if clearly needed.

When administered by continuous IV infusion (especially for more than 24 hours preceding delivery) to control convulsions in toxemic mothers, the newborn may show signs of magnesium toxicity, including neuromuscular or respiratory depression (see **OVERDOSAGE**).

Nursing Mothers

Since magnesium is distributed into milk during parenteral magnesium sulfate administration, the drug should be used with caution in nursing women.

Geriatrics

Geriatric patients often require reduced dosage because of impaired renal function. In patients with severe impairment, dosage should not exceed 20 g in 48 hours. Serum magnesium should be monitored in such patients.

ADVERSE REACTIONS

The adverse effects of parenterally administered magnesium usually are the result of magnesium intoxication. These include flushing, sweating, hypotension, depressed reflexes, flaccid paralysis, hypothermia, circulatory collapse, cardiac and CNS depression proceeding to respiratory paralysis. Hypocalcemia with signs of tetany secondary to magnesium sulfate therapy for eclampsia has been reported.

OVERDOSAGE

Magnesium intoxication is manifested by a sharp drop in blood pressure and respiratory paralysis. Disappearance of the patellar reflex is a useful clinical sign to detect the onset of magnesium intoxication. In the event of overdosage, artificial ventilation must be provided until a calcium salt can be injected IV to antagonize the effects of magnesium.

For Treatment of Overdose

Artificial respiration is often required. Intravenous calcium, 10 to 20 mL of a 5% solution (diluted if desirable with isotonic sodium chloride for injection) is used to counteract effects of hypermagnesemia. Subcutaneous physostigmine, 0.5 to 1 mg may be helpful.

Hypermagnesemia in the newborn may require resuscitation and assisted ventilation via endotracheal intubation or intermittent positive pressure ventilation as well as IV calcium.

DOSAGE AND ADMINISTRATION

Dosage of magnesium sulfate must be carefully adjusted according to individual requirements and response, and administration of the drug should be discontinued as soon as the desired effect is obtained.

Both IV and IM administration are appropriate. IM administration of the undiluted 50% solution results in therapeutic plasma levels in 60 minutes, whereas IV doses will provide a therapeutic level almost immediately. The rate of IV injection should generally not exceed 150 mg/minute, except in severe eclampsia with seizures (see below).

Solutions for IV infusion must be diluted to a concentration of 20% or less prior to administration. The diluents commonly used are 5% Dextrose Injection, USP and 0.9% Sodium Chloride Injection, USP.

Deep IM injection of the undiluted (50%) solution is appropriate for adults, but the solution should be diluted to a 20% or less concentration prior to such injection in children.

In Magnesium Deficiency

In the treatment of mild magnesium deficiency, the usual adult dose is 1 g, equivalent to 8.12 mEq of magnesium (2 mL of the 50% solution) injected IM every six hours for four doses (equivalent to a total of 32.5 mEq of magnesium per 24 hours). For severe hypomagnesemia, as much as 250 mg (approximately 2 mEq) per kg of body weight (0.5 mL of the 50% solution) may be given IM within a period of four hours if necessary. Alternatively, 5 g (approximately 40 mEq) can be added to one liter of 5% Dextrose Injection, USP or 0.9% Sodium Chloride Injection, USP for slow IV infusion over a three-hour period. In the treatment of deficiency states, caution must be observed to prevent exceeding the renal excretory capacity.

In Hyperalimentation

In TPN, maintenance requirements for magnesium are not precisely known. The maintenance dose used in adults ranges from 8 to 24 mEq (1 to 3 g) daily; for infants, the range is 2 to 10 mEq (0.25 to 1.25 g) daily.

In Eclampsia

In severe pre-eclampsia or eclampsia, the total initial dose is 10 to 14 g of magnesium sulfate. Intravenously, a dose of 4 to 5 g in 250 mL of 5% Dextrose Injection, USP or 0.9% Sodium Chloride Injection, USP may be infused. Simultaneously, IM doses of up to 10 g (5 g or 10 mL of the undiluted 50% solution in each buttock) are given. Alternatively, the initial IV dose of 4 g may be given by diluting the 50% solution to a 10 or 20% concentration; the diluted fluid (40 mL of a 10% solution or 20 mL of a 20% solution) may then be injected IV over a period of three to four minutes. Subsequently, 4 to 5 g (8 to 10 mL of the 50% solution) are injected IM into alternate buttocks every four hours as needed, depending on the continuing presence of the patellar reflex and adequate respiratory function. Alternatively, after the initial IV dose, some clinicians administer 1 to 2 g/hour by constant IV infusion. Therapy should continue until paroxysms cease. A serum magnesium level of 6 mg/100 mL is considered optimal for control of seizures. A total daily (24 hr) dose of 30 to 40 g should not be exceeded. In the presence of severe renal insufficiency, the maximum dosage of magnesium sulfate is 20 grams/48 hours and frequent serum magnesium concentrations must be obtained.

Other uses

In counteracting the muscle-stimulating effects of barium poisoning, the usual dose of magnesium sulfate is 1 to 2 g given IV.

For controlling seizures associated with epilepsy, glomerulonephritis or hypothyroidism, the usual adult dose is 1 g administered IM or IV.

In paroxysmal atrial tachycardia, magnesium should be used only if simpler measures have failed and there is no evidence of myocardial damage. The usual dose is 3 to 4 g (30 to 40 mL of a 10% solution) administered IV over 30 seconds with extreme caution.

For reduction of cerebral edema, 2.5 g (25 mL of a 10% solution) is given IV.

Incompatibilities

Magnesium sulfate in solution may result in a precipitate formation when mixed with solutions containing:

Alcohol (in high concentrations)	Heavy metals
Alkali carbonates and bicarbonates	Hydrocortisone sodium succinate
Alkali hydroxides	Phosphates
Arsenates	Polymyxin B sulfate
Barium	Procaine hydrochloride
Calcium	Salicylates
Clindamycin phosphate	Strontium
	Tartrates

The potential incompatibility will often be influenced by the changes in the concentration of reactants and the pH of the solutions.

It has been reported that magnesium may reduce the antibiotic activity of streptomycin, tetracycline and tobramycin when given together.

Parenteral drug products should be inspected visually for particulate matter and discoloration prior to administration, whenever solution and container permit.

HOW SUPPLIED

Product No.	NDC No.	% Magnesium Sulfate Heptahydrate	Volume Fill	Magnesium mg/mL	Sulfate mg/mL
6420	63323-064-20	50	20 mL	49.3	194.7
6450	63323-064-50	50	50 mL	49.3	194.7

Do not administer unless solution is clear and seal intact.

Preservative Free. Discard unused portion.

Store at 20° to 25°C (68° to 77°F) [see USP Controlled Room Temperature].

45808E

Revised: January 2008

Cardinal Health

Zanesville, OH 43701

IA66840313

PACKAGE LABEL - PRINCIPAL DISPLAY

Magnesium Sulfate Injection, USP

50% (10 grams/20mL)

1 x 20 mL Single Dose Vial



**MAGNESIUM SULFATE INJECTION, USP
50% (10 grams/20 mL)**

1 x 20 mL SINGLE DOSE VIAL

For IM or IV Use

MUST BE DILUTED BEFORE IV USE.

Sterile, Nonpyrogenic

Preservative Free

Each mL contains: 500 mg (4.06 mEq) of magnesium sulfate heptahydrate; Water for Injection q.s. Sulfuric acid and/or sodium hydroxide may have been added for pH adjustment.

4.06 mEq/mL 4.06 mOsmol/mL

Contains no more than 300 mcg/L of aluminum.

For magnesium replenishment: Usual dosage range
1-3 grams (8-24 mEq) per day.

See product insert for prescribing information, precautions, warnings and for additional information.

Use only if solution is clear and seal intact and undamaged. Cleanse stopper with antiseptic. The entire contents should be dispensed promptly under a laminar flow hood.

Discard unused portion.

RX ONLY

STORAGE: Store at 20 to 25 C (68 to 77 F) [see USP Controlled Room Temperature].

Bulk NDC 63323-064-20

APP Pharmaceuticals, LLC
Schaumburg, IL 60173

Repackaged by Cardinal Health
Zanesville, OH 43701
LA66840613

Cardinal Health Item # AR6684

LOT #: XXXX

EXP. DATE: 12/34

MAGNESIUM SULFATE

magnesium sulfate heptahydrate injection, solution

Product Information

Product Type	HUMAN PRESCRIPTION DRUG LABEL	Item Code (Source)	NDC:55154- 9394(NDC:63323-064)
Route of Administration	INTRAVENOUS, INTRAMUSCULAR	DEA Schedule	

Active Ingredient/Active Moiety				
Ingredient Name		Basis of Strength		Strength
MAGNESIUM SULFATE HEPTAHYDRATE (MAGNESIUM CATION)		MAGNESIUM SULFATE HEPTAHYDRATE		500 mg in 1 mL
Inactive Ingredients				
Ingredient Name			Strength	
WATER				
SULFURIC ACID				
SODIUM HYDRO XIDE				
Packaging				
#	Item Code	Package Description	Marketing Start Date	Marketing End Date
1	NDC:55154-9394-8	1 in 1 BAG		
1		20 mL in 1 VIAL, SINGLE-DOSE		
Marketing Information				
Marketing Category	Application Number or Monograph Citation		Marketing Start Date	Marketing End Date
Unapproved drug other			08/08/2000	

Labeler - Cardinal Health (188557102)

Establishment			
Name	Address	ID/FEI	Business Operations
Cardinal Health		188557102	REPACK(55154-9394)