

GOOD NEIGHBOR PHARMACY ANTI DIARRHEAL ANTI GAS- loperamide hydrochloride, simethicone tablet
Amerisource Bergen

Amerisource Bergen Anti-Diarrheal Anti-Gas Drug Facts

Active ingredients (in each caplet)

Loperamide HCl 2 mg

Simethicone 125 mg

Purposes

Anti-diarrheal

Anti-gas

Uses

relieves symptoms of diarrhea plus bloating, pressure and cramps, commonly referred to as gas

Warnings

Allergy alert: Do not use if you have ever had a rash or other allergic reaction to loperamide HCl

Heart alert: Taking more than directed can cause serious heart problems or death

Do not use

- if you have bloody or black stool
- if you have difficulty swallowing

Ask a doctor before use if you have

- fever
- mucus in the stool
- a history of liver disease
- a history of abnormal heart rhythm

Ask a doctor or pharmacist before use if you are

taking a prescription drug. Loperamide may interact with certain prescription drugs.

When using this product

tiredness, drowsiness or dizziness may occur. Be careful when driving or operating machinery.

Stop use and ask a doctor if

- symptoms get worse
- diarrhea lasts for more than 2 days
- you get abdominal swelling or bulging. These may be signs of a serious condition.
- you have difficulty swallowing the caplet

If pregnant or breast-feeding,

ask a health professional before use.

Keep out of reach of children.

In case of overdose, get medical help or contact a Poison Control Center right away. (1-800-222-1222)

Directions

- **drink plenty of clear fluids to help prevent dehydration caused by diarrhea**
- take only on an empty stomach (1 hour before or 2 hours after a meal)
- take with a full (8 oz.) glass of water
- find right dose on chart below. If possible, use weight to dose; otherwise, use age.

| | |
|---------------------------------------|---|
| adults and children 12 years and over | 2 caplets after the first loose stool; 1 caplet after each subsequent loose stool; but no more than 4 caplets in 24 hours |
| children 9-11 years (60-95 lbs) | 1 caplet after the first loose stool; ½ caplet after each subsequent loose stool; but no more than 3 caplets in 24 hours |
| children 6-8 years (48-59 lbs) | 1 caplet after the first loose stool; ½ caplet after each subsequent loose stool; but no more than 2 caplets in 24 hours |
| children 2-5 years (34 to 47 lbs) | ask a doctor |
| children under 2 years (up to 33 lbs) | do not use |

Other information

- **each caplet contains:** calcium 165 mg, sodium 3 mg
- store between 20°-25°C (68°-77°F). Protect from light.
- **do not use if printed blister unit is broken or torn**

Inactive ingredients

acesulfame potassium, croscarmellose sodium, dibasic calcium phosphate, maltodextrin, microcrystalline cellulose, natural and artificial flavor, propylene glycol, stearic acid

Questions or comments?

1-800-719-9260

Package/Label Principal Display Panel

Compare to Imodium® Multi-Symptom Relief

active ingredients

loperamide hydrochloride and simethicone tablets, 2 mg/125 mg

Anti-Diarrheal/Anti-Gas

Multi-Symptom Relief Caplet

Relieves symptoms of diarrhea plus

Cramps & Pressure - Bloating - Gas

actual size

24 Caplets*

*Capsule-Shaped Tablets



GOOD NEIGHBOR PHARMACY ANTI DIARRHEAL ANTI GAS

loperamide hydrochloride, simethicone tablet

Product Information

| | | | |
|--------------------------------|----------------|---------------------------|---------------|
| Product Type | HUMAN OTC DRUG | Item Code (Source) | NDC:46122-620 |
| Route of Administration | ORAL | | |

Active Ingredient/Active Moiety

| Ingredient Name | Basis of Strength | Strength |
|---|--------------------------|----------|
| LOPERAMIDE HYDROCHLORIDE (UNII: 77TI35393C) (LOPERAMIDE - UNII:6X9OC3H4II) | LOPERAMIDE HYDROCHLORIDE | 2 mg |
| DIMETHICONE (UNII: 92RU3N3Y1O) (DIMETHICONE - UNII:92RU3N3Y1O) | DIMETHICONE | 125 mg |

Inactive Ingredients

| Ingredient Name | Strength |
|---|----------|
| SILICON DIOXIDE (UNII: ETJ7Z6XBU4) | |
| ACESULFAME POTASSIUM (UNII: 23OV73Q5G9) | |
| CROSCARMELLOSE SODIUM (UNII: M28OL1HH48) | |
| ANHYDROUS DIBASIC CALCIUM PHOSPHATE (UNII: L11K75P92J) | |
| MALTODEXTRIN (UNII: 7CVR7L4A2D) | |
| MICROCRYSTALLINE CELLULOSE (UNII: OP1R32D61U) | |
| PROPYLENE GLYCOL (UNII: 6DC9Q167V3) | |
| STEARIC ACID (UNII: 4ELV7Z65AP) | |

Product Characteristics

| | | | |
|-----------------|-------|---------------------|------------|
| Color | WHITE | Score | 2 pieces |
| Shape | OVAL | Size | 17mm |
| Flavor | | Imprint Code | LOR1;2;125 |
| Contains | | | |

Packaging

| # | Item Code | Package Description | Marketing Start Date | Marketing End Date |
|---|------------------|--|----------------------|--------------------|
| 1 | NDC:46122-620-62 | 24 in 1 CARTON | 09/24/2019 | |
| 1 | | 1 in 1 BLISTER PACK; Type 0: Not a Combination Product | | |

Marketing Information

| Marketing Category | Application Number or Monograph Citation | Marketing Start Date | Marketing End Date |
|--------------------|--|----------------------|--------------------|
| ANDA | ANDA209837 | 09/24/2019 | |

