

**ICON 5000 II- iodine liquid  
IBA**

*Disclaimer: This drug has not been found by FDA to be safe and effective, and this labeling has not been approved by FDA. For further information about unapproved drugs, click here.*

-----

**ICON 5000 II - NPE FREE**

**ACTIVE INGREDIENTS:**

Iodine Complex:.....25.00%

(Provide a full .5% available iodine when used at the recommended dilution ratio)

**INERT INGREDIENTS:**

Triple Emollient Syste (As Humectants And Skin  
Conditioners):.....20.00%

(Provides a total of 2.0% emollients at the recommended dilution ratio)

Other  
Inerts.....55.00%

(Includes Buffering Agents, Stabilizing Agents, And Surfactants)

TOTAL:.....100.00%

**GENERAL RECOMMENDATIONS**

**GENERAL RECOMMENDATIONS**

1. Use teat dip applicators made of plastic or other non-corrosive materials. Do not use cups made of aluminum.
2. When soluiton in the teat dip cup becomes dirty and/or noticeably loses color. discard remaining contents and refill cup with fresh teat dip.
3. Do not use this product for cleaning or sanitizing equipment.
4. Wash teat dip cups after each milking.

**CAUTION**

**KEEP OUT OF REACH OF CHILDREN. AVOID CONTACT WITH FOOD. DO NOT TAKE INTERNALLY.**

**FIRST AID**

**FIRST AID:** IF TAKEN INTERNALLY, DRINK LARGE QUANTITIES OF WATER AND GET IMMEDIATE MEDICAL ATTENTION. IN CASE OF CONTACT WITH EYES, FLUSH LIBERALLY WITH TEPID WATER AND GET MEDICAL ATTENTION. NEVER GIVE ANYHTING BY MOUTH TO AN UNCONSCIOUS PERSON.

FOR MEDICAL EMERGENCIES CALL 1-303-623-5716

## **RECOMMENDED PROCEDURES FOR TEAT SANITATION**

## **RECOMMENDED PROCEDURES FOR TEAT SANITATION**

### **A. PRE-MILKING (PRE-DIPPING)**

1. Remove visible soils from the teats.
2. Onserve foremilk by stripping two or three streams of milk into a strip cup.
3. Dip or spray teh cow's teats full length with diluted **ICON 5000 II** Teat Dip.
4. Wait approximately 30 seconds.
5. Remove all teat dip by thoroughly drying the teats and teat orifices with a clean, single service paper towel(s).
6. Attach milking unit

### **B.POST -MILKING (POST-DIPPING)**

1. Immediately after milking, dip or spray all teats full length with diluted **ICON 5000 II** Teat Dip. **Ensure good coverage of all sides of the teats and teat orifices.**

Dilute 1 part ICON 5000 II TEAT DIP per 9 parts potable , tepid water.

### **WARNING**

**WARNING:** ICON 5000 II TEAT DIP CONCENTRATE MUST BE DILUTED BEFORE USE. DO NOT MIX WITH ANYTHING BUT POTABLE WATER. DO NOT RETURN TEAT DIP LEFT IN THE TEAT DIPPER TO STORAGE CONTAINER. ENSURE THAT DIPPED TEATS ARE DRY BEFORE EXPOSING THE ANIMALS TO WEATHER CONDITIONS WHICH MAY CAUSE DAMGE TO THE TEATS.

### **Icon 5000 II**

Iodine teat dip concentrate

NDC 29983-1018-8



A HIGHLY EFFECTIVE SANITIZING TEAT DIP



AN AID IN THE PREVENTION OF NEW INTRAMAMMARY INFECTIONS

**IODINE TEAT DIP CONCENTRATE**

Especially Formulated For Dairies Where **NPE FREE** Teat Dips Are Preferred Or Required

**ACTIVE INGREDIENT:**

Iodine Complex..... 25.00%  
 (Provides A Full .5% Available Iodine When Used At The Recommended Dilution Ratio)  
**INERT INGREDIENTS:**  
 Triple Emollients (As Humectants And Skin Conditioners)..... 20.00%  
 (Provides A Total Of 2.0% Emollients At The Recommended Dilution Ratio)  
 Other Inerts..... 55.00%  
 (Includes Buffering Agents, Stabilizing Agents And Surfactants)  
**TOTAL:**..... 100.00 %

When mixed according to label directions, this product provides highly effective levels of free iodine for fast kill of mastitis causing organisms.

**GENERAL RECOMMENDATIONS**

1. Use teat dip applicators made of plastic or other non-corrosive materials. Do not use cups made of aluminum.
2. When solution in the teat dip cup becomes dirty and/or noticeably loses color, discard remaining contents and refill cup with fresh teat dip.
3. Do not use this product for cleaning or sanitizing equipment.
4. Wash teat dip cups after each milking.

**FIRST AID PROCEDURES**

**EYE CONTACT:** IMMEDIATELY FLUSH EYE WITH LARGE AMOUNTS OF FRESH, TEPID WATER FOR 15 MINUTES. HOLD EYELID OPEN TO ENSURE COMPLETE IRRIGATION OF EYE AND LID TISSUES. TILT HEAD TO THE SIDE AND IRRIGATE THE EYE FROM THE BRIDGE OF THE NOSE TO THE OUTSIDE OF THE FACE.

**SKIN CONTACT:** FOR CONTACT WITH CONCENTRATED PRODUCTS, FLUSH THOROUGHLY WITH FRESH, TEPID WATER. DISCARD CONTAMINATED CLOTHING AND FOOTWEAR OR WASH BEFORE REUSE. FOR READY-TO-USE PRODUCTS, WASH ANY SENSITIVE AREAS.

**INGESTION:** DO NOT INDUCE VOMITING. NEVER GIVE ANYTHING BY MOUTH TO AN UNCONSCIOUS PERSON. DRINK LARGE AMOUNTS OF WATER. IF VOMITING OCCURS, KEEP AIRWAYS OPEN. KEEP HEAD LOWER THAN THE HIPS TO PREVENT ASPIRATION INTO THE LUNGS.

**ALWAYS SEEK IMMEDIATE MEDICAL ATTENTION FOR MEDICAL EMERGENCIES CALL 1-800-424-9300**

Dilute 1 Part  
**ICON 5000 II NPE FREE**  
**TEAT DIP**  
 Per 9 Parts Potable,  
 Tepid Water.

**RECOMMENDED PROCEDURES FOR TEAT SANITATION**

**A. PRE-MILKING (PRE-DIPPING)**

1. Remove visible soils from the teats.
2. Observe foremilk by stripping two or three streams of milk into a strip cup.
3. Dip, spray or foam the cow's teats full length with **DILUTED** **ICON 5000 II NPE FREE TEAT DIP**.
4. Wait approximately 30 seconds.
5. Remove all teat dip by thoroughly drying the teats and teat orifices with a clean, single service paper towel(s).
6. Attach milking unit.

**B. POST-MILKING (POST-DIPPING)**

1. Immediately after milking, dip or spray all teats full length with **DILUTED** **ICON 5000 II NPE FREE TEAT DIP**. Ensure good coverage of all sides of the teats and teat orifices.

**CAUTION: KEEP OUT OF REACH OF CHILDREN. AVOID CONTACT WITH FOOD. DO NOT TAKE INTERNALLY.**

**WARNING**

**ICON 5000 II NPE FREE TEAT DIP CONCENTRATE MUST BE DILUTED** BEFORE USE. DO NOT MIX WITH ANYTHING BUT POTABLE WATER.

PROTECT THIS PRODUCT FROM FREEZING. DO NOT RETURN DILUTED TEAT DIP TO STORAGE CONTAINER.

ALWAYS ENSURE THAT DIPPED TEATS ARE DRY BEFORE EXPOSING THE ANIMALS TO WEATHER CONDITIONS WHICH MAY CAUSE DAMAGE TO THE TEATS.

**A PART OF THE COMPLETE IBA SANITATION PROGRAM**

EXP. DATE:      LOT NO.:      NET CONTENTS: 55 GALLONS (208.197 LITERS)      IBA STOCK # 261042      1121W

MANUFACTURED EXCLUSIVELY FOR IBA INC., SANITATION DIVISION, 103 GILMORE DRIVE, SUTTON, MA 01590

**ICON 5000 II**

iodine liquid

**Product Information**

<b>Product Type</b>	OTC ANIMAL DRUG	<b>Item Code (Source)</b>	NDC:29983-1018
<b>Route of Administration</b>	TOPICAL		

**Active Ingredient/Active Moiety**

Ingredient Name	Basis of Strength	Strength
<b>IODINE</b> (UNII: 9679TC07X4) (IODINE - UNII:9679TC07X4)	IODINE	0.5 kg in 100 kg

**Packaging**

#	Item Code	Package Description	Marketing Start Date	Marketing End Date
1	NDC:29983-1018-9	1188.75 kg in 1 CONTAINER		

**Marketing Information**

Marketing Category	Application Number or Monograph Citation	Marketing Start Date	Marketing End Date
unapproved drug other		09/27/2014	

**Labeler** - IBA (019494160)

**Registrant** - IBA (019494160)

### Establishment

Name	Address	ID/FEI	Business Operations
Morgan Gallacher Inc. DBA Custom Chemical Formulators Inc.		028311595	manufacture, api manufacture, pack

Revised: 11/2023

IBA