

**TUSSIN MULTI SYMPTOM COLD CF ADULT- dextromethorphan hbr,  
guaifenesin, phenylephrine liquid  
AAFES/Your Military Exchanges**

-----

**Drug Facts**

**Active ingredients (in each 10 mL)**

Dextromethorphan HBr 20 mg

Guaifenesin 200 mg

Phenylephrine HCl 10 mg

**Purposes**

Cough suppressant

Expectorant

Nasal decongestant

**Uses**

- helps loosen phlegm (mucus) and thin bronchial secretions to drain bronchial tubes
- temporarily relieves these symptoms occurring with a cold:
  - nasal congestion
  - cough due to minor throat and bronchial irritation

**Warnings**

**Do not use**

- if you are now taking a prescription monoamine oxidase inhibitor (MAOI) (certain drugs for depression, psychiatric or emotional conditions, or Parkinson's disease), or for 2 weeks after stopping the MAOI drug. If you do not know if your prescription drug contains an MAOI, ask a doctor or pharmacist before taking this product.

**Ask a doctor before use if you have**

- heart disease
- diabetes
- high blood pressure
- thyroid disease
- cough that lasts or is chronic such as occurs with smoking, asthma, chronic bronchitis or emphysema
- cough that occurs with too much phlegm ( mucus)
- trouble urinating due to enlarged prostate gland

**Ask a doctor or pharmacist before use if you are**

taking any other oral nasal decongestant or stimulant.

**When using this product,  
do not use more than directed.**

**Stop use and ask a doctor if**

- nervousness, dizziness, or sleeplessness occur
- symptoms do not get better within 7 days or are accompanied by fever
- cough lasts more than 7 days, comes back, or is accompanied by fever, rash, or persistent headache.

These could be signs of a serious condition.

**If pregnant or breast-feeding,**

ask a health professional before use.

**Keep out of reach of children.**

In case of overdose, get medical help or contact a Poison Control Center (1-800-222-1222) right away.

**Directions**

- do not take more than 6 doses in any 24-hour period
- measure only with dosing cup provided. Do not use any other dosing device.
- keep dosing cup with product
- mL=milliliter
- this adult product is not intended for use in children under 12 years of age
- adults and children 12 years and over: 10 mL every 4 hours
- children under 12 years: do not use

**Other information**

- store between 20-25°C (68°-77°F). Do not refrigerate.

**Inactive ingredients**

anhydrous citric acid, FD&C red #40, flavor, glycerin, lactic acid, menthol, propylene glycol, purified water, sodium benzoate, sorbitol, sucralose

**Questions or comments?**

Call **1-877-753-3935** Monday-Friday 9AM-5PM EST

**Principal Display Panel**

\*Compare to the active ingredients in Robitussin® Peak Cold Multi-Symptom Cold CF

Adult

**Tussin**

Multi-Symptom Cold CF

Dextromethorphan HBr

Cough Suppressant

Guaifenesin

Expectorant

Phenylephrine HCl

Nasal Decongestant

Relieves:

- Cough
- Mucus
- Nasal Congestion

Non-drowsy

Alcohol-Free

For Ages 12 Years & over

Dosing Cup Included

FL OZ (mL)

\*This product is not manufactured or distributed by Pfizer Consumer Healthcare, distributors of Robitussin® Peak Cold Multi-Symptom Cold CF.

**TAMPER EVIDENT: DO NOT USE IF CARTON IS OPENED OR IF PRINTED SAFETY SEAL AROUND BOTTLE OR UNDER CAP IS BROKEN OR MISSING.**

Manufactured for Your Military Exchanges

By: PL Developments, 11865 S. Alameda St

Lynwood, CA 90262

**Package Label**

**Drug Facts (continued)**

**Inactive ingredients** citric acid, FD&C red #40, flavor, glycerin, lactic acid, menthol, propylene glycol, purified water, sodium benzoate, sorbitol, sucralose

**Questions or comments?**  
Call 1-877-753-3835 Monday-Friday 9AM-5PM EST

\*This product is not manufactured or distributed by Pfizer Consumer Healthcare, distributor of Robitussin® Peak Cold Multi-Symptom Cold CF.

TAMPER EVIDENT: DO NOT USE IF CARTON IS OPENED OR IF PRINTED SAFETY SEAL AROUND BOTTLE OR UNDER CAP IS BROKEN OR MISSING.

**PARENTS:**  
Learn about teen medicine abuse  
[www.StopMedicineAbuse.org](http://www.StopMedicineAbuse.org)



Dosing Cup Included

\*SATISFACTION GUARANTEED OR YOUR MONEY BACK.\*  
Manufactured For Your Military Exchanges  
By: **PL Developments**, 11865 S. Alameda St  
Lynwood, CA 90262

\*Compare to active ingredients in Robitussin® Peak Cold Multi-Symptom Cold CF

exchange select

## ADULT Tussin Multi-Symptom Cold CF

Dextromethorphan HBr Cough Suppressant  
Guaifenesin Expectorant  
Phenylephrine HCl Nasal Decongestant

Relieves:

- Cough
- Mucus
- Nasal Congestion

Non-drowsy Alcohol-Free

For Ages 12 Years & Over  
8 fl oz (237 mL)

quality value

\*Compare to active ingredients in Robitussin® Peak Cold Multi-Symptom Cold CF

exchange select

## ADULT Tussin Multi-Symptom Cold CF

Dextromethorphan HBr Cough Suppressant  
Guaifenesin Expectorant  
Phenylephrine HCl Nasal Decongestant

Relieves:

- Cough
- Mucus
- Nasal Congestion

Non-drowsy Alcohol-Free

For Ages 12 Years & Over  
8 fl oz (237 mL)

quality value

**Drug Facts**

<b>Active ingredients (in each 10 mL)</b>	<b>Purposes</b>
Dextromethorphan HBr 20 mg	Cough suppressant
Guaifenesin 200 mg	Expectorant
Phenylephrine HCl 10 mg	Nasal decongestant

**Uses**

- helps loosen phlegm (mucus) and thin bronchial secretions to drain bronchial tubes
- temporarily relieves these symptoms occurring with a cold
- nasal congestion
- cough due to minor throat and bronchial irritation

**Warnings**

Do not use if you are now taking a prescription monoamine oxidase inhibitor (MAOI) (certain drugs for depression, psychiatric, or emotional conditions, or Parkinson's disease), or for 2 weeks after stopping the MAOI drug. If you do not know if your prescription drug contains an MAOI, ask a doctor or pharmacist before taking this product.

Ask a doctor before use if you have

- heart disease
- high blood pressure
- diabetes
- thyroid disease
- trouble urinating due to an enlarged prostate gland
- cough that occurs with too much phlegm (mucus)
- cough that lasts or is chronic such as occurs with smoking, asthma, chronic bronchitis or emphysema

Ask a doctor or pharmacist before use if you are taking any other oral nasal decongestant or stimulant.

When using this product, do not use more than directed.

Stop use and ask a doctor if

- nervousness, dizziness, or sleeplessness occur
- symptoms do not get better within 7 days or are accompanied by fever
- cough lasts more than 7 days, comes back, or is accompanied by fever, rash, or persistent headache. These could be signs of a serious condition.

If pregnant or breast-feeding, ask a health professional before use.

Keep out of reach of children.  
In case of overdose, get medical help or contact a Poison Control Center (1-800-222-1222) right away.

**Directions**

- do not take more than 6 doses in any 24-hour period
- measure only with dosing cup provided. Do not use any other dosing device.
- keep dosing cup with product
- mL = milliliter
- this adult product is not intended for use in children under 12 years of age
- adults and children 12 years and over: 10 mL every 4 hours
- children under 12 years: do not use

**Other information**

- store between 20-25°C (68-77°F). Do not refrigerate.



PLD-F322C F0007426

List No.:  
Exp. Date:

### EXCHANGE SELECT Adult Tussin Multi-Symptom Cold

## TUSSIN MULTI SYMPTOM COLD CF ADULT

dextromethorphan hbr, guaifenesin, phenylephrine liquid

### Product Information

<b>Product Type</b>	HUMAN OTC DRUG	<b>Item Code (Source)</b>	NDC:55301-382
<b>Route of Administration</b>	ORAL		

### Active Ingredient/Active Moiety

Ingredient Name	Basis of Strength	Strength
<b>DEXTROMETHORPHAN HYDROBROMIDE</b> (UNII: 9D2RTI9KYH) (DEXTROMETHORPHAN - UNII:7355X3ROTS)	DEXTROMETHORPHAN HYDROBROMIDE	20 mg in 10 mL

<b>GUAIFENESIN</b> (UNII: 495W7451VQ) (GUAIFENESIN - UNII:495W7451VQ)	GUAIFENESIN	200 mg in 10 mL
<b>PHENYLEPHRINE HYDROCHLORIDE</b> (UNII: 04JA59TNSJ) (PHENYLEPHRINE - UNII:1WS297W6MV)	PHENYLEPHRINE HYDROCHLORIDE	10 mg in 10 mL

### Inactive Ingredients

Ingredient Name	Strength
<b>ANHYDROUS CITRIC ACID</b> (UNII: XF417D3PSL)	
<b>GLYCERIN</b> (UNII: PDC6A3C0OX)	
<b>PROPYLENE GLYCOL</b> (UNII: 6DC9Q167V3)	
<b>WATER</b> (UNII: 059QF0KO0R)	
<b>SODIUM BENZOATE</b> (UNII: OJ245FE5EU)	
<b>FD&amp;C RED NO. 40</b> (UNII: WZB9127XOA)	
<b>LACTIC ACID</b> (UNII: 33X04XA5AT)	
<b>MENTHOL</b> (UNII: L7T10EIP3A)	
<b>SORBITOL</b> (UNII: 506T60A25R)	
<b>SUCRALOSE</b> (UNII: 96K6UQ3ZD4)	

### Packaging

#	Item Code	Package Description	Marketing Start Date	Marketing End Date
1	NDC:55301-382-08	1 in 1 BOX	08/31/2015	08/31/2025
1		237 mL in 1 BOTTLE, PLASTIC; Type 0: Not a Combination Product		

### Marketing Information

Marketing Category	Application Number or Monograph Citation	Marketing Start Date	Marketing End Date
OTC Monograph Drug	M012	08/31/2015	08/31/2025

**Labeler** - AAFES/Your Military Exchanges (001695568)

Revised: 2/2024

AAFES/Your Military Exchanges