REGULAR STRENGTH ANTACID- aluminum hydroxide, magnesium hydroxide, dimethicone suspension CVS

CVS ANTACID original

Active ingredients (in each 10 mL dose)

Aluminum hydroxide 400 mg (equivalent to dried gel, USP) Magnesium hydroxide 400 mg Simethicone 40 mg

Purposes

Antacid Antigas

Uses

relieves

- heartburn
- sour stomach
- acid indigestion
- the symptoms referred to as gas

Warnings

Ask a doctor before use if you have

- kidney disease
- a magnesium-restricted diet

Ask a doctor or pharmacist before use if you are taking a prescription drug. Antacids may interact with certain prescription drugs.

Stop use and ask a doctor if symptoms last more than 2 weeks **If pregnant or breast-feeding,** ask a health professional before use.

Keep out of reach of children.

Directions

- shake well before use
- adults and children 12 years and older: take 10 mL to 20 mL (1 to 2 doses) between meals as needed, at bedtime, or as directed by a doctor
- do not exceed 120 mL (12 doses) in a 24 hour period or use the maximum dosage for more than 2 weeks
- children under 12 years: ask a doctor

Other information

- each 10 mL dose contains: magnesium 165 mg, sodium 6 mg
- store at room temperature
- protect from freezing
- keep tightly closed

Inactive ingredients

benzyl alcohol, butylparaben, flavor (contains alcohol), hydroxyethylcellulose, propylparaben, purified water, saccharin sodium, sorbitol solution

Questions or comments? 1-800-540-3765

package Label



Regular Strength

Antacid Liquid

ALUMINUM HYDROXIDE / MAGNESIUM HYDROXIDE / SIMETHICONE Antacid / Anti-Gas

Fast-Acting

Soothing relief of:

- Heartburn
- Acid indigestion
- Pressure & bloating

Plus gas relief

Original Flavor ALCOHOL 0.15%

12 FL 0Z (355 mL)

205-05112-1 **REV 0721**

Drug Facts TAMPER-EVIDENT: Do not use if breakaway band on bottle cap is broken or missing.

Purpose

Active ingredients (in each 10 mL dose)

uminum hydroxide

(equivalent to dried gel, USP) 400 mg Antacid Magnesium hydroxide 400 mg ------ Antacid Simethicone 40 mg ------ Antigas

Uses relieves

- · heartburn · sour stomach
- acid indigestion
 the symptoms referred to as gas

Warnings

Ask a doctor before use if you have

kidney disease + a magnesium-restricted diet

Ask a doctor or pharmacist before use if you are presently taking a prescription drug. Antacids may interact with certain prescription drugs.

Stop use and ask a doctor if symptoms last more than 2 weeks

If pregnant or breast-feeding, ask a health professional before use. Keep out of reach of children,

Directions • shake well before use

- adults and children 12 years of age and older: take 10 mL to 20 mL.
 (1 to 2 doses) between meals as needed, at bedtime, or as directed by a doctor
- do not exceed 120 ml. (12 doses) in a 24 hour period or use the maximum dosage for more than 2 weeks children under 12 years of age: ask a doctor

Other information

- each 10 mL dose contains: magnesium 165 mg, sodium 6 mg
- store at room temperature protect from freezing
- keep tightly closed

Inactive ingredients benzyl alcohol, butylparaben, flavor (contains alcohol), hydroxyethylcellulose, propylparaben, purified water, saccharin sodium, sorbito solution

Questions or comments? 1-800-540-3765

Distributed by: CVS Pharmacy, Inc. One CVS Drive

Woonsocket, RI 02895 © 2021 CVS/pharmacy 0 CVS.com®

1-800-SHOP CVS V-37297

905-05112-1 REV GC0721

REGULAR STRENGTH ANTACID

aluminum hydroxide, magnesium hydroxide, dimethicone suspension

Product Information

Product Type HUMAN OTC DRUG Item Code (Source) NDC:69842-286

Route of Administration ORAL

Active Ingredient/Active Moiety

Ingredient Name	Basis of Strength	Strength
ALUMINUM HYDROXIDE (UNII: 5QB0T2IUN0) (ALUMINUM HYDROXIDE - UNII:5QB0T2IUN0)	ALUMINUM HYDROXIDE	400 mg in 10 mL
MAGNESIUM HYDROXIDE (UNII: NBZ3QY004S) (MAGNESIUM CATION - UNII:T6V3LHY838, HYDROXIDE ION - UNII:9159UV381P)	MAGNESIUM HYDROXIDE	400 mg in 10 mL

DIMETHICONE (UNII: 92RU3N3Y10) (DIMETHICONE - UNII:92RU3N3Y10) DIMETHICONE	in 10 mL
---	----------

Inactive Ingredients		
Ingredient Name	Strength	
BENZYL ALCOHOL (UNII: LKG8494WBH)		
BUTYLPARABEN (UNII: 3QPI1U3FV8)		
HYDROXYETHYL CELLULOSE, UNSPECIFIED (UNII: T4V6TWG28D)		
PROPYLPARABEN (UNII: Z8IX2SC10H)		
WATER (UNII: 059QF0KO0R)		
SACCHARIN SODIUM (UNII: SB8ZUX40TY)		
SORBITOL SOLUTION (UNII: 8KW3E207O2)		

l	Packaging				
	#	Item Code	Package Description	Marketing Start Date	Marketing End Date
	1	NDC:69842-286- 12	355 mL in 1 BOTTLE; Type 0: Not a Combination Product	06/01/2017	

Marketing Information			
Marketing Category	Application Number or Monograph Citation	Marketing Start Date	Marketing End Date
OTC Monograph Drug	M001	06/01/2015	

Labeler - CVS (062312574)

Registrant - GCP Laboratories (965480861)

Establishment				
Name	Address	ID/FEI	Business Operations	
GCP Laboratories		965480861	manufacture(69842-286) , pack(69842-286)	

Revised: 10/2023 CVS