

## **LIL DRUG STORE CHILDRENS PAIN AND FEVER- acetaminophen suspension Lil' Drug Store Products, Inc.**

*Disclaimer: Most OTC drugs are not reviewed and approved by FDA, however they may be marketed if they comply with applicable regulations and policies. FDA has not evaluated whether this product complies.*

-----  
**Children's Pain & Fever, Lil' Drug Store ®**

### **Drug Facts**

#### **Active ingredient (in each 5 mL)**

Acetaminophen 160 mg

#### **Purpose**

Pain reliever/fever reducer

#### **Uses**

- temporarily reduces fever
- temporarily relieves minor aches and pains due to
  - the common cold
  - flu
  - headache
  - sore throat
  - toothache

#### **Warnings**

##### **Liver warning**

This product contains acetaminophen. Severe liver damage may occur if your child takes:

- more than 5 doses in 24 hours, which is the maximum daily amount
- with other drugs containing acetaminophen

##### **Allergy alert**

Acetaminophen may cause severe skin reactions. Symptoms may include:

- skin reddening
- blisters
- rash

If a skin reaction occurs, stop use and seek medical help right away.

##### **Sore throat warning**

If sore throat is severe, persists for more than 2 days, is accompanied or followed by fever, headache, rash, nausea, or vomiting, consult a doctor promptly.

### **Do not use**

- with any other drug containing acetaminophen (prescription or nonprescription). If you are not sure whether a drug contains acetaminophen, ask a doctor or pharmacist.
- if your child is allergic to acetaminophen or any of the inactive ingredients in this product

**Ask a doctor before use if your child has** liver disease.

**Ask a doctor or pharmacist before use if your child is** taking the blood thinning drug warfarin.

**When using this product, do not exceed recommended dose (see Overdose warning).**

### **Stop use and ask a doctor if**

- new symptoms occur
- redness or swelling is present
- pain gets worse or lasts more than 5 days
- fever gets worse or lasts more than 3 days

These could be signs of a serious condition.

**Keep out of reach of children.**

### **Overdose warning**

Taking more than the recommended dose (overdose) may cause liver damage. In case of overdose, get medical help or contact a Poison Control Center (1-800-222-1222) right away. Quick medical attention is critical even if you do not notice any signs or symptoms.

### **Directions**

- **this product does not contain directions or complete warnings for adult use**
- **do not take more than directed (see Overdose warning)**
- **shake well before using**
- mL = milliliter
- use only the enclosed dosing cup designed for use with this product. Do not use any other dosing device.
- if needed, repeat dose every 4 hours while symptoms last
- do not give more than 5 times in 24 hours
- do not give more than 5 days unless directed by a doctor
- find the right dose on chart below. If possible, use weight to dose; otherwise, use age.

<b>Weight (lb)</b>	<b>Age (year)</b>	<b>Dose (mL) *</b>
--------------------	-------------------	--------------------

under 24	under 2	ask a doctor
24-35	2-3	5 mL
36-47	4-5	7.5 mL
48-59	6-8	10 mL
60-71	9-10	12.5 mL
72-95	11	15 mL

\* or as directed by a doctor

### Other information

- **each 5 mL contains:** sodium 2 mg
- store between 20-25°C (68-77°F)
- do not refrigerate
- see bottom panel for lot number and expiration date

### Inactive ingredients

acesulfame potassium, butylparaben, carboxymethylcellulose sodium, citric acid, FD&C red #40, flavors, glycerin, high fructose corn syrup, microcrystalline cellulose, propylene glycol, purified water, sodium benzoate, sorbitol, xanthan gum

### Questions or comments?

Call toll free **1-877-507-6516 (M-F 8AM-4:30PM CST)**

### Product distributed by:

Lil' Drug Store Products, Inc.  
1201 Continental Place NE  
Cedar Rapids, IA 52402

### PRINCIPAL DISPLAY PANEL - 118 mL Bottle Carton

QUALITY

\* GUARANTEED \*

Compare to the Active Ingredient  
in Children's Tylenol®  
Oral Suspension\*\*

Children's  
Pain & Fever

Acetaminophen  
160 mg per 5 mL  
Pain Reliever/Fever Reducer

Oral Suspension  
For ages 2-11 years

Cherry

4 FL OZ (118 mL)

Lil'  
Drug Store®

# Children's Pain & Fever

## Acetaminophen

160 mg per 5 mL  
Pain Reliever/Fever Reducer

4 FL OZ (118 mL)



Compare to the Active Ingredient  
in Children's Tylenol®  
Oral Suspension\*\*

# Children's Pain & Fever

## Acetaminophen

160 mg per 5 mL  
Pain Reliever/Fever Reducer

Oral Suspension  
For ages 2-11 years



Cherry

**TAMPER EVIDENT: DO NOT USE IF CARTON IS  
OPENED OR IF PRINTED SAFETY SEAL AROUND  
BOTTLE OR UNDER CAP IS BROKEN OR MISSING.**

### Drug Facts

Active ingredient (in each 5 mL)	Purpose
Acetaminophen 160 mg.....	Pain reliever/fever reducer

**Uses** ■ temporarily reduces fever  
■ temporarily relieves minor aches and pains due to  
■ the common cold ■ flu ■ headache  
■ sore throat ■ toothache

### Warnings

**Liver warning:** This product contains acetaminophen. Severe liver damage may occur if your child takes:

- more than 5 doses in 24 hours, which is the maximum daily amount
- with other drugs containing acetaminophen

**Allergy alert:** Acetaminophen may cause severe skin reactions. Symptoms may include: ■ skin reddening ■ blisters ■ rash. If a skin reaction occurs, stop use and seek medical help right away.

**Sore throat warning:** If sore throat is severe, persists for more than 2 days, is accompanied or followed by fever, headache, rash, nausea, or vomiting, consult a doctor promptly.

### Do not use

- with any other drug containing acetaminophen (prescription or nonprescription). If you are not sure whether a drug contains acetaminophen, ask a doctor or pharmacist.
- if your child is allergic to acetaminophen or any of the inactive ingredients in this product

Ask a doctor before use if your child has liver disease.

Ask a doctor or pharmacist before use if your child is taking the blood thinning drug warfarin.

When using this product, do not exceed recommended dose (see Overdose warning).

Stop use and ask a doctor if ■ new symptoms occur  
■ redness or swelling is present  
■ pain gets worse or lasts more than 5 days  
■ fever gets worse or lasts more than 3 days  
These could be signs of a serious condition.

Keep out of reach of children.

4 FL OZ (118 mL)



Keep out of reach of children.

**Overdose warning:** Taking more than the recommended dose (overdose) may cause liver damage. In case of overdose, get medical help or contact a Poison Control Center (1-800-222-1222) right away. Quick medical attention is critical even if you do not notice any signs or symptoms.

PLD-D231D FC004017 68564C-US-07-16-1



Lot No.:

Exp. Date:

NTOC

**Drug Facts** (continued)

**Directions**

- this product does not contain directions or complete warnings for adult use
- do not take more than directed (see Overdose warning)
- shake well before using
- mL = milliliter
- use only the enclosed dosing cup designed for use with this product. Do not use any other dosing device.
- if needed, repeat dose every 4 hours while symptoms last
- do not give more than 5 times in 24 hours
- do not give more than 5 days unless directed by a doctor
- find the right dose on chart below. If possible, use weight to



Compare to the Active Ingredient in Children's Tylenol® Oral Suspension\*\*

Children's  
**Pain & Fever**

NIOC

dose; otherwise, use age.

Weight (lb)	Age (year)	Dose (mL)*
under 24	under 2	ask a doctor
24-35	2-3	5 mL
36-47	4-5	7.5 mL
48-59	6-8	10 mL
60-71	9-10	12.5 mL
72-95	11	15 mL

\*or as directed by a doctor

**Other information**

- each 5 mL contains: sodium 2 mg
- store between 20-25°C (68-77°F) ■ do not refrigerate
- see bottom panel for lot number and expiration date

**Inactive ingredients** acesulfame potassium, butylparaben, carboxymethylcellulose sodium, citric acid, FD&C red #40, flavors, glycerin, high fructose corn syrup, microcrystalline cellulose, propylene glycol, purified water, sodium benzoate, sorbitol, xanthan gum

**Questions or comments?** Call toll free 1-877-507-6516 (M-F 8AM-4:30PM CST)

\*\*This product is not manufactured or distributed by McNeil Consumer Healthcare, distributor of Children's Tylenol® Oral Suspension.

**Product distributed by:**  
Lil' Drug Store Products, Inc.  
1201 Continental Place NE  
Cedar Rapids, IA 52402

**Acetaminophen**

**160 mg per 5 mL**

Pain Reliever/Fever Reducer

**Oral Suspension**  
**For ages 2-11 years**



Cherry

4 FL OZ (118 mL)



NIOC

**LIL DRUG STORE CHILDRENS PAIN AND FEVER**

acetaminophen suspension

**Product Information**

<b>Product Type</b>	HUMAN OTC DRUG	<b>Item Code (Source)</b>	NDC:66715-6956
<b>Route of Administration</b>	ORAL		

### Active Ingredient/Active Moiety

Ingredient Name	Basis of Strength	Strength
ACETAMINOPHEN (UNII: 362O9ITL9D) (ACETAMINOPHEN - UNII:362O9ITL9D)	ACETAMINOPHEN	160 mg in 5 mL

### Inactive Ingredients

Ingredient Name	Strength
ACESULFAME POTASSIUM (UNII: 23OV73Q5G9)	
ANHYDROUS CITRIC ACID (UNII: XF417D3PSL)	
BUTYLPARABEN (UNII: 3QPI1U3FV8)	
CARBOXYMETHYLCELLULOSE SODIUM, UNSPECIFIED FORM (UNII: K679OBS311)	
FD&C RED NO. 40 (UNII: WZB9127XOA)	
GLYCERIN (UNII: PDC6A3C0OX)	
HIGH FRUCTOSE CORN SYRUP (UNII: XY6UN3QB6S)	
MICROCRYSTALLINE CELLULOSE (UNII: OP1R32D61U)	
PROPYLENE GLYCOL (UNII: 6DC9Q167V3)	
WATER (UNII: 059QF0KO0R)	
SODIUM BENZOATE (UNII: OJ245FE5EU)	
SORBITOL (UNII: 506T60A25R)	
XANTHAN GUM (UNII: TTV12P4NEE)	

### Product Characteristics

<b>Color</b>		<b>Score</b>	
<b>Shape</b>		<b>Size</b>	
<b>Flavor</b>	CHERRY	<b>Imprint Code</b>	
<b>Contains</b>			

### Packaging

#	Item Code	Package Description	Marketing Start Date	Marketing End Date
1	NDC:66715-6956-4	1 in 1 CARTON	12/01/2017	
1		118 mL in 1 BOTTLE, PLASTIC; Type 0: Not a Combination Product		

### Marketing Information

Marketing Category	Application Number or Monograph Citation	Marketing Start Date	Marketing End Date
OTC monograph not final	part343	12/01/2014	

Revised: 12/2022

Lil' Drug Store Products, Inc.