

**IBUPROFEN 200- ibuprofen capsule, liquid filled**  
**WALGREEN COMPANY**

-----  
***Drug Facts***

***Active ingredient (in each capsule)***

Solubilized ibuprofen equal to 200 mg ibuprofen (NSAID)\*  
(present as the free acid and potassium salt)  
\*nonsteroidal anti-inflammatory drug

***Purpose***

Pain reliever/Fever reducer

***Uses***

- temporarily relieves minor aches and pains due to:
- headache
- toothache
- backache
- menstrual cramps
- the common cold
- muscular aches
- minor pain of arthritis
- temporarily reduces fever

***Warnings***

**Allergy alert:** Ibuprofen may cause a severe allergic reaction, especially in people allergic to aspirin. Symptoms may include:

- hives
- facial swelling
- asthma (wheezing)
- shock
- skin reddening
- rash
- blisters

If an allergic reaction occurs, stop use and seek medical help right away.

**Stomach bleeding warning:** This product contains an NSAID, which may cause severe stomach bleeding. The chance is higher if you

- are age 60 or older
- have had stomach ulcers or bleeding problems
- take a blood thinning (anticoagulant) or steroid drug
- take other drugs containing prescription or nonprescription NSAIDs [aspirin, ibuprofen, naproxen, or others]
- have 3 or more alcoholic drinks every day while using this product
- take more or for a longer time than directed

**Heart attack and stroke warning:** NSAIDs, except aspirin, increase the risk of heart attack, heart failure, and stroke. These can be fatal. The risk is higher if you use more than directed or for longer than directed.

**Do not use**

- if you have ever had an allergic reaction to any other pain reliever/fever reducer
- right before or after heart surgery

**Ask a doctor before use if**

- stomach bleeding warning applies to you
- you have problems or serious side effects from taking pain relievers or fever reducers
- you have a history of stomach problems, such as heartburn
- you have high blood pressure, heart disease, liver cirrhosis, kidney disease, asthma, or had a stroke
- you are taking a diuretic

**Ask a doctor or pharmacist before use if you are**

- under a doctor's care for any serious condition
- taking aspirin for heart attack or stroke, because ibuprofen may decrease this benefit of aspirin
- taking any other drug

**When using this product**

- take with food or milk if stomach upset occurs

**Stop use and ask a doctor if**

- you experience any of the following signs of stomach bleeding:
  - feel faint
  - vomit blood
  - have bloody or black stools
  - have stomach pain that does not get better
- you have symptoms of heart problems or stroke:
  - chest pain
  - trouble breathing
  - weakness in one part or side of body
  - slurred speech
  - leg swelling
- pain gets worse or lasts more than 10 days
- fever gets worse or lasts more than 3 days
- redness or swelling is present in the painful area
- any new symptoms appear

**If pregnant or breast-feeding,**

ask a health professional before use. It is especially important not to use ibuprofen during the last 3 months of pregnancy unless definitely directed to do so by a doctor because it may cause problems in the unborn child or complications during delivery.

**Keep out of reach of children.**

In case of overdose, get medical help or contact a Poison Control Center right away.

### ***Directions***

- **do not take more than directed**
- **the smallest effective dose should be used**
- adults and children 12 years and over: take 1 capsule every 4 to 6 hours while symptoms persist
- if pain or fever does not respond to 1 capsule, 2 capsules may be used
- do not exceed 6 capsules in 24 hours, unless directed by a doctor
- children under 12 years: ask a doctor

### ***Other information***

- each capsule contains: **potassium 20 mg**
- read all warnings and directions before use
- store at 20° to 25°C (68° to 77°F)
- avoid excessive heat above 40°C (104°F). Protect from light.

### ***Inactive ingredients***

FD&C Red #40, FD&C Yellow #6, gelatin, pharmaceutical ink, polyethylene glycol, potassium hydroxide, purified water, sorbitan and sorbitol

### ***Questions or comments?***

call toll free **1-888-235-2466**

**DO NOT USE IF TAMPER-EVIDENT SEAL UNDER BOTTLE CAP IMPRINTED WITH “SEALED for YOUR PROTECTION” IS BROKEN OR MISSING.**

††This product is not manufactured or distributed by Johnson & Johnson Corporation, owner of the registered trademark Motrin® IB Liquid Gels.

DISTRIBUTED BY: WALGREEN CO.

200 WILMOT RD., DEERFIELD, IL 60015

**100% SATISFACTION GUARANTEED**

walgreens.com ©2019 Walgreen Co.

ITEM 295646

W00000-0000-0 L0000219

Rev # 05/19

**ORG0719-F**

### **120's Bottle Label**

*Compare to Motrin® IB Liquid Gels active ingredient††*

**NDC 0363-0222-22**

**Ibuprofen 200**

**IBUPROFEN CAPSULES 200 mg /**

**PAIN RELIEVER / FEVER REDUCER (NSAID)**

**SOFTGELS**

120 SOFTGELS\*\*

(\*\*LIQUID-FILLED CAPSULES)



Compare to Motrin® IB  
Liquid Gels active ingredient††

NDC 0363-0222-22

# Ibuprofen 200

IBUPROFEN CAPSULES 200 mg /  
PAIN RELIEVER / FEVER REDUCER (NSAID)

SOFTGELS

120 SOFTGELS\*\*  
(\*\*LIQUID-FILLED CAPSULES)

**DO NOT USE IF TAMPER-EVIDENT SEAL UNDER BOTTLE CAP IMPRINTED WITH "SEALED FOR YOUR PROTECTION" IS BROKEN OR MISSING**

|   |   |  |   |
|---|---|--|---|
| <p><b>Drug Facts</b></p> <p><b>Active ingredient (in each capsule)</b><br/>Solubilized ibuprofen equal to 200 mg ibuprofen (NSAID)* ..... Pain reliever/fever reducer (present as the free acid and potassium salt) *nonsteroidal anti-inflammatory drug</p> <p><b>Uses</b> ■ temporarily relieves minor aches and pains due to:<br/>■ headache ■ toothache ■ backache ■ menstrual cramps ■ the common cold ■ muscular aches ■ minor pain of arthritis ■ temporarily reduces fever</p> <p><b>Warnings</b><br/><b>Allergy alert:</b> Ibuprofen may cause a severe allergic reaction, especially in people allergic to aspirin. Symptoms may include:<br/>■ hives ■ facial swelling ■ asthma (wheezing) ■ shock ■ skin redness ■ rash ■ blisters. If an allergic reaction occurs, stop use and seek medical help right away.<br/><b>Stomach bleeding warning:</b> This product contains an NSAID which may cause severe stomach bleeding. The chance is higher if you ■ are age 60 or older ■ have had stomach ulcers or bleeding problems ■ take a blood thinning (anticoagulant) or steroid drug ■ take other drugs containing prescription or nonprescription NSAIDs (aspirin, ibuprofen, naproxen, or others) ■ have 3 or more alcoholic drinks every day while using this product ■ take more or for a longer time than directed<br/><b>Heart attack and stroke warning:</b> NSAIDs, except aspirin, increase the risk of heart attack, heart failure, and stroke. These can be fatal. The risk is higher if you use more than directed or for longer than directed.</p> <p><b>Do not use</b> ■ if you have ever had an allergic reaction to any other pain reliever/fever reducer ■ right before or after heart surgery</p> | <p><b>120 SOFTGELS**</b><br/>(**LIQUID-FILLED CAPSULES)</p> | <p>11917 21029 2</p> <p>Exp. Date: ORG0719-E</p> | <p>PEEL HERE</p> <p style="text-align: center; color: red; font-weight: bold;">Non Varnish Area</p> |
|---|---|--|---|

††STANDARD BY WALGREEN CO.  
20 WILMUTAIL DRIVE, DEERFIELD, ILLINOIS  
100% SATISFACTION GUARANTEED  
Walgreens.com ©2019 Walgreen Co.  
ITEM 2935946  
NDC 0363-0222-22 1000219 3  
Net Wt. 0.979 g    Lot No.

|  |  |
|--|--|
| <p><b>Drug Facts (continued)</b></p> <p><b>Ask a doctor before use if</b> ■ stomach bleeding warning applies to you ■ you have problems or serious side effects from taking pain relievers or fever reducers ■ you have a history of stomach problems, such as heartburn ■ you have high blood pressure, heart disease, liver cirrhosis, kidney disease, asthma, or had a stroke ■ you are taking a diuretic</p> <p><b>Ask a doctor or pharmacist before use if you are</b> ■ under a doctor's care for any serious condition ■ taking aspirin for heart attack or stroke, because ibuprofen may decrease this benefit of aspirin ■ taking any other drug</p> <p><b>When using this product</b> ■ take with food or milk if stomach upset occurs</p> <p><b>Stop use and ask a doctor if</b> ■ you experience any of the following signs or stomach bleeding: ■ feel faint ■ vomit blood ■ have bloody or black stools ■ you have stomach pain that does not get better ■ you have symptoms of heart problems or stroke: ■ chest pain ■ trouble breathing ■ weakness in one part or side of body ■ slurred speech ■ leg swelling ■ pain gets worse or lasts more than 10 days ■ fever gets worse or lasts more than 3 days ■ redness or swelling is present in the painful area ■ any new symptoms appear</p> <p><b>If pregnant or breast-feeding</b>, ask a health professional before use. It is especially important not to use ibuprofen during the last 3 months of pregnancy unless definitely directed to do so by a doctor because it may cause problems in the unborn child or complications during delivery.<br/><b>Keep out of reach of children.</b> In case of overdose, get medical help or contact a Poison Control Center right away.</p> <p><b>Directions</b></p> <p>■ do not take more than directed ■ the smallest effective dose should be used ■ adults and children 12 years and over: take 1 capsule every 4 to 6 hours while symptoms persist ■ if pain or fever does not respond to 1 capsule, 2 capsules may be used ■ do not exceed 6 capsules in 24 hours, unless directed by a doctor ■ children under 12 years: ask a doctor</p> <p><b>Other information</b></p> <p>■ each capsule contains: potassium 20 mg ■ read all warnings and directions before use ■ store at 20° to 25°C (68° to 77°F) ■ avoid excessive heat above 40°C (104°F). Protect from light.</p> <p><b>Inactive ingredients</b> FD&amp;C Red #40, FD&amp;C Yellow #6, gelatin, pharmaceutical ink, polyethylene glycol, potassium hydroxide, purified water, sorbitan and sorbitol</p> <p><b>Questions or comments?</b> call toll free 1-888-235-2466</p> | <p>††This product is not manufactured or distributed by Johnson &amp; Johnson Corporation, owner of the registered trademark Motrin® IB Liquid Gels.</p> |
|--|--|

|  |                |                    |               |
|--|----------------|--------------------|---------------|
| <h2 style="margin: 0;">IBUPROFEN 200</h2> <p style="margin: 0;">ibuprofen capsule, liquid filled</p> |                |                    |               |
| <h3 style="margin: 0;">Product Information</h3>  |                |                    |               |
| Product Type   | HUMAN OTC DRUG | Item Code (Source) | NDC:0363-0222 |
| Route of Administration  | ORAL           |                    |               |
| <h3 style="margin: 0;">Active Ingredient/Active Moiety</h3>  |                |                    |               |

| Ingredient Name  |  | Basis of Strength                                  | Strength             |                    |
|--|--|--|----------------------|--------------------|
| IBUPROFEN (UNII: WK2XYI10QM) (IBUPROFEN - UNII:WK2XYI10QM) |  | IBUPROFEN  | 200 mg               |                    |
| <b>Inactive Ingredients</b>                                |  |  |                      |                    |
| Ingredient Name  |  |  | Strength             |                    |
| FD&C RED NO. 40 (UNII: WZB9127XOA)                         |  |  |                      |                    |
| FD&C YELLOW NO. 6 (UNII: H77VEI93A8)                       |  |  |                      |                    |
| GELATIN (UNII: 2G86QN327L)                                 |  |  |                      |                    |
| POLYETHYLENE GLYCOL, UNSPECIFIED (UNII: 3WJQ0SDW1A)        |  |  |                      |                    |
| POTASSIUM HYDROXIDE (UNII: WZH3C48M4T)                     |  |  |                      |                    |
| WATER (UNII: 059QF0K00R)                                   |  |  |                      |                    |
| SORBITAN (UNII: 6O92ICV9RU)                                |  |  |                      |                    |
| SORBITOL (UNII: 506T60A25R)                                |  |  |                      |                    |
| <b>Product Characteristics</b>                             |  |  |                      |                    |
| Color  | orange                                   | Score  | no score             |                    |
| Shape  | capsule                                  | Size   | 20mm                 |                    |
| Flavor   |  | Imprint Code                                       | IB2                  |                    |
| Contains   |  |  |                      |                    |
| <b>Packaging</b>   |  |  |                      |                    |
| #  | Item Code                                | Package Description                                | Marketing Start Date | Marketing End Date |
| 1  | NDC:0363-0222-22                         | 120 in 1 BOTTLE; Type 0: Not a Combination Product | 08/02/2019           |                    |
| <b>Marketing Information</b>                               |  |  |                      |                    |
| Marketing Category   | Application Number or Monograph Citation | Marketing Start Date                               | Marketing End Date   |                    |
| ANDA   | ANDA078682                               | 08/02/2019   |                      |                    |

**Labeler** - WALGREEN COMPANY (008965063)

**Registrant** - Bionpharma Inc. (079637826)

### Establishment

| Name                  | Address | ID/FEI    | Business Operations    |
|-----------------------|---------|-----------|------------------------|
| Patheon Softgels Inc. |         | 002193829 | manufacture(0363-0222) |