IBUPROFEN- ibuprofen tablet, film coated Stephen L. LaFrance Pharmacy, Inc.

Select Brand 44-392

Active ingredient (in each tablet)

Ibuprofen USP, 200 mg (NSAID)*

*nonsteroidal anti-inflammatory drug

Purpose

Pain reliever/fever reducer

Uses

- temporarily relieves minor aches and pains due to:
 - headache
 - toothache
 - menstrual cramps
 - the common cold
 - backache
 - muscular aches
 - minor pain of arthritis
- temporarily reduces fever

Warnings

Allergy alert: Ibuprofen maay cause a severe allergic reaction, especially in people allergy to aspirin. Sympoms may include:

- facial swelling
- rash
- hives
- asthma (wheezing)
- shock
- skin reddening
- blisters

If an allergic reaction occurs, stop use and seek medical help right away.

Stomach bleeding warning: This product contains an NSAID, which may cause severe stomach bleeding. The chance is higher if you:

- are age 60 or older
- have had stomach ulcers or bleeding problems
- take a blood thinning (anticoagulant) or steroid drug
- take other drugs containing prescription or nonprescription NSAIDs [aspirin, ibuprofen, naproxen, or others]
- have 3 or more alcoholic drinks every day while using this product
- take more or for a longer time than directed

Do not use

- if you have ever had an allergic reaction to any other pain reliever/fever reducer
- right before or after heart surgery

Ask a doctor before use if

- stomach bleeding warning applies to you
- you have a history of stomach problems, such as heartburn
- you have high blood pressure, heart disease, liver cirrhosis, or kidney disease
- you are taking a diuretic
- you have problems or serious side effects from taking pain relievers or fever reducers
- vou have asthma

Ask a doctor or pharmacist before use if you are

- under a doctor's care for any serious condition
- taking aspirin for heart attack or stroke, because ibuprofen may decrease this benefit of aspirin
- taking any other drug

When using this product

- take with food or milk if stomach upset occurs
- the risk of heart attack or stroke may increase if you use more than directed or for longer than directed

Stop use and ask a doctor if

- you experience any of the following signs of stomach bleeding:
- feel faint
- have bloody or black stools
- vomit blood
- have stomach pain that does not get better
- pain gets worse or lasts more than 10 days
- any new symptoms appear
- redness or swelling is present in the painful area
- fever gets worse or lasts more than 3 days

If pregnant or breast-feeding,

ask a health professional before use. It is especially important not to use ibuprofen during the last 3 months of pregnancy unless definitely directed to do so by a doctor because it may cause problems in the unborn child or complications during delivery.

Keep out of reach of children.

In case of overdose, get medical help or contact a Poison Control Center right away.

Directions

- do not take more than directed
- the smallest effective dose should be used
- do not take longer than 10 days, unless directed by a doctor (see Warnings)
- adults and children 12 years and over: take 1 tablet every 4 to 6 hours while symptoms persist
- if pain or fever does not respond to 1 tablet, 2 tablets may be used
- do not exceed 6 tablets in 24 hours, unless directed by a doctor
- children under 12 years: ask a doctor

Other information

- store between 20°-25°C (68°-77°F)
- avoid excessive heat 40°C (104°F)
- used by expiration date on package

Inactive ingredients

carnauba wax, cellulose, corn starch, FD&C yellow #6 aluminum lake, fumed silica gel, hypromellose, lactose, magnesium stearate, polydextrose, polyethylene glycol, sodium starch glycolate, stearic acid, titanium dioxide

Questions

Call 1-800-426-9391 8:30 AM-4:00 PM ET, Monday-Friday

Principal Display Panel

select brand®

the lower price name brand

NDC 15127-905-50

IBUPROFEN
IBUPROFEN TABLETS USP, 200 mg
PAIN RELIEVER/FEVER REDUCER

†Compare to the Active Ingredient of Motrin® IB

50 Film Coated Orange Tablets

TAMPER EVIDENT: DO NOT USE IF IMPRINTED SAFETY SEAL UNDER CAP IS BROKEN OR MISSING


†This product is not manufactured or distributed by McNeil Consumer Healthcare, owner of the registered trademark Motrin® IB.

50844 REV0411B39215

Distributed by:

SELECT BRAND® DISTRIBUTORS

Pine Bluff, AR 71603 USA



Select Brand 44-392

IBUPROFEN

ibuprofen tablet, film coated Product Information Product Type HUMAN OTC DRUG LABEL Item Code (Source) NDC:15127-905 Route of Administration ORAL DEA Schedule

Active Ingredient/Active Moiety		
Ingredient Name	Basis of Strength	Strength
IBUPRO FEN (IBUPRO FEN)	IBUPROFEN	200 mg

Inactive Ingredients		
Ingredient Name	Strength	
CARNAUBA WAX		
FD&C YELLOW NO. 6		
HYPROMELLOSES		
LACTOSE		
MAGNESIUM STEARATE		
POLYDEXTROSE		
POLYETHYLENE GLYCOLS		
SODIUM STARCH GLYCOLATE TYPE A POTATO		
STEARIC ACID		
TITANIUM DIO XIDE		

Product Characteristics			
Color	ORANGE	Score	no score
Shape	ROUND	Size	10 mm
Flavor		Imprint Code	44;392
Contains			

Packaging			
# Item Code	Package Description	Marketing Start Date	Marketing End Date
1 NDC:15127-905-50	1 in 1 CARTON		
1	50 in 1 BOTTLE		

Marketing Information				
Marketing Category	Application Number or Monograph Citation	Marketing Start Date	Marketing End Date	
ANDA	ANDA075139	03/30/2002		

Labeler - Stephen L. LaFrance Pharmacy, Inc. (043562370)

Establishment				
Name	Address	ID/FEI	Business Operations	
LNK International, Inc.		038154464	PACK(15127-905)	

Establishment			
Name	Address	ID/FEI	Business Operations

LNK International, Inc. 832867894 MANUFACTURE(15127-905)

Revised: 10/2012 Stephen L. LaFrance Pharmacy, Inc.