

**TOPCARE PAIN RELIEF- acetaminophen tablet**  
**Topco Associates LLC**

*Disclaimer: Most OTC drugs are not reviewed and approved by FDA, however they may be marketed if they comply with applicable regulations and policies. FDA has not evaluated whether this product complies.*

-----

**Topco Associates LLC. Pain Relief Drug Facts**

**Active ingredient (in each caplet)**

Acetaminophen 500 mg

**Purpose**

Pain reliever/fever reducer

**Uses**

- temporarily relieves minor aches and pains due to:
- the common cold
- headache
- backache
- minor pain of arthritis
- toothache
- muscular aches
- premenstrual and menstrual cramps
- temporarily reduces fever

**Warnings**

Liver warning: This product contains acetaminophen. Severe liver damage may occur if you take

- more than 4,000 mg of acetaminophen in 24 hours
- with other drugs containing acetaminophen
- 3 or more alcoholic drinks every day while using this product

Allergy alert: Acetaminophen may cause severe skin reactions. Symptoms may include:

- skin reddening
- blisters
- rash

If a skin reaction occurs, stop use and seek medical help right away.

**Do not use**

- with any other drug containing acetaminophen (prescription or nonprescription). If you are not sure whether a drug contains acetaminophen, ask a doctor or pharmacist.
- if you have ever had an allergic reaction to this product or any of its ingredients

**Ask a doctor before use if you have**

liver disease

**Ask a doctor or pharmacist before use if you are**

taking the blood thinning drug warfarin

**Stop use and ask a doctor if**

- pain gets worse or lasts more than 10 days
- fever gets worse or lasts more than 3 days
- new symptoms occur
- redness or swelling is present

These could be signs of a serious condition.

**If pregnant or breast-feeding,**

ask a health professional before use.

**Keep out of reach of children.**

Overdose warning: In case of overdose, get medical help or contact a Poison Control Center right away (1-800-222-1222). Quick medical attention is critical for adults as well as for children even if you do not notice any signs or symptoms.

**Directions**

- do not take more than directed (see overdose warning)

adults and children 12 years and over	<ul style="list-style-type: none"><li>• take 2 caplets every 6 hours while symptoms last</li><li>• do not take more than 6 caplets in 24 hours, unless directed by a doctor</li><li>• do not use for more than 10 days unless directed by a doctor</li></ul>
children under 12 years	ask a doctor

**Other information**

- store at 20-25°C (68-77°F)

**Inactive ingredients**

carnauba wax, corn starch\*, croscarmellose sodium\*, hypromellose, polyethylene glycol, povidone, pregelatinized starch, sodium starch glycolate\*, stearic acid

\*may contain one or more of these ingredients

**Questions or comments?**

1-888-423-0139

**Principal Display Panel**

EXTRA STRENGTH

Pain Relief

ACETAMINOPHEN 500 mg

Pain Reliever / Fever Reducer

For Adults

SEE NEW WARNINGS

THIS PACKAGE FOR HOUSEHOLDS WITHOUT YOUNG CHILDREN

COMPARE TO EXTRA STRENGTH TYLENOL® CAPLETS active ingredient

actual size

150 CAPLETS



**DO NOT USE IF PRINTED SEAL UNDER  
CAP IS BROKEN OR MISSING**

: 97547 88 F5

## **Drug Facts**

<b>Active ingredient (in each caplet)</b>	<b>Purpose</b>
---	----------------

Acetaminophen 500 mg.....	Pain reliever/ fever reducer
---------------------------	---------------------------------

## **Uses**

- temporarily relieves minor aches and pains due to:
  - the common cold
  - minor pain of arthritis
  - muscular aches
  - premenstrual and menstrual cramps
  - headache
  - backache
  - toothache
- temporarily reduces fever

## **Warnings**

**Liver warning:** This product contains acetaminophen. Severe liver damage may occur if you take

- more than 4,000 mg of acetaminophen in 24 hours
- with other drugs containing acetaminophen
- 3 or more alcoholic drinks every day while using this product

**Allergy alert:** Acetaminophen may cause severe skin reactions. Symptoms may include:

- skin reddening
- blisters
- rash

If a skin reaction occurs, stop use and seek medical help right away.

## **Do not use**

- with any other drug containing acetaminophen (prescription or nonprescription). If you are not sure whether a drug contains acetaminophen, ask a doctor or pharmacist.
- if you have ever had an allergic reaction to this product or any of its ingredients

**Ask a doctor before use if you have liver disease**

**Ask a doctor or pharmacist before use if you are taking the blood thinning drug warfarin**

## **Stop use and ask a doctor if**

- pain gets worse or lasts more than 10 days
  - fever gets worse or lasts more than 3 days
  - new symptoms occur
  - redness or swelling is present
- These could be signs of a serious condition.

**If pregnant or breast-feeding, ask a health professional before use.**

**Keep out of reach of children. Overdose warning:** In case of overdose, get medical help or contact a Poison Control Center right away (1-800-222-1222). Quick medical attention is



critical for adults  
as well as for  
children even if  
you do not notice  
any signs or  
symptoms. ▶

**PEEL BACK HERE ▶**



## ***Drug Facts*** (continued)

### ***Directions***

■ do not take more than directed  
(see overdose warning)

adults and children 12 years and over	<ul style="list-style-type: none"><li>■ take 2 caplets every 6 hours while symptoms last</li><li>■ do not take more than 6 caplets in 24 hours, unless directed by a doctor</li><li>■ do not use for more than 10 days unless directed by a doctor</li></ul>
children under 12 years	ask a doctor

### ***Other information***

■ store at 20-25°C (68-77°F)

***Inactive ingredients*** carnauba wax, corn starch\*, croscarmellose sodium\*, hypromellose, polyethylene glycol, povidone, pregelatinized starch, sodium starch glycolate\*, stearic acid  
\*may contain one or more of these ingredients

### ***Questions or comments?***

1-888-423-0139

# TOPCARE PAIN RELIEF

acetaminophen tablet

## Product Information

Product Type	HUMAN OTC DRUG LABEL	Item Code (Source)	NDC:36800-484
Route of Administration	ORAL	DEA Schedule	

## Active Ingredient/Active Moiety

Ingredient Name	Basis of Strength	Strength
ACETAMINOPHEN (ACETAMINOPHEN)	ACETAMINOPHEN	500 mg

## Inactive Ingredients

Ingredient Name	Strength
CARNAUBA WAX	
STARCH, CORN	
HYPROMELLOSES	
POLYETHYLENE GLYCOLS	
POVIDONES	
STEARIC ACID	
CROSCARMELLOSE SODIUM	

## Product Characteristics

Color	WHITE	Score	no score
Shape	CAPSULE (caplet)	Size	16mm
Flavor		Imprint Code	L484
Contains			

## Packaging

#	Item Code	Package Description	Marketing Start Date	Marketing End Date
1	NDC:36800-484-76	1 in 1 CARTON		
1		120 in 1 BOTTLE		
2	NDC:36800-484-62	1 in 1 CARTON		
2		24 in 1 BOTTLE		
3	NDC:36800-484-71	1 in 1 CARTON		
3		50 in 1 BOTTLE		
4	NDC:36800-484-78	1 in 1 CARTON		
4		100 in 1 BOTTLE		
5	NDC:36800-484-90	500 in 1 BOTTLE		
6	NDC:36800-484-82	2 in 1 CARTON		
6		100 in 1 BOTTLE		
7	NDC:36800-484-47	150 in 1 BOTTLE		

## Marketing Information

Marketing Category	Application Number or Monograph Citation	Marketing Start Date	Marketing End Date
OTC monograph not final	part343	07/15/1987	

**Labeler** - Topco Associates LLC (006935977)

Revised: 12/2014

Topco Associates LLC