

IBUPROFEN- ibuprofen capsule, liquid filled
Winco Foods, LLC

WNC - 1185A - 2019-1016

Drug Facts

Active ingredient (in each capsule)

Solubilized ibuprofen equal to
200 mg ibuprofen (NSAID)*

* nonsteroidal anti-inflammatory drug

Purpose

Pain reliever/fever reducer

Uses

- temporarily relieves minor aches and pains due to:
 - headache
 - toothache
 - backache
 - menstrual cramps
 - the common cold
 - muscular aches
 - minor pain of arthritis
- temporarily reduces fever

Warnings

Allergy alert

Ibuprofen may cause a severe allergic reaction, especially in people allergic to aspirin. Symptoms may include:

- hives
- facial swelling
- asthma (wheezing)
- shock
- skin reddening
- rash
- blisters

If an allergic reaction occurs, stop use and seek medical help right away.

Stomach bleeding warning

This product contains an NSAID, which may cause severe stomach bleeding. The chance is higher if you:

- are age 60 or older
- have had stomach ulcers or bleeding problems

- take a blood thinning (anticoagulant) or steroid drug
- take other drugs containing prescription or nonprescription NSAIDs (aspirin, ibuprofen, naproxen, or others)
- have 3 or more alcoholic drinks every day while using this product
- take more or for a longer time than directed

Heart attack and stroke warning

NSAIDs, except aspirin, increase the risk of heart attack, heart failure, and stroke. These can be fatal. The risk is higher if you use more than directed or for longer than directed.

Do not use

- if you have ever had an allergic reaction to any other pain reliever/fever reducer
- right before or after heart surgery

Ask a doctor before use if

- stomach bleeding warning applies to you
- you have problems or serious side effects from taking pain relievers or fever reducers
- you have a history of stomach problems, such as heartburn
- you have high blood pressure, heart disease, liver cirrhosis, kidney disease, asthma, or had a stroke
- you are taking a diuretic

Ask a doctor or pharmacist before use if you are

- under a doctor's care for any serious condition
- taking aspirin for heart attack or stroke, because ibuprofen may decrease this benefit of aspirin
- taking any other drug

When using this product

- take with food or milk if stomach upset occurs

Stop use and ask a doctor if

- side effects occur. You may report side effects to FDA at 1-800-FDA-1088.
- you experience any of the following signs of stomach bleeding:
 - feel faint
 - vomit blood
 - have bloody or black stools
 - have stomach pain that does not get better
- you have symptoms of heart problems or stroke:
 - chest pain
 - trouble breathing
 - weakness in one part or side of body
 - slurred speech
 - leg swelling
- pain gets worse or lasts more than 10 days
- fever gets worse or lasts more than 3 days
- redness or swelling is present in the painful area

- any new symptoms appear

If pregnant or breast-feeding, ask a health professional before use. It is especially important not to use ibuprofen during the last 3 months of pregnancy unless definitely directed to do so by a doctor because it may cause problems in the unborn child or complications during delivery.

Keep out of reach of children. In case of overdose, get medical help or contact a Poison Control Center right away (1-800-222-1222).

Directions

- **do not take more than directed**
- **the smallest effective dose should be used**
- adults and children 12 years and over:
 - take 1 capsule every 4 to 6 hours while symptoms persist
 - if pain or fever does not respond to 1 capsule, 2 capsules may be used
 - do not exceed 6 capsules in 24 hours, unless directed by a doctor
- children under 12 years: ask a doctor

Other information

- each capsule contains: **potassium 20 mg**
- store at 20-25°C (68-77°F)
- avoid excessive heat above 40°C (104°F)
- read all warnings and directions before use
- retain carton for complete product information and warnings

Inactive ingredients

ammonium hydroxide, FD&C green #3, gelatin, iron oxide black, medium chain triglycerides, polyethylene glycol, propylene glycol, potassium hydroxide, purified water, shellac, sorbitan monooleate, sorbitol sorbitan

Questions?

Call 800-824-1706 Monday - Friday 9am - 4pm MST

PRINCIPAL DISPLAY PANEL

WinCo Foods

Ibuprofen

Compare to the active ingredient in ADVIL Liqui-Gels

Relieves: Minor Aches & Pains, Fever

IBUPROFEN CAPSULES

200 mg (NSAID)/

PAIN RELIEVER – FEVER REDUCER

80 Softgels**

** Liquid Filled Capsules

INK AND COATING FREE
FOR LOT AND
EXPIRATION STAMPING



0R_NC 11854MWC03095R11F



Drug Facts

Active Ingredient (in each capsule)

Sulfalated ibuprofen sodium to 200 mg ibuprofen (NSAID).....fever reducer (present as the free acid and potassium salt) nonsteroidal anti-inflammatory drug

Purpose

Pain reliever/fever reducer

Uses

- temporarily relieves minor aches and pains due to:
 - headache
 - toothache
 - backache
 - muscular cramps
 - the common cold
- muscular aches
- minor pain of arthritis
- temporarily reduces fever

Warnings

Allergy alert: Ibuprofen may cause a severe allergic reaction, especially in people allergic to aspirin. Symptoms may include:

- hives
- facial swelling
- shock
- skin reddening
- asthma (wheezing)
- rash
- blisters

If an allergic reaction occurs, stop use and seek medical help right away.

Stomach bleeding warning: This product contains an NSAID, which may cause severe stomach bleeding. The chance is higher if you:

- have had stomach ulcers or bleeding problems
- are age 60 or older
- take a blood thinning (anticoagulant) or steroid drug
- take other drugs containing prescription or over-the-counter NSAIDs (aspirin, ibuprofen, naproxen, or celecoxib)
- have a more alcoholic drinks every day while using this product
- take more or for a longer time than directed

Drug Facts (continued)

Heart attack and stroke warning: NSAIDs, except aspirin, increase the risk of heart attack, heart failure, and stroke. These can be fatal. The risk is higher if you use more than directed or for longer than directed.

Do not use if you have ever had an allergic reaction to any other pain reliever/fever reducer right before or after heart surgery

Ask a doctor before use if

- you have problems or serious side effects from taking pain relievers or fever reducers
- you have high blood pressure, heart disease, liver cirrhosis, kidney disease, asthma, or had a stroke
- you are taking a diuretic
- you have a history of stomach problems, such as heartburn
- you have had a stomach ulcer, stomach pain, or stomach bleeding
- you are taking any other drug

When using this product

- take with food or milk if stomach upset occurs
- stop use and ask a doctor if side effects occur. You may report side effects to FDA at 1-800-FDA-1088.
- you experience any of the following signs of stomach bleeding:
 - feel faint
 - vomit blood
 - have bloody or black stools
- have stomach pain that does not get better
- you have symptoms of heart problems or stroke:
 - chest pain
 - windness in one part or side of body
 - slurred speech
 - leg swelling
- pain gets worse or lasts more than 10 days

Drug Facts (continued)

Fever gets worse or lasts more than 3 days
redness or swelling is present in the painful area
any new symptoms appear

If pregnant or breast-feeding, ask a health professional before use. It is especially important not to use ibuprofen during the last 3 months of pregnancy unless specifically directed to do so by a doctor because it may cause problems in the unborn child or complications during delivery. Keep out of reach of children. In case of overdose, get medical help or contact a Poison Control Center right away (1-800-222-1222).

Directions

- do not take more than directed
- the smallest effective dose should be used
- adults and children 12 years and over:
 - if pain or fever does not respond to 1 capsule, 2 capsules may be used
 - do not exceed 6 capsules in 24 hours, unless directed by a doctor
- children under 12 years: ask a doctor

Other information

- each capsule contains potassium 20 mg
- store at 20-25°C (68-77°F)
- store excursions permitted to 15-30°C (59-86°F)
- read all warnings and directions before use
- retain carton for complete product information and warnings

Inactive ingredients

ammonium hydroxide, FD&C green #3, gelatin, iron oxide black, medium chain triglycerides, polyethylene glycol, propylene glycol, potassium hydroxide, purified water, stibic acid, sorbitan monostearate, sorbitol sorbitan

Questions? Call 800-824-1786 Monday - Friday 9am - 4pm MST

The Supermarket Low Price Leader®
100% SATISFACTION GUARANTEE



MADE IN INDIA
BOISE, ID 83704
WINCO FOODS, LLC.

DO NOT USE IF IMPRINTED SEAL UNDER CAP IS BROKEN OR MISSING.
Consumer Healthcare, distributor of Advil® Liqui-Gels®.



WinCo FOODS

Ibuprofen

IBUPROFEN CAPSULES
200 mg (NSAID)/
PAIN RELIEVER - FEVER REDUCER

80 Softgels**

**Liquid Filled Capsules

†Compare to the active ingredient in ADVIL® Liqui-Gels®
Relieves: Minor Aches & Pains, Fever

IBUPROFEN
ibuprofen capsule, liquid filled

Product Information			
Product Type	HUMAN OTC DRUG	Item Code (Source)	NDC:67091-185
Route of Administration	ORAL		

Active Ingredient/Active Moiety

Ingredient Name	Basis of Strength	Strength
IBUPROFEN (UNII: WK2XYI10QM) (IBUPROFEN - UNII:WK2XYI10QM)	IBUPROFEN	200 mg

Inactive Ingredients

Ingredient Name	Strength
AMMONIA (UNII: 5138Q19F1X)	
FD&C GREEN NO. 3 (UNII: 3P3ONR6O1S)	
GELATIN (UNII: 2G86QN327L)	
FERROUS OXIDE (UNII: G7036X8B5H)	
MEDIUM-CHAIN TRIGLYCERIDES (UNII: C9H2L21V7U)	
POLYETHYLENE GLYCOL, UNSPECIFIED (UNII: 3WJQ0SDW1A)	
PROPYLENE GLYCOL (UNII: 6DC9Q167V3)	
POTASSIUM HYDROXIDE (UNII: WZH3C48M4T)	
WATER (UNII: 059QF0K00R)	
SHELLAC (UNII: 46N107B71O)	
SORBITAN MONOOLEATE (UNII: 06XEA2VD56)	
SORBITOL (UNII: 506T60A25R)	

Product Characteristics

Color	green	Score	no score
Shape	CAPSULE	Size	19mm
Flavor		Imprint Code	133
Contains			

Packaging

#	Item Code	Package Description	Marketing Start Date	Marketing End Date
1	NDC:67091-185-03	1 in 1 CARTON	05/01/2018	
1		80 in 1 BOTTLE, PLASTIC; Type 0: Not a Combination Product		

Marketing Information

Marketing Category	Application Number or Monograph Citation	Marketing Start Date	Marketing End Date
ANDA	ANDA079205	05/01/2018	

Labeler - Winco Foods, LLC (056098817)