

**SMART SENSE IBUPROFEN PM- diphenhydramine citrate and ibuprofen tablet, coated**  
**Kmart Corporation**

-----  
**Kmart Corporation Ibuprofen PM Drug Facts**

**Active ingredients (in each caplet)**

Diphenhydramine citrate 38 mg

Ibuprofen 200 mg (NSAID)\*

\*nonsteroidal anti-inflammatory drug

**Purposes**

Nighttime sleep-aid

Pain reliever

**Uses**

- for relief of occasional sleeplessness when associated with minor aches and pains
- helps you fall asleep and stay asleep

**Warnings**

Allergy alert: Ibuprofen may cause a severe allergic reaction, especially in people allergic to aspirin. Symptoms may include:

- hives
- facial swelling
- asthma (wheezing)
- shock
- skin reddening
- rash
- blisters

If an allergic reaction occurs, stop use and seek medical help right away.

Stomach bleeding warning: This product contains an NSAID, which may cause severe stomach bleeding. The chance is higher if you

- are age 60 or older
- have had stomach ulcers or bleeding problems
- take a blood thinning (anticoagulant) or steroid drug
- take other drugs containing prescription or nonprescription NSAIDs [aspirin, ibuprofen, naproxen, or others]
- have 3 or more alcoholic drinks every day while using this product
- take more or for a longer time than directed

**Do not use**

- if you have ever had an allergic reaction to any other pain reliever/fever reducer
- unless you have time for a full night's sleep
- in children under 12 years of age
- right before or after heart surgery
- with any other product containing diphenhydramine, even one used on skin
- if you have sleeplessness without pain

#### **Ask a doctor before use if**

- stomach bleeding warning applies to you
- you have problems or serious side effects from taking pain relievers or fever reducers
- you have a history of stomach problems, such as heartburn
- you have high blood pressure, heart disease, liver cirrhosis, kidney disease, or asthma
- you are taking a diuretic
- you have a breathing problem such as emphysema or chronic bronchitis
- you have glaucoma
- you have trouble urinating due to an enlarged prostate gland

#### **Ask a doctor or pharmacist before use if you are**

- taking sedatives or tranquilizers, or any other sleep-aid
- under a doctor's care for any continuing medical illness
- taking any other antihistamines
- taking aspirin for heart attack or stroke, because ibuprofen may decrease this benefit of aspirin
- taking any other drug

#### **When using this product**

- drowsiness will occur
- avoid alcoholic drinks
- do not drive a motor vehicle or operate machinery
- take with food or milk if stomach upset occurs
- the risk of heart attack or stroke may increase if you use more than directed or for longer than directed

#### **Stop use and ask a doctor if**

- you experience any of the following signs of stomach bleeding:
- feel faint
- vomit blood
- have bloody or black stools
- have stomach pain that does not get better
- pain gets worse or lasts more than 10 days
- sleeplessness persists continuously for more than 2 weeks. Insomnia may be a symptom of a serious underlying medical illness.
- redness or swelling is present in the painful area
- any new symptoms appear

**If pregnant or breast-feeding,**

ask a health professional before use. It is especially important not to use ibuprofen during the last 3 months of pregnancy unless definitely directed to do so by a doctor because it may cause problems in the unborn child or complications during delivery.

**Keep out of reach of children.**

In case of overdose, get medical help or contact a Poison Control Center right away. (1-800-222-1222)

**Directions**

- do not take more than directed
- adults and children 12 years and over: take 2 caplets at bedtime
- do not take more than 2 caplets in 24 hours

**Other information**

- read all warnings and directions before use. Keep carton.
- store at 20-25°C (68-77°F)
- avoid excessive heat above 40°C (104°F)

**Inactive ingredients**

colloidal silicon dioxide, croscarmellose sodium, FD&C blue no. 2 aluminum lake, glyceryl behenate, hydroxypropyl cellulose, iron oxide black, lactose monohydrate, magnesium stearate, microcrystalline cellulose, polyethylene glycol, polyvinyl alcohol, pregelatinized starch, talc, titanium dioxide

**Questions or comments?**

1-800-719-9260

**Principal Display Panel**

Compare to the active ingredients of Advil® PM Caplets

PAIN RELIEVER/NIGHTTIME SLEEP-AID

Ibuprofen PM

Ibuprofen 200 mg / Pain Reliever (NSAID)

Diphenhydramine Citrate 38 mg

Nighttime Sleep-Aid Tablets

ACTUAL SIZE

(\*\*Capsule-Shaped Tablets)

20 COATED CAPLETS\*\*

NDC 49738-191-60

Compare to the active ingredients  
of Advil® PM Caplets\*\*\*

Smart  
sense®

PAIN RELIEVER/NIGHTTIME SLEEP-AID

**Ibuprofen PM**

Ibuprofen 200 mg / Pain Reliever (NSAID)  
Diphenhydramine Citrate 38 mg  
Nighttime Sleep-Aid Tablets

ACTUAL SIZE



(\*\*\* Capsule-Shaped Tablets) **20 COATED CAPLETS\*\*\***

DO NOT USE IF PRINTED FOIL UNDER CAP  
IS BROKEN OR MISSING



LOT NO.

EXP.

: OSDBO FT CH

**Drug Facts (continued)**

**Inactive ingredients** colloidal silicon dioxide, croscarmellose sodium, FD&C blue no. 2 aluminum lake, glyceryl behenate, hydroxypropyl cellulose, iron oxide black, lactose monohydrate, magnesium stearate, microcrystalline cellulose, polyethylene glycol, polyvinyl alcohol, pregelatinized starch, talc, titanium dioxide

**Questions or comments?** 1-800-719-9260

**READ AND KEEP CARTON FOR COMPLETE WARNINGS AND INFORMATION.**

\*\*\*This product is not manufactured or distributed by Wyeth Consumer Healthcare, distributor of Advil® PM Caplets.

**SATISFACTION  
GUARANTEE**

GLUTEN  
FREE

Distributed by: Kmart Corporation  
Hoffman Estates, IL 60179  
**Shop kmart.com**

Thank you for purchasing this quality product. If you are unsatisfied for any reason, return the unused portion to the store for a full refund or call 1-800-642-7886.

**Drug Facts (continued)**

**Stop use and ask a doctor if**

- you experience any of the following signs of stomach bleeding:
    - feel faint
    - vomit blood
    - have bloody or black stools
  - have stomach pain that does not get better
  - pain gets worse or lasts more than 10 days
  - sleeplessness persists continuously for more than 2 weeks.
- Insomnia may be a symptom of a serious underlying medical illness.
- redness or swelling is present in the painful area
  - any new symptoms appear

**if pregnant or breast-feeding**, ask a health professional before use. It is especially important not to use ibuprofen during the last 3 months of pregnancy unless definitely directed to do so by a doctor because it may cause problems in the unborn child or complications during delivery. **Keep out of reach of children.** In case of overdose, get medical help or contact a Poison Control Center right away. (1-800-222-1222)

**Directions** ■ do not take more than directed ■ adults and children 12 years and over: take 2 caplets at bedtime ■ do not take more than 2 caplets in 24 hours

**Other information** ■ read all warnings and directions before use. Keep carton. ■ store at 20°-25° C (68°-77° F) ■ avoid excessive heat above 40° C (104° F)



### Drug Facts

Active Ingredients (In each caplet)	Purposes
Diphenhydramine citrate 38 mg.....	Nighttime sleep-aid
Ibuprofen 200 mg (NSAID)*.....	Pain reliever
*nonsteroidal anti-inflammatory drug	

**Uses** ■ for relief of occasional sleeplessness when associated with minor aches and pains  
 ■ helps you fall asleep and stay asleep

### Warnings

**Allergy alert:** Ibuprofen may cause a severe allergic reaction, especially in people allergic to aspirin. Symptoms may include:  
 ■ hives ■ facial swelling ■ asthma (wheezing)  
 ■ shock ■ skin reddening ■ rash ■ blisters  
 If an allergic reaction occurs, stop use and seek medical help right away.

**Stomach bleeding warning:** This product contains an NSAID, which may cause severe stomach bleeding.

- The chance is higher if you
- are age 60 or older
  - have had stomach ulcers or bleeding problems
  - take a blood thinning (anticoagulant) or steroid drug
  - take other drugs containing prescription or nonprescription NSAIDs [aspirin, ibuprofen, naproxen, or others]
  - have 3 or more alcoholic drinks every day while using this product ■ take more or for a longer time than directed

**Do not use** ■ if you have ever had an allergic reaction to any other pain reliever/fever reducer  
 ■ unless you have time for a full night's sleep ▶

### Drug Facts (continued)

- in children under 12 years of age
- right before or after heart surgery
- with any other product containing diphenhydramine, even one used on skin
- if you have sleeplessness without pain

### Ask a doctor before use if

- stomach bleeding warning applies to you
- you have problems or serious side effects from taking pain relievers or fever reducers
- you have a history of stomach problems, such as heartburn
- you have high blood pressure, heart disease, liver cirrhosis, kidney disease, or asthma ■ you are taking a diuretic
- you have a breathing problem such as emphysema or chronic bronchitis ■ you have glaucoma
- you have trouble urinating due to an enlarged prostate gland

### Ask a doctor or pharmacist before use if you are

- taking sedatives or tranquilizers, or any other sleep-aid
- under a doctor's care for any continuing medical illness
- taking any other antihistamines
- taking aspirin for heart attack or stroke, because ibuprofen may decrease this benefit of aspirin
- taking any other drug

### When using this product

- drowsiness will occur ■ avoid alcoholic drinks
- do not drive a motor vehicle or operate machinery
- take with food or milk if stomach upset occurs
- the risk of heart attack or stroke may increase if you use more than directed or for longer than directed ▶





# SMART SENSE IBUPROFEN PM

diphenhydramine citrate, ibuprofen tablet, coated

## Product Information

<b>Product Type</b>	HUMAN OTC DRUG LABEL	<b>Item Code (Source)</b>	NDC:49738-191
<b>Route of Administration</b>	ORAL	<b>DEA Schedule</b>	

## Active Ingredient/Active Moiety

Ingredient Name	Basis of Strength	Strength
DIPHENHYDRAMINE CITRATE (DIPHENHYDRAMINE)	DIPHENHYDRAMINE CITRATE	38 mg
IBUPROFEN (IBUPROFEN)	IBUPROFEN	200 mg

## Inactive Ingredients

Ingredient Name	Strength
SILICON DIOXIDE	
CROSCARMELLOSE SODIUM	
GLYCERYL DIBEHENATE	
HYDROXYPROPYL CELLULOSE (TYPE H)	
FERROSFERRIC OXIDE	
LACTOSE MONOHYDRATE	
MAGNESIUM STEARATE	
CELLULOSE, MICROCRYSTALLINE	
POLYETHYLENE GLYCOLS	
POLYVINYL ALCOHOL	
TALC	
TITANIUM DIOXIDE	

## Product Characteristics

<b>Color</b>	BLUE	<b>Score</b>	no score
<b>Shape</b>	CAPSULE (tablet)	<b>Size</b>	15mm
<b>Flavor</b>		<b>Imprint Code</b>	L050
<b>Contains</b>			

## Packaging

#	Item Code	Package Description	Marketing Start Date	Marketing End Date
1	NDC:49738-191-60	1 in 1 CARTON		
1		20 in 1 BOTTLE		
2	NDC:49738-191-27	1 in 1 CARTON		
2		80 in 1 BOTTLE		

## Marketing Information

Marketing Category	Application Number or Monograph Citation	Marketing Start Date	Marketing End Date
ANDA	ANDA079113	05/27/2010	

**Labeler** - Kmart Corporation (008965873)

Revised: 6/2014

Kmart Corporation