

MILK OF MAGNESIA- magnesium hydroxide liquid

HyVee Inc

Disclaimer: Most OTC drugs are not reviewed and approved by FDA, however they may be marketed if they comply with applicable regulations and policies. FDA has not evaluated whether this product complies.

Hy-Vee, Inc. Milk of Magnesia Drug Facts

Active ingredient (in each 15 mL Tablespoon)

Magnesium hydroxide 1200 mg

Purpose

Saline laxative

Uses

- relieves occasional constipation (irregularity)
- generally produces bowel movement in ½ to 6 hours

Warnings

Ask a doctor before use if you have

- kidney disease
- a magnesium-restricted diet
- stomach pain, nausea, or vomiting
- a sudden change in bowel habits that lasts more than 2 weeks

Ask a doctor or pharmacist before use if you are

taking any other drug. Take this product two or more hours before or after other drugs. Laxatives may affect how other drugs work.

Stop use and ask a doctor if

- you have rectal bleeding or no bowel movement after using this product. These could be signs of a serious condition.
- you need to use a laxative for more than 1 week

If pregnant or breast-feeding,

ask a health professional before use.

Keep out of reach of children.

In case of overdose, get medical help or contact a Poison Control Center right away. (1-800-222-1222)

Directions

- do not exceed the maximum recommended daily dose in a 24 hour period

- shake well before use
- dose may be taken once a day preferably at bedtime, in divided doses, or as directed by a doctor
- drink a full glass (8 oz) of liquid with each dose
- for accurate dosing, use dose cup provided
- mL = milliliter; TBSP = Tablespoon

adults and children 12 years and older	30 mL (2 TBSP) to 60 mL (4 TBSP)
children 6 to 11 years	15 mL (1 TBSP) to 30 mL (2 TBSP)
children under 6 years	ask a doctor

Other information

- each tablespoon (15 mL) contains: calcium 20 mg, magnesium 595 mg, potassium 2 mg and sodium 3 mg
- does not meet USP requirements for preservative effectiveness
- store at 20-25°C (68-77°F)
- close cap tightly after use
- avoid freezing

Inactive ingredients

anhydrous citric acid, D&C red #28, flavor, glycerin, purified water, sorbitol solution, sucrose, xanthan gum

Questions or comments?

1-800-719-9260

Principal Display Panel

COMPARE TO the Active Ingredient of Phillips'® Milk of Magnesia

Milk of Magnesia

Saline Laxative

Magnesium Hydroxide

Wild Cherry Flavor

#1 Doctor Recommended Saline Laxative Active Ingredient

Comfortable, Cramp Free Relief

Stimulant Free

ALCOHOL 0.4%

12 FL OZ (355 mL)

HyVee[®]
health

***COMPARE TO**
the Active Ingredient
of Phillips[®] Milk
of Magnesia

Milk of Magnesia

NDC 42507-949-40

Saline Laxative
Magnesium Hydroxide



Wild Cherry Flavor

- #1 Doctor Recommended Saline Laxative Active Ingredient**
- Comfortable, Cramp Free Relief
- Stimulant Free



ALCOHOL 0.4%

12 FL OZ (355 mL)

: 94940 2H F7

Drug Facts

DO NOT USE IF PRINTED NECKBAND IS BROKEN OR MISSING.

Active ingredient (in each 15 mL Tablespoon)

Magnesium hydroxide 1200 mg.....Saline laxative

Purpose

Uses ■ relieves occasional constipation (irregularity)
■ generally produces bowel movement in ½ to 6 hours

Warnings

Ask a doctor before use if you have

- kidney disease ■ a magnesium-restricted diet ■ stomach pain, nausea, or vomiting
- a sudden change in bowel habits that lasts more than 2 weeks

Ask a doctor or pharmacist before use if you are taking any other drug.

Take this product two or more hours before or after other drugs. Laxatives may affect how other drugs work.

Stop use and ask a doctor if ■ you have rectal bleeding or no bowel movement after using this product. These could be signs of a serious condition.

- you need to use a laxative for more than 1 week

If pregnant or breast-feeding, ask a health professional before use.

Keep out of reach of children. In case of overdose, get medical help or contact a Poison Control Center right away (1-800-222-1222).

Directions

- do not exceed the maximum recommended daily dose in a 24 hour period
- shake well before use
- dose may be taken once a day preferably at bedtime, in divided doses, or as directed by a doctor
- drink a full glass (8 oz) of liquid with each dose
- for accurate dosing, use dose cup provided ■ mL = milliliter; TBSP = Tablespoon

adults and children 12 years and older	30 mL (2 TBSP) to 60 mL (4 TBSP)
children 6 to 11 years	15 mL (1 TBSP) to 30 mL (2 TBSP)
children under 6 years	ask a doctor

Other information

- each tablespoon (15 mL) contains: calcium 20 mg, magnesium 595 mg, potassium 2 mg and sodium 3 mg
- does not meet USP requirements for preservative effectiveness
- store at 20-25°C (68-77°F) ■ close cap tightly after use ■ avoid freezing

Inactive ingredients anhydrous citric acid, D&C red #28, flavor, glycerin, purified water, sorbitol solution, sucrose, xanthan gum


Questions or comments? 1-800-719-9260

*This product is not manufactured or distributed by Bayer HealthCare LLC, distributor of Phillips'® Milk of Magnesia.

**In the U.S. Source: Symphony Health Solutions.

GLUTEN FREE

DISTRIBUTED BY HY-VEE, INC.
WEST DES MOINES, IA 50399



<div>MILK OF MAGNESIA</div> <div>magnesium hydroxide liquid</div>				
Product Information				
Product Type		HUMAN OTC DRUG LABEL	Item Code (Source)	
Route of Administration		ORAL	DEA Schedule	
Active Ingredient/Active Moiety				
Ingredient Name			Basis of Strength	
MAGNESIUM HYDRO XIDE (MAGNESIUM CATION and HYDRO XIDE ION)			MAGNESIUM HYDRO XIDE	
			1200 mg in 15 mL	
Inactive Ingredients				
Ingredient Name			Strength	
ANHYDRO US CITRIC ACID				
GLYCERIN				
WATER				
SORBITOL				
SUCROSE				
XANTHAN GUM				
D&C RED NO. 28				
Product Characteristics				
Color	PINK (Opaque Light pink)		Score	
Shape			Size	
Flavor	CHERRY (wild cherry)		Imprint Code	
Contains				
Packaging				
#	Item Code	Package Description	Marketing Start Date	Marketing End Date
1	NDC:42507-949-40	355 mL in 1 BOTTLE		
Marketing Information				
Marketing Category	Application Number or Monograph Citation		Marketing Start Date	Marketing End Date
OTC monograph not final	part334		07/18/2005	

