

MILK OF MAGNESIA- magnesium hydroxide liquid

HyVee Inc

Disclaimer: Most OTC drugs are not reviewed and approved by FDA, however they may be marketed if they comply with applicable regulations and policies. FDA has not evaluated whether this product complies.

Hy-Vee, Inc. Milk of Magnesia Drug Facts

Active ingredient (in each 15 mL TBSP)

Magnesium hydroxide 1200 mg

Purpose

Saline laxative

Uses

- relieves occasional constipation (irregularity)
- generally produces bowel movement in ½ to 6 hours

Warnings

Ask a doctor before use if you have

- kidney disease
- a magnesium-restricted diet
- stomach pain, nausea, or vomiting
- a sudden change in bowel habits that lasts more than 2 weeks

Ask a doctor or pharmacist before use if you are

taking any other drug. Take this product two or more hours before or after other drugs. Laxatives may affect how other drugs work.

Stop use and ask a doctor if

- you have rectal bleeding or no bowel movement after using this product. These could be signs of a serious condition.
- you need to use a laxative for more than 1 week

If pregnant or breast-feeding,

ask a health professional before use.

Keep out of reach of children.

In case of overdose, get medical help or contact a Poison Control Center right away. (1-800-222-1222)

Directions

- shake well before use

- do not exceed the maximum recommended daily dose in a 24 hour period
- dose may be taken once a day preferably at bedtime, in divided doses, or as directed by a doctor
- drink a full glass (8 oz) of liquid with each dose
- for accurate dosing, use dose cup provided
- TBSP = tablespoonful

adults and children 12 years and older	2 to 4 TBSP.
children 6 to 11 years	1 to 2 TBSP.
children under 6 years	ask a doctor

Other information

- each tablespoon (15 mL) contains: calcium 20 mg, magnesium 595 mg, potassium 2 mg and sodium 3 mg
- store at 20-25°C (68-77°F)
- keep tightly closed
- does not meet USP requirements for preservative effectiveness

Inactive ingredients

anhydrous citric acid, D&C red #28, flavor, glycerin, purified water, sorbitol solution, sucrose, xanthan gum

Questions or comments?

1-800-719-9260

Principal Display Panel

Compare to Phillips'® Milk of Magnesia active ingredient

Milk of Magnesia

SALINE LAXATIVE

Magnesium Hydroxide

Comfortable, Cramp Free Relief

Stimulant Free

Wild Cherry Flavor

ALCOHOL 0.4%

12 FL OZ (355 mL)

Compare to **Phillips'® Milk of Magnesia**
active ingredient

NDC 42507-949-40

HyVee® Milk of Magnesia

SALINE LAXATIVE

Magnesium Hydroxide

- Comfortable, Cramp Free Relief
Stimulant Free



Wild Cherry Flavor

ALCOHOL 0.4%

12 FL OZ (355 mL)

: 94940 2H F6

Drug Facts

**DO NOT USE IF PRINTED NECKBAND IS
BROKEN OR MISSING.**

Active ingredient (in each 15 mL TBSP) Magnesium hydroxide 1200 mg.....**Purpose** Saline laxative

Uses ■ relieves occasional constipation (irregularity)
■ generally produces bowel movement in ½ to 6 hours

Warnings

Ask a doctor before use if you have

- kidney disease ■ a magnesium-restricted diet
- stomach pain, nausea, or vomiting
- a sudden change in bowel habits that lasts more than 2 weeks

Ask a doctor or pharmacist before use if you are taking any other drug.
Take this product two or more hours before or after other drugs.
Laxatives may affect how other drugs work.

Stop use and ask a doctor if

- you have rectal bleeding or no bowel movement after using this product. These could be signs of a serious condition.
- you need to use a laxative for more than 1 week

If pregnant or breast-feeding, ask a health professional before use.
Keep out of reach of children. In case of overdose, get medical help or contact a Poison Control Center right away. (1-800-222-1222)

Directions

- shake well before use
 - do not exceed the maximum recommended daily dose in a 24 hour period
 - dose may be taken once a day preferably at bedtime, in divided doses, or as directed by a doctor
 - drink a full glass (8 oz) of liquid with each dose
 - for accurate dosing, use dose cup provided ■ TBSP = tablespoonful
- | | |
|--|--------------|
| adults and children 12 years and older | 2 to 4 TBSP. |
| children 6 to 11 years | 1 to 2 TBSP. |
| children under 6 years | ask a doctor |

Other information

- each tablespoon (15 mL) contains: calcium 20 mg, magnesium 595 mg, potassium 2 mg and sodium 3 mg
- does not meet USP requirements for preservative effectiveness
- store at 20-25°C (68-77°F) ■ keep tightly closed

Inactive ingredients anhydrous citric acid, D&C red #28, flavor, glycerin, purified water, sorbitol solution, sucrose, xanthan gum

Questions or comments? 1-800-719-9260

98 H2 0h6h6 :

DISTRIBUTED BY HY-VEE, INC.
5820 WESTOWN PARKWAY
WEST DES MOINES, IA 50266
SATISFACTION GUARANTEED
OR PURCHASE PRICE REFUNDED
FOR INFORMATION CALL 1-800-289-8343
www.hy-vee.com

GLUTEN FREE



MILK OF MAGNESIA

magnesium hydroxide liquid

Product Information

Product Type

HUMAN OTC DRUG LABEL

Item Code (Source)

NDC:42507-949

Route of Administration	ORAL	DEA Schedule		
Active Ingredient/Active Moiety				
Ingredient Name	Basis of Strength	Strength		
MAGNESIUM HYDRO XIDE (MAGNESIUM CATION and HYDROXIDE ION)	MAGNESIUM HYDROXIDE	1200 mg in 15 mL		
Inactive Ingredients				
Ingredient Name	Strength			
ANHYDROUS CITRIC ACID				
GLYCERIN				
WATER				
SORBITOL				
SUCROSE				
XANTHAN GUM				
D&C RED NO. 28				
Product Characteristics				
Color	PINK (Opaque Light pink)	Score		
Shape		Size		
Flavor	CHERRY (wild cherry)	Imprint Code		
Contains				
Packaging				
#	Item Code	Package Description	Marketing Start Date	Marketing End Date
1	NDC:42507-949-40	355 mL in 1 BOTTLE		
Marketing Information				
Marketing Category	Application Number or Monograph Citation	Marketing Start Date	Marketing End Date	
OTC monograph not final	part334	07/18/2005		

Labeler - HyVee Inc (006925671)

Revised: 4/2014

HyVee Inc