

## **HEADACHE RELIEF EXTRA STRENGTH- acetaminophen, aspirin, caffeine tablet AmerisourceBergen (Good Neighbor Pharmacy) 46122**

*Disclaimer: Most OTC drugs are not reviewed and approved by FDA, however they may be marketed if they comply with applicable regulations and policies. FDA has not evaluated whether this product complies.*

-----

### **Drug Facts**

#### **Active ingredients (in each caplet)**

**Acetaminophen 250 mg**

**Aspirin 250 mg (NSAID)\***

Caffeine 65 mg

\*nonsteroidal anti-inflammatory drug

### **Purposes**

**Pain reliever**

**Pain reliever**

Pain reliever aid

### **Uses**

- temporarily relieves minor aches and pains due to:
  - headache
  - a cold
  - arthritis
  - muscular aches
  - toothache
  - premenstrual and menstrual cramps

### **Warnings**

**Reye's syndrome:** Children and teenagers who have or are recovering from chicken pox or flu-like symptoms should not use this product. When using this product, if changes in behavior with nausea and vomiting occur, consult a doctor because these symptoms could be an early sign of Reye's syndrome, a rare but serious illness

**Allergy alert:** Acetaminophen may cause severe skin reactions. Symptoms may include:

- skin reddening
- blisters
- rash

If a skin reaction occurs, stop use and seek medical help right away.

**Allergy alert:** Aspirin may cause a severe allergic reaction, which may include:

- hives
- facial swelling
- shock
- asthma (wheezing)

**Liver warning:** This product contains acetaminophen. Severe liver damage may occur if you take:

- more than 8 caplets in 24 hours, which is the maximum daily amount
- with other drugs containing acetaminophen
- 3 or more alcoholic drinks every day while using this product

**Stomach bleeding warning:** This product contains an NSAID, which may cause severe stomach bleeding. The chance is higher if you:

- are age 60 or older
- have had stomach ulcers or bleeding problems
- take a blood thinning (anticoagulant) or steroid drug
- take other drugs containing prescription or nonprescription NSAIDs (aspirin, ibuprofen, naproxen, or others)
- have 3 or more alcoholic drinks every day while using this product
- take more or for a longer time than directed

**Caffeine warning:** The recommended dose of this product contains about as much caffeine as a cup of coffee. Limit the use of caffeine-containing medications, foods, or beverages while taking this product because too much caffeine may cause nervousness, irritability, sleeplessness, and, occasionally, rapid heartbeat.

### **Do not use**

- if you have ever had an allergic reaction to acetaminophen, aspirin or any other pain reliever/fever reducer
- with any other drug containing acetaminophen (prescription or nonprescription). If you are not sure whether a drug contains acetaminophen, ask a doctor or pharmacist.

### **Ask a doctor before use if**

- you have liver disease
- stomach bleeding warning applies to you
- you have asthma
- you are taking a diuretic
- you have a history of stomach problems, such as heartburn
- you have high blood pressure, heart disease, liver cirrhosis, or kidney disease

### **Ask a doctor or pharmacist before use if you are**

- taking a prescription drug for diabetes, gout, or arthritis
- taking any other drug or are under a doctor's care for any serious condition

### **Stop use and ask doctor if**

- an allergic reaction occurs. Seek medical help right away.

- any new symptoms appear
- you experience any of the following signs of stomach bleeding
  - feel faint
  - vomit blood
  - have bloody or black stools
  - have stomach pain that does not get better
- ringing in the ears or loss of hearing occurs
- pain gets worse or lasts for more than 10 days
- fever gets worse or lasts for more than 3 days
- painful area is red or swollen

These could be signs of a serious condition.

### **If pregnant or breast-feeding,**

ask a health professional before use. It is especially important not to use aspirin during the last 3 months of pregnancy unless definitely directed to do so by a doctor because it may cause problems in the unborn child or complications during delivery.

### **Keep out of reach of children.**

In case of overdose, get medical help or contact a Poison Control Center right away. Quick medical attention is critical for adults as well as for children even if you do not notice any signs or symptoms.

### **Directions**

- **do not use more than directed**
- drink a full glass of water with each dose
- adults and children 12 years and over: take 2 caplets every 6 hours; do not take more than 8 caplets in 24 hours
- children under 12 years: do not use unless directed by a doctor

### **Other information**

- store between 20°-25°C (68°-77°F)

### **Inactive ingredients**

carnauba wax\*\*, colloidal silicon dioxide, corn starch, hypromellose, microcrystalline cellulose, mineral oil\*\*, polyethylene glycol\*\*, povidone, sodium starch glycolate\*\*, stearic acid, talc\*\*, titanium dioxide

\*\*contains one or more of these ingredients

### **Questions or comments?**

Call **1-877-753-3935 Monday-Friday 9AM-5PM EST**

## **Principal Display Panel**

Compare to Excedrin® Extra Strength active ingredients†

Extra Strength

### **Headache Relief**

Acetaminophen, 250 mg

Aspirin, 250 mg, (NSAID),

Caffeine, 65 mg

Pain Reliever / Pain Reliever Aid

CAPLETS

†This product is not manufactured or distributed by GSK Consumer Healthcare distributor of Excedrin® Extra Strength.

**TAMPER EVIDENT: DO NOT USE IF PRINTED SAFETY SEAL UNDER CAP IS BROKEN OR MISSING.**

**KEEP OUTER CARTON FOR COMPLETE WARNINGS AND PRODUCT INFORMATION.**

Distributed by AmerisourceBergen

1300 Morris Drive, Chesterbrook, PA 19087

Questions or comments?

Visit us at [www.mygnp.com](http://www.mygnp.com)

## **Package Label**

Lot / Exp.:  
Product of:

PLD-E1036 F0003700



ABC# 10168088

Distributed By AmerisourceBergen  
1300 Morris Drive, Chesletbrook, PA 19067  
Questions or Concerns?  
Visit us at [www.mygrip.com](http://www.mygrip.com)



## Drug Facts

### Active ingredients (in each caplet)

Acetaminophen 250 mg ..... Pain reliever  
Aspirin 250 mg (NSAID)\* ..... Pain reliever  
Caffeine 65 mg ..... Pain reliever aid  
\*nonsteroidal anti-inflammatory drug

### Uses

- temporarily relieves minor aches and pains due to
  - headache
  - a cold
  - muscular aches
  - toothache
- premenstrual and menstrual cramps

### Warnings

**Reye's syndrome:** Children and teenagers who have or are recovering from chicken pox or flu-like symptoms should not use this product. When using this product, if changes in behavior with nausea and vomiting occur, consult a doctor because these symptoms could be an early sign of Reye's syndrome, a rare but serious illness.

**Allergy alert:** Aspirin may cause a severe allergic reaction, which may include:
 

- hives
- facial swelling
- shock
- asthma (wheezing)

**Allergy alert:** Acetaminophen may cause severe skin reactions. Symptoms may include:
 

- skin redness
- skin peeling
- blisters

**Liver warning:** This product contains acetaminophen. Severe liver damage may occur if you take:
 

- more than 8 caplets in 24 hours, which is the maximum daily amount

with other drugs containing acetaminophen

3 or more alcoholic drinks every day while using this product

**Stomach bleeding warning:** This product contains an NSAID, which may cause severe stomach bleeding. The chance is higher if you:
 

- are age 60 or older
- have had stomach ulcers or bleeding problems
- take a blood thinning (anticoagulant) or steroid drug
- take other drugs containing prescription or nonprescription NSAIDs (aspirin, ibuprofen, naproxen, or others)
- have 3 or more alcoholic drinks every day while using this product
- take more or for a longer time than directed

## Drug Facts (continued)

**Keep out of reach of children.** In case of overdose, get medical help or contact a Poison Control Center right away. Quick medical attention is critical for adults as well as for children even if you do not notice any signs or symptoms.

### Directions

- do not use more than directed
- drink a full glass of water with each dose
- adults and children 12 years and over: take 2 caplets every 6 hours; do not take more than 8 caplets in 24 hours
- children under 12 years: do not use unless directed by a doctor

### Other information

- store between 20°-25°C (68°-77°F)

**Inactive ingredients** carnauba wax\*\*, colloidal silicon dioxide, corn starch, hypromellose, microcrystalline cellulose, mineral oil\*\*, polyethylene glycol\*\*, povidone, sodium starch glycolate\*\*, stearic acid, talc\*\*, titanium dioxide\*\*

\*\*contains one or more of these ingredients

### Questions or comments?

Call 1-877-753-3835 Monday-Friday 9AM-5PM EST

\*This product is not manufactured or distributed by GSK Consumer Healthcare, distributor of Excedrin® Extra Strength.

TAMPER EVIDENT DO NOT USE IF PRINTED SAFETY SEAL UNDER CAP IS BROKEN OR MISSING.

KEEP OUTER CARTON FOR COMPLETE WARNINGS AND PRODUCT INFORMATION.



Compare to Excedrin® Extra Strength active ingredients\*

NDC 46122-382-78

Extra Strength

# Headache Relief

acetaminophen, 250 mg

aspirin, 250 mg (NSAID), caffeine, 65 mg

Pain Reliever/Pain Reliever Aid

100 caplets



**GOOD NEIGHBOR PHARMACY Extra Strength Headache Relief****HEADACHE RELIEF EXTRA STRENGTH**

acetaminophen, aspirin, caffeine tablet

**Product Information**

<b>Product Type</b>	HUMAN OTC DRUG	<b>Item Code (Source)</b>	NDC:46122-382
<b>Route of Administration</b>	ORAL		

**Active Ingredient/Active Moiety**

<b>Ingredient Name</b>	<b>Basis of Strength</b>	<b>Strength</b>
<b>ACETAMINOPHEN</b> (UNII: 362O9ITL9D) (ACETAMINOPHEN - UNII:362O9ITL9D)	ACETAMINOPHEN	250 mg
<b>ASPIRIN</b> (UNII: R16CO5Y76E) (ASPIRIN - UNII:R16CO5Y76E)	ASPIRIN	250 mg
<b>CAFFEINE</b> (UNII: 3G6A5W338E) (CAFFEINE - UNII:3G6A5W338E)	CAFFEINE	65 mg

**Inactive Ingredients**

<b>Ingredient Name</b>	<b>Strength</b>
<b>CARNAUBA WAX</b> (UNII: R12CBM0EIZ)	
<b>SILICON DIOXIDE</b> (UNII: ETJ7Z6XBU4)	
<b>STARCH, CORN</b> (UNII: O8232NY3SJ)	
<b>HYPROMELLOSES</b> (UNII: 3NXW29V3WO)	
<b>CELLULOSE, MICROCRYSTALLINE</b> (UNII: OP1R32D61U)	
<b>MINERAL OIL</b> (UNII: T5L8T28FGP)	
<b>POVIDONE</b> (UNII: FZ989GH94E)	
<b>SODIUM STARCH GLYCOLATE TYPE A CORN</b> (UNII: AG9B65PV6B)	
<b>STEARIC ACID</b> (UNII: 4ELV7Z65AP)	
<b>TALC</b> (UNII: 7SEV7J4R1U)	
<b>TITANIUM DIOXIDE</b> (UNII: 15FIX9V2JP)	
<b>POLYETHYLENE GLYCOL, UNSPECIFIED</b> (UNII: 3WJQ0SDW1A)	

**Product Characteristics**

<b>Color</b>	white	<b>Score</b>	no score
<b>Shape</b>	CAPSULE	<b>Size</b>	17mm
<b>Flavor</b>		<b>Imprint Code</b>	TCL370;S53
<b>Contains</b>			

**Packaging**

<b>#</b>	<b>Item Code</b>	<b>Package Description</b>	<b>Marketing Start Date</b>	<b>Marketing End Date</b>
1	NDC:46122-382-78	1 in 1 BOX	09/30/2016	09/30/2024
1		100 in 1 BOTTLE, PLASTIC; Type 0: Not a Combination Product		

## Marketing Information

Marketing Category	Application Number or Monograph Citation	Marketing Start Date	Marketing End Date
OTC monograph not final	part343	09/30/2016	09/30/2024

**Labeler** - AmerisourceBergen (Good Neighbor Pharmacy) 46122 (007914906)

Revised: 7/2023

AmerisourceBergen (Good Neighbor Pharmacy) 46122