

**REGNECY POST-MILKING BARRIER TEAT DIP- phenol teat dip solution**  
**ABS Corporation**

*Disclaimer: This drug has not been found by FDA to be safe and effective, and this labeling has not been approved by FDA. For further information about unapproved drugs, click [here](#).*

-----

**Regency Post-Milking Barrier Teat Dip**

**ACTIVE INGREDIENTS:**

<b>Active Ingredients</b>	<b>Purpose</b>
PHENOL (1.56% w/v)	Disinfectant
PHENOLATE SODIUM (0.06% w/v)	Disinfectant

**PURPOSE:**

<b>Active Ingredients</b>	<b>Purpose</b>
PHENOL (1.56% w/v)	Disinfectant
PHENOLATE SODIUM (0.06% w/v)	Disinfectant

**Keep Out of Reach of Children.**

If swallowed, get medical help or contact Poison Control Center right away.

**Uses:**

**FOR ANIMAL USE ONLY**

Aids in controlling mastitis organisms with a softening effect on teats

**Directions for Use:**

Before milking, wash udder and teats with a damp cloth.

Immediately after milking, dip each teat 1 to 2 inches into Regency Teat Dip.

Allow each teat to air dry. Do not wipe.

During the winter months, keep cows inside until the teats have completely dried to prevent freezing and chapping.

When a cow is being dried out, dip each teat in Regency Post-Milking Barrier Teat Dip once a day for 3 to 4 days after the last milking.

**CAUTION:**

**Avoid eye contact. Do not inhale fumes. For external use only.**

**FIRST AID:**

**For Ingestion - Immediately give castor oil or vegetable oil (15-30cc). Get medical attention immediately.**

**Be ready to induce vomiting at the advice of a physician or poison control center.**

**For Eyes - Immediately flush eyes with water for at least 15 minutes. Wash affected skin with water.**

**When using this product:**

Do Not Store in Direct Sunlight.

Avoid Excessive Heat.

**USER SAFETY WARNINGS:**

Do Not Allow Product To Freeze. If frozen, allow the product to completely thaw in a warm room.

Shake and thoroughly mix product before using.


Rinse teat dip cup with water after each milking and invert to dry.

Never pour used teat dip into original container.

Keep cap on container when not in use.

**INACTIVE INGREDIENTS:**

Water, Glycerin, Linear Alkylbenzene Sulfonate, Sodium Borate, Xanthan Gum, Fragrance, FD&C Blue #1



Aids in controlling mastitis organisms with a softening effect on the teats.  
**1.56% (w/v) Phenol with 0.06% (w/v) Sodium Phenate**  
Teat Dip Will Not Burn, Dry, Irritate Or Chap Teats. Contains a special blend of emollients and phenol.  
NOTE: Store at room temperature away from direct sunlight.

FOR ANIMAL USE ONLY  
KEEP OUT OF REACH OF CHILDREN  
DO NOT STORE IN DIRECT SUNLIGHT  
AVOID EXCESSIVE HEAT



**NET CONTENTS: 270 U.S. Gallons**

**FIRST AID:** For Ingestion - Immediately give castor oil or vegetable oil (15 - 30cc). Get medical attention immediately. Be ready to induce vomiting at the advice of a physician or poison control center. For Eyes - Immediately flush eyes with water for at least 15 minutes. Wash affected skin with water.

**CAUTION:** Avoid eye contact. For external use only.

**DIRECTIONS:** Before milking, wash udder and teats with damp cloth. Immediately after milking, dip teat entirely into Regency® Masticide® Post-Milking Barrier Teat Dip. Allow each teat to air dry. Do not wipe. During the winter months, keep cows inside until the teats have completely dried to prevent freezing and chapping.

If the solution in the dip cup becomes contaminated, discard and use fresh solution. Rinse teat dip cup with water after each milking and invert to dry. Never pour used teat dip into the original container. Keep cap on container when not in use.

When a cow is being dried out, dip each teat in Regency® Masticide® Post-Milking Barrier Teat Dip once a day for 3 to 4 days after the last milking.

Do not allow product to freeze. If frozen, allow the product to completely thaw in a warm room. Shake and thoroughly mix product before using.

**INGREDIENTS:** Water, Glycerin, Phenol, LAS (Linear Alkylbenzene Sulfonate), Sodium Borate, Xanthan Gum, Fragrance, FD&C Blue #1.

Distributed Exclusively By:  
**CLSolutions, LLC™**  
7031 North 16th St.  
Omaha, NE 68112  
(402) 502-3958  
www.clsag.com  
CLS-B-REV-5

Sporicidin® is a registered trademark of Contec, Inc.  
United States Patent # US6,586,477 B1

**LOT NO.** \_\_\_\_\_

**Regency** Post-Milking Barrier Teat Dip

Masticide® with **Sporicidin™ Technology**. "Ready To Use Barrier Teat Dip"

Aids in controlling mastitis organisms with a softening effect on the teats.

**1.56% (w/v) Phenol with 0.06% (w/v) Sodium Phenate**

Teat Dip Will Not Burn, Dry, Irritate or Chap Teats. Contains a special blend of emollients and phenol.

**NET CONTENTS: 270 U.S. Gallon**

# Regency<sup>®</sup>

## Masticide<sup>®</sup>

"Ready To Use Barrier Teat Dip"

Post-Milking Barrier Teat Dip  
(10% Glycerin)



Aids in controlling mastitis organisms with a softening effect on the teats.

**1.56% (w/v) Phenol with 0.06% (w/v) Sodium Phenate**  
Teat Dip Will Not Burn, Dry, Irritate Or Chap Teats. Contains a special blend of emollients and phenol.

NOTE: Store at room temperature away from direct sunlight.

FOR ANIMAL USE ONLY  
KEEP OUT OF REACH OF CHILDREN  
DO NOT STORE IN DIRECT SUNLIGHT  
AVOID EXCESSIVE HEAT



NET CONTENTS: 55 U.S. Gallons (208.2 L)

**FIRST AID:** For Ingestion - Immediately give castor oil or vegetable oil (15 - 30cc). Get medical attention immediately. Be ready to induce vomiting at the advice of a physician or poison control center. For Eyes - Immediately flush eyes with water for at least 15 minutes. Wash affected skin with water.

**CAUTION:** Avoid eye contact. For external use only.

**DIRECTIONS:** Before milking, wash udder and teats with damp cloth. Immediately after milking, dip teat entirely into Regency<sup>®</sup> Masticide<sup>®</sup> Post-Milking Barrier Teat Dip. Allow each teat to air dry. Do not wipe. During the winter months, keep cows inside until the teats have completely dried to prevent freezing and chapping.

If the solution in the dip cup becomes contaminated, discard and use fresh solution. Rinse teat dip cup with water after each milking and invert to dry. Never pour used teat dip into the original container. Keep cap on container when not in use.


When a cow is being dried out, dip each teat in Regency<sup>®</sup> Masticide<sup>®</sup> Post-Milking Barrier Teat Dip once a day for 3 to 4 days after the last milking.

Do not allow product to freeze. If frozen, allow the product to completely thaw in a warm room. Shake and thoroughly mix product before using.

**INGREDIENTS:** Water, Glycerin, Phenol, LAS (Linear Alkylbenzene Sulfonate), Sodium Borate, Xanthan Gum, Fragrance, FD&C Blue #1.

Distributed Exclusively By:  
CLSolutions, LLC  
PO BOX 837  
Fulton, TX 78358  
(361) 729-3993  
www.CoastalLivestock.com

Sporicidin<sup>®</sup> is a registered trademark of Contec, Inc.  
United States Patent # US6,586,477 B1  
CLS-B-55GAL



## Regency Post-Milking Barrier Teat Dip

Masticide<sup>®</sup> with **Sporicidin<sup>™</sup> Technology**. "Ready To Use Barrier Teat Dip"

Aids in controlling mastitis organisms with a softening effect on the teats.

**1.56% (w/v) Phenol with 0.06% (w/v) Sodium Phenate**

Teat Dip Will Not Burn, Dry, Irritate or Chap Teats. Contains a special blend of emollients and phenol.

NET CONTENTS: 55 U.S. Gallon (208.2 L)

# Regency<sup>®</sup>

## Masticide<sup>®</sup>

"Ready To Use Barrier Teat Dip"

Post-Milking Barrier Teat Dip  
(10% Glycerin)



Aids in controlling mastitis organisms with a softening effect on the teats.

**1.56% (w/v) Phenol with 0.06% (w/v) Sodium Phenate**  
Teat Dip Will Not Burn, Dry, Irritate Or Chap Teats. Contains a special blend of emollients and phenol.

NOTE: Store at room temperature away from direct sunlight.

FOR ANIMAL USE ONLY  
KEEP OUT OF REACH OF CHILDREN  
DO NOT STORE IN DIRECT SUNLIGHT  
AVOID EXCESSIVE HEAT



NET CONTENTS: 5 U.S. Gallons (18.927 L)

**FIRST AID:** For Ingestion - Immediately give castor oil or vegetable oil (15 - 30cc). Get medical attention immediately. Be ready to induce vomiting at the advice of a physician or poison control center. For Eyes - Immediately flush eyes with water for at least 15 minutes. Wash affected skin with water.

**CAUTION:** Avoid eye contact. For external use only.

**DIRECTIONS:** Before milking, wash udder and teats with damp cloth. Immediately after milking, dip teat entirely into Regency<sup>®</sup> Masticide<sup>®</sup> Post-Milking Barrier Teat Dip. Allow each teat to air dry. Do not wipe. During the winter months, keep cows inside until the teats have completely dried to prevent freezing and chapping.

If the solution in the dip cup becomes contaminated, discard and use fresh solution. Rinse teat dip cup with water after each milking and invert to dry. Never pour used teat dip into the original container. Keep cap on container when not in use.

When a cow is being dried out, dip each teat in Regency<sup>®</sup> Masticide<sup>®</sup> Post-Milking Barrier Teat Dip once a day for 3 to 4 days after the last milking.

Do not allow product to freeze. If frozen, allow the product to completely thaw in a warm room. Shake and thoroughly mix product before using.

**INGREDIENTS:** Water, Glycerin, Phenol, LAS (Linear Alkylbenzene Sulfonate), Sodium Borate, Xanthan Gum, Fragrance, FD&C Blue #1.

Distributed Exclusively By:  
CLSolutions, LLC  
PO BOX 837  
Fulton, TX 78358  
(361) 729-3993  
www.CoastalLivestock.com

Sporicidin<sup>®</sup> is a registered trademark of Contec, Inc.  
United States Patent # US6,586,477 B1  
CLS-B-55GAL



## Regency Post-Milking Barrier Teat Dip

Masticide<sup>®</sup> with **Sporicidin<sup>™</sup> Technology**. "Ready To Use Barrier Teat Dip"

Aids in controlling mastitis organisms with a softening effect on the teats.

**1.56% (w/v) Phenol with 0.06% (w/v) Sodium Phenate**

Teat Dip Will Not Burn, Dry, Irritate or Chap Teats. Contains a special blend of emollients and phenol.  
**NET CONTENTS: 5 U.S. Gallon ( 18.927 L)**



**Post-Milking Barrier Teat Dip**  
*(10% Glycerin)*



Aids in controlling mastitis organisms with a softening effect on the teats.

**1.56% (w/v) Phenol with 0.06% (w/v) Sodium Phenate**

Teat Dip Will Not Burn, Dry, Irritate Or Chap Teats. Contains a special blend of emollients and phenol.

NOTE: Store at room temperature away from direct sunlight.

**NET CONTENTS: 1 U.S. Gallon (3.785 L)**

FOR ANIMAL USE ONLY  
 KEEP OUT OF REACH OF CHILDREN  
 DO NOT STORE IN DIRECT SUNLIGHT  
 AVOID EXCESSIVE HEAT



**FIRST AID:** For Ingestion - Immediately give castor oil or vegetable oil (15 - 30cc). Get medical attention immediately. Be ready to induce vomiting at the advice of a physician or poison control center. For Eyes - Immediately flush eyes with water for at least 15 minutes. Wash affected skin with water.

**CAUTION:** Avoid eye contact. For external use only.

**DIRECTIONS:** Before milking, wash udder and teats with damp cloth. Immediately after milking, dip teat entirely into Regency® Masticide® Post-Milking Barrier Teat Dip. Allow each teat to air dry. Do not wipe. During the winter months, keep cows inside until the teats have completely dried to prevent freezing and chapping.

If the solution in the dip cup becomes contaminated, discard and use fresh solution. Rinse teat dip cup with water after each milking and invert to dry. Never pour used teat dip into the original container. Keep cap on container when not in use.

When a cow is being dried out, dip each teat in Regency® Masticide® Post-Milking Barrier Teat Dip once a day for 3 to 4 days after the last milking.

Do not allow product to freeze. If frozen, allow the product to completely thaw in a warm room. Shake and thoroughly mix product before using.

**INGREDIENTS:** Water, Glycerin, Phenol, LAS (Linear Alkylbenzene Sulfonate), Sodium Borate, Xanthan Gum, Fragrance, FD&C Blue #1.

Distributed Exclusively By:

CLSolutions, LLC  
 PO BOX 837  
 Fulton, TX 78358  
 (361) 729-3993

www.CoastalLivestock.com

Sporicidin® is a registered trademark of Contec, Inc.  
 United States Patent # US6,586,477 B1

CLS-B-1GAL



### Regency Post-Milking Barrier Teat Dip

Masticide® with **Sporicidin™ Technology**. "Ready To Use Barrier Teat Dip"

Aids in controlling mastitis organisms with a softening effect on the teats.

**1.56 % (w/v) Phenol with 0.06 % (w/v) Sodium Phenate**

Teat Dip Will Not Burn, Dry, Irritate or Chap Teats. Contains a special blend of emollients and phenol.

**NET CONTENTS: 1 U.S. Gallon (3.785 L)**

## REGNECY POST-MILKING BARRIER TEAT DIP

phenol teat dip solution

### Product Information

<b>Product Type</b>	OTC ANIMAL DRUG	<b>Item Code (Source)</b>	NDC:18472-100
<b>Route of Administration</b>	TOPICAL		

### Active Ingredient/Active Moiety

Ingredient Name	Basis of Strength	Strength
PHENOL (UNII: 339NCG44TV) (PHENOL - UNII:339NCG44TV)	PHENOL	156 g in 1 L
PHENOLATE SODIUM (UNII: 4NC0T56V35) (PHENOL - UNII:339NCG44TV)	PHENOLATE SODIUM	0.06 g in 1 L

### Inactive Ingredients

Ingredient Name	Strength
SODIUM HYDROXIDE (UNII: 55X04QC32I)	
SODIUM BORATE (UNII: 91MBZ8H3QO)	

<b>GLYCERIN</b> (UNII: PDC6A3C0OX)	
<b>XANTHAN GUM</b> (UNII: TTV12P4NEE)	
<b>P-DODECYLBENZENESULFONIC ACID</b> (UNII: OC21S23N1O)	
<b>FD&amp;C BLUE NO. 1</b> (UNII: HBR47K3TBD)	
<b>WATER</b> (UNII: 059QF0KO0R)	

### Packaging

#	Item Code	Package Description	Marketing Start Date	Marketing End Date
1	NDC:18472-100-03	1021.95 L in 1 CONTAINER		
2	NDC:18472-100-04	3.785 L in 1 JUG		
3	NDC:18472-100-05	18.925 L in 1 PAIL		
4	NDC:18472-100-06	208.175 L in 1 DRUM		

### Marketing Information

Marketing Category	Application Number or Monograph Citation	Marketing Start Date	Marketing End Date
unapproved drug other		06/13/2011	

**Labeler** - ABS Corporation (048022727)

**Registrant** - ABS Corporation (048022727)

### Establishment

Name	Address	ID/FEI	Business Operations
ABS Corporation		048022727	api manufacture, pack, label