## SINECCH- arnica montana capsule Hahnemann Laboratories, Inc. dba Alpine Pharmaceuticals

Disclaimer: This homeopathic product has not been evaluated by the Food and Drug Administration for safety or efficacy. FDA is not aware of scientific evidence to support homeopathy as effective.

-----

# **Active Ingredient**

Active Ingredients: SinEcch 01 500mg Arnica Montana 1M, SinEcch 02 500mg Arnica Montana 12C

# **Purpose**

surgical recovery regimen

Reduces Pain, Bruising, and Swelling after Surgery or Major Trauma.

### Keep Out of Reach of Children

Keep this and all medication out of the reach of children.

#### Uses

Surgery or Major Trauma

## Warnings

Warnings: If you are pregnant or nursing, consult your physician before taking SinEcch. For multiple procedures during a 4 week interval, take only 1 SinEcch regimen with the final procedure.

Do not drink or eat anything for 15 minutes before and after taking SinEcch.

# **Drug Interactions**

Drug Interactions: Do not take together with Talwin tm or Talwin NX tm

#### Storage

Storage: Store at room temperature.

#### **Directions**

Directions: For Adults and children over 6. Swallow the capsule with water, or twist open the capsule and pour contents into your mouth. Let the small pellets dissolve.

SinEcch SURGERY MAJOR TRAUMA

Directions for Using the Facial Surgery, Liposuction, Major injuries from

Accidents,

Capsule Labeled: Body Contouring, or Falls, Sports Injuries, or when

	General Surgery	Starting after Surgery
PRE-OP (SinEcch 01	1-24 hours	As soon as possible after
Orange/Peach Color	before surgery	the surgery or trauma
POST-OP (SinEcch 01)	1-6 hours	4 hours after the
Orange/Peach Color	after surgery	first dose
BEDTIME (SinEcch 01)	Bedtime on the day of	Bedtime the same day as
Orange/Peach Color	surgery (Day 1)	the above doses (Day 1)
DAYS 2, 3, AND 4	1 capsule every morning	1 capsule every morning
(SinEcch 02)	1 capsule every afternoon	1 capsule every afternoon
Peach/Peach Color	1 capsule every evening	1 capsule every evening

#### **Inactive Ingredients**

Inactive Ingredients : Sucrose 400mg, Lactose 100 mg, Capsules contain Hypromellose, Red Iron Oxide, Titanium Dioxide

#### **Package Label**

SinEcch tm Homeopathic Arnica Montanan

surgical recovery regimen Professional Strength

Clinically Proven to Reduce Bruising and Swelling after Surgery

ALL NATURAL 12 capsules THIS PRODUCT IS SAFETY SEALED. THE BOX IS SEALED AND THE CAPSULES ARE INDIVIDUALLY BLISTER SEALED. DO NOT USE IF SEALS ARE BROKEN.

Printed with soy based inks.

Developed and distributed by Alpine Pharmaceuticals, San Rafael, CA 94901 a wholly owned subsidiary of Hahnemann Laboratories, Inc. All rights reserved. Made in the USA

Questions or comments call 888.SINECCH (888-746-3224) email: info@alpinepharm.com or visit

A Product of Alpine Pharmaceuticals The Arnica Experts! tm



# SINECCH

**Professional Strength** Homeopathic Arnica

surgical recovery regimen

Reduces Pain, Bruising, and Swelling after Surgery or Major Trauma.

0.		_		
SI	N			HM.
U	IN			

Directions for Using the Capsule Labeled:

PRE-OP (SinEcch 01) Orange/Peach Color

POST-OP (SinEcch 01) Orange/Peach Color

BEDTIME (SinEcch 01) Orange/Peach Color

DAYS 2, 3, AND 4 (SinEcch 02) Peach/Peach Color

# SURGERY

Facial Surgery, Liposuction, Body Contouring, or General Surgery

1 to 24 hours before surgery

1 to 6 hours after surgery

Bedtime on the day of surgery (Day 1)

1 capsule every morning 1 capsule every afternoon 1 capsule every evening

# **MAJOR TRAUMA**

Major injuries from Accidents, Falls, Sports Injuries, or when Starting after Surgery

As soon as possible after the surgery or trauma

4 hours after the first dose

Bedtime the same day as the above doses (Day 1)

1 capsule every morning 1 capsule every afternoon 1 capsule every evening

Directions: For Adults and children over 6. Swallow the capsule with water, or twist open the capsule and pour the contents into your mouth. Let the small pellets dissolve.

Active Ingredients: SinEcch 01 500mg Arnica Montana 1M, SinEcch 02 500mg Arnica Montana 12C Inactive Ingredients: Sucrose 400mg, Lactose 100mg. Capsules contain Hypromellose, Red Iron Oxide, Titanium Dioxide

Warnings: If you are pregnant or nursing, consult your physician before taking SinEcch. Keep this and all medication out of the reach of children. For multiple procedures during a 4 week interval, take only 1 SinEcch regimen with final procedure. Storage: Store at room temperature.

Drug Interactions: Do not take together with

Talwin™ or Talwin NX™. Do not eat or drink anything for 15 minutes before and after taking SinEcch.



The Arnica Experts!"

#### **SINECCH**

arnica montana capsule

#### **Product Information**

ı	Product Type	HUMAN OTC DRUG	Item Code (Source)	NDC:63545-142

Route of Administration ORAL

#### Active Ingredient/Active Moiety

Ingredient Name	Basis of Strength	Strength
APNICA MONTANA (UNII: 080TY2087W) (APNICA MONTANA - UNII: 080TY2087W)	ΔΡΝΙΟΔ ΜΟΝΤΔΝΔ	1 [hp M]

Inactive .	Ingre	dients
------------	-------	--------

	Ingredient Name	Strength
SI	UCRO SE (UNII: C151H8 M554)	

LACTOSE (UNII: J2B2A4N98G)	
HYPROMELLOSES (UNII: 3NXW29V3WO)	
FERRIC OXIDE RED (UNII: 1K09F3G675)	
TITANIUM DIO XIDE (UNII: 15FIX9 V2JP)	

Product Characteristics				
Color	orange (orange/peach)	Score	no score	
Shape	CAPSULE (capsule)	Size	18 mm	
Flavor		Imprint Code	SinEcch01	
Contains				

P	ackaging			
#	Item Code	Package Description	<b>Marketing Start Date</b>	<b>Marketing End Date</b>
1	NDC:63545-142-01	12 in 1 BLISTER PACK; Type 0: Not a Combination Product	06/30/1998	

Marketing Information				
Marketing Category	Application Number or Monograph Citation	Marketing Start Date	Marketing End Date	
unapproved homeopathic		06/30/1998		

# Labeler - Hahnemann Laboratories, Inc. dba Alpine Pharmaceuticals (147098081)

# Registrant - Hahnemann Laboratories, Inc. dba Alpine Pharmaceuticals (147098081)

Establishment					
Name	Address	ID/FEI	<b>Business Operations</b>		
Hahnemann Laboratories, Inc. dba Alpine Pharmaceuticals		147098081	manufacture(63545-142)		

Revised: 9/2015 Hahnemann Laboratories, Inc. dba Alpine Pharmaceuticals