

LAXATIVE- bisacodyl tablet, delayed release
ARMY AND AIR FORCE EXCHANGE SERVICE

Exchange Select 44-676

Active ingredient (in each tablet)

Bisacodyl USP, 5 mg

Purpose

Stimulant laxative

Uses

- for relief of occasional constipation (irregularity)
- this product generally produces bowel movement in 6 to 12 hours

Warnings

Do not use

if you cannot swallow without chewing.

Ask a doctor before use if you have

- a sudden change in bowel habits that lasts more than 2 weeks
- stomach pain, nausea or vomiting

When using this product

- do not chew or crush tablet(s)
- do not use within 1 hour after taking an antacid or milk
- you may have stomach discomfort, faintness and cramps

Stop use and ask a doctor if

- you have rectal bleeding or fail to have a bowel movement after use of a laxative. These could be signs of a serious condition.
- you need to use a laxative for more than 1 week

If pregnant or breast-feeding,

ask a health professional before use.

Keep out of reach of children.

In case of overdose, get medical help or contact a Poison Control Center (1-800-222-1222) right away.

Directions

- take with a glass of water

adults and children 12 years and over	take 1 to 3 tablets in a single daily dose
children 6 to under 12 years	take 1 tablet in a single daily dose
children under 6 years	ask a doctor

Other information

- **TAMPER EVIDENT: DO NOT USE IF OUTER PACKAGE IS OPENED OR BLISTER IS TORN OR BROKEN**
- avoid excessive humidity
- store at 25°C (77°F); excursions permitted between 15°C-30°C (59°F-86°F)
- see end flap for expiration date and lot number

Inactive ingredients

carmine, colloidal silicon dioxide, corn starch, FD&C red #40 aluminum lake, hypromellose, lactose anhydrous, magnesium stearate, methacrylic acid, microcrystalline cellulose, polydextrose, polyethylene glycol, shellac wax, simethicone, sodium bicarbonate, sodium lauryl sulfate, stearic acid, talc, titanium dioxide, triacetin, triethyl citrate

Questions or comments?

1-800-426-9391

Principal Display Panel

exchange/**select**™

Women's Gentle

Laxative

Bisacodyl USP, 5 mg

STIMULANT LAXATIVE

Gentle, predictable
overnight relief

Actual Size

60 Tablets

Enteric Coated

quality
value

**TAMPER EVIDENT:
DO NOT USE IF
PACKAGE IS OPENED
OR IF BLISTER UNIT
IS TORN, BROKEN
OR SHOWS ANY
SIGNS OF
TAMPERING**

50844 REV0119A67611

"SATISFACTION GUARANTEED OR YOUR MONEY BACK"

Manufactured For Your Military Exchanges

Distributed by: LNK International, Inc., Hauppauge, NY 11788

1-800-426-9391

exchange **select**™

Women's
Gentle

Laxative

Bisacodyl USP, 5 mg

STIMULANT LAXATIVE

exchange **select**™

Women's
Gentle

Laxative

Bisacodyl USP, 5 mg

STIMULANT LAXATIVE

Gentle, predictable
overnight relief

60 Tablets
Enteric Coated

Actual Size

quality
value

exchange **select**™

Women's
Gentle

Laxative

Bisacodyl USP, 5 mg
STIMULANT LAXATIVE

TAMPER EVIDENT:
DO NOT USE IF
PACKAGE IS OPENED
OR IF BLISTER UNIT
IS TORN, BROKEN
OR SHOWS ANY
SIGNS OF
TAMPERING



SATISFACTION GUARANTEED OR YOUR MONEY BACK™

Manufactured by: Linc International, Inc.
Hempstead, NY 11788 1-800-426-9391

No Print Area
Lot & Exp Date

508444 REV0119A67611

Drug Facts (continued)
Inactive ingredients: carmine, colloidal silicon dioxide, corn starch, FD&C red #40 aluminum lake, hypromellose, lactose anhydrous, magnesium stearate, methacrylic acid, microcrystalline cellulose, polydextrose, polyethylene glycol, shellac wax, simethicone, sodium bicarbonate, sodium lauryl sulfate, stearic acid, talc, titanium dioxide, triacetin, triethyl citrate
Questions or comments? 1-800-426-9391

Drug Facts (continued)
Other information
TAMPER EVIDENT: DO NOT USE IF OUTER PACKAGE IS OPENED OR BLISTER IS TORN OR BROKEN
avoid excessive humidity
store at 25°C (77°F); excursions permitted between 15°-30°C (59°-86°F)
see end flap for expiration date and lot number

B-0066F-676A-11-R
REV0119A67611

Drug Facts (continued) Active ingredient Bisacodyl USP, 5 mg.....Stimulant laxative (in each tablet) Purpose ■ you have rectal bleeding or fail to have a bowel movement after use of a laxative. These could be signs of a serious condition. ■ you need to use a laxative for more than 1 week If pregnant or breast-feeding, ask a health professional before use. Keep out of reach of children. In case of overdose, get medical help or contact a Poison Control Center (1-800-222-1222) right away. Directions ■ take with a glass of water adults and children 12 years and over take 1 to 3 tablets in a single daily dose children 6 to under 12 years take 1 tablet in a single daily dose children under 6 years ask a doctor	
Drug Facts (continued) Uses ■ for relief of occasional constipation (irregularity) to 12 hours ■ this product generally produces bowel movement in 6 to 12 hours Warnings Do not use if you cannot swallow without chewing. Ask a doctor before use if you have ■ a sudden change in bowel habits that lasts more than 2 weeks ■ stomach pain, nausea or vomiting When using this product ■ do not chew or crush tablet(s) ■ do not use within 1 hour after taking an antacid or milk ■ you may have stomach discomfort, faintness and cramps	

E1048

Exchange Select 44-676A

LAXATIVE

bisacodyl tablet, delayed release

Product Information

Product Type	HUMAN OTC DRUG	Item Code (Source)	NDC:55301-676
Route of Administration	ORAL		

Active Ingredient/Active Moiety

Ingredient Name	Basis of Strength	Strength
BISACODYL (UNII: 10X0709Y6I) (DEACETYLBISACODYL - UNII:R09078E41Y)	BISACODYL	5 mg

Inactive Ingredients

Ingredient Name	Strength
SILICON DIOXIDE (UNII: ETJ7Z6XBU4)	
STARCH, CORN (UNII: O8232NY3SJ)	
FD&C RED NO. 40 (UNII: WZB9127XOA)	
HYPROMELLOSE, UNSPECIFIED (UNII: 3NXW29V3WO)	
ANHYDROUS LACTOSE (UNII: 3SY5LH9PMK)	
MAGNESIUM STEARATE (UNII: 70097M6I3O)	
METHACRYLIC ACID (UNII: 1CS02G8656)	
MICROCRYSTALLINE CELLULOSE (UNII: OP1R32D61U)	
POLYDEXTROSE (UNII: VH2XOU12IE)	
POLYETHYLENE GLYCOL, UNSPECIFIED (UNII: 3WJQ0SDW1A)	
SHELLAC (UNII: 46N107B71O)	
DIMETHICONE (UNII: 92RU3N3Y1O)	
WATER (UNII: 059QF0KO0R)	
SODIUM BICARBONATE (UNII: 8MDF5V39QO)	
SODIUM LAURYL SULFATE (UNII: 368GB5141J)	
STEARIC ACID (UNII: 4ELV7Z65AP)	
TALC (UNII: 7SEV7J4R1U)	
TITANIUM DIOXIDE (UNII: 15FIX9V2JP)	
TRIACETIN (UNII: XHX3C3X673)	
TRIETHYL CITRATE (UNII: 8Z96QXD6UM)	

Product Characteristics

Color	pink	Score	no score
Shape	ROUND	Size	8mm
Flavor		Imprint Code	B
Contains			

Packaging

#	Item Code	Package Description	Marketing Start Date	Marketing End Date
1	NDC:55301-676-11	4 in 1 CARTON	09/16/2015	

1	15 in 1 BLISTER PACK; Type 0: Not a Combination Product		
Marketing Information			
Marketing Category	Application Number or Monograph Citation	Marketing Start Date	Marketing End Date
OTC Monograph Drug	505G(a)(3)	09/16/2015	

Labeler - ARMY AND AIR FORCE EXCHANGE SERVICE (001695568)

Establishment

Name	Address	ID/FEI	Business Operations
LNK International, Inc.		038154464	pack(55301-676)

Establishment

Name	Address	ID/FEI	Business Operations
LNK International, Inc.		832867837	manufacture(55301-676) , pack(55301-676)

Establishment

Name	Address	ID/FEI	Business Operations
LNK International, Inc.		832867894	manufacture(55301-676)

Establishment

Name	Address	ID/FEI	Business Operations
LNK International, Inc.		967626305	pack(55301-676)

Establishment

Name	Address	ID/FEI	Business Operations
LNK International, Inc.		117025878	manufacture(55301-676)

Revised: 12/2023

ARMY AND AIR FORCE EXCHANGE SERVICE