

CIRCLE K COLD- acetaminophen, dextromethorphan hydrobromide, guaifenesin, and phenylephrine hydrochloride tablet
Lil' Drug Store Products, Inc

Disclaimer: Most OTC drugs are not reviewed and approved by FDA, however they may be marketed if they comply with applicable regulations and policies. FDA has not evaluated whether this product complies.

Circle K ® Cold

Drug Facts

Active ingredients (in each tablet)	Purpose
Acetaminophen 325 mg	Pain reliever/fever reducer
Dextromethorphan HBr 15 mg	Cough suppressant
Guaifenesin 200 mg	Expectorant
Phenylephrine HCl 5 mg	Nasal decongestant

Uses

- for the temporary relief of the following cold/flu symptoms:
 - minor aches and pains
 - headache
 - sore throat
 - nasal congestion
 - cough
 - impulse to cough
- helps loosen phlegm (mucus) and thin bronchial secretions to make coughs more productive
- temporarily reduces fever

Warnings

Liver warning

This product contains acetaminophen. Severe liver damage may occur if you take:

- more than 4,000 mg of acetaminophen in 24 hours
- with other drugs containing acetaminophen
- 3 or more alcoholic drinks every day while using this product.

Allergy alert

Acetaminophen may cause severe allergic reactions. Symptoms may include:

- skin reddening
- blisters
- rash

If a skin reaction occurs, stop use and seek medical help right away.

Sore throat warning

If sore throat is severe, persists for more than 2 days, is accompanied or followed by fever, headache, rash, nausea or vomiting, consult a doctor promptly.

A persistent cough may be a sign of a serious condition. If cough persists for more than 1 week, tends to recur, or is accompanied by fever, rash, or persistent headache, consult a doctor.

Do not use

- with any other drug containing acetaminophen (prescription or nonprescription). If you are not sure whether a drug contains acetaminophen, ask a doctor or pharmacist.
- if you are now taking a prescription monoamine oxidase inhibitor (MAOI) (certain drugs for depression, psychiatric or emotional conditions, or Parkinson's disease), or for 2 weeks after stopping the MAOI drug. If you do not know if your prescription drug contains an MAOI, ask a doctor or pharmacist before taking this product.

Ask a doctor before use if you have

- liver disease
- heart disease
- diabetes
- thyroid disease
- high blood pressure
- trouble urinating due to an enlarged prostate gland
- cough that occurs with too much phlegm (mucus)
- persistent or chronic cough such as occurs with smoking, asthma, chronic bronchitis or emphysema

Ask a doctor or pharmacist before use if you are

- taking the blood thinning drug warfarin

When using this product

- **do not exceed recommended dosage**

Stop use and ask a doctor if

- pain, nasal congestion or cough gets worse or lasts more than 7 days
- fever gets worse or lasts more than 3 days
- redness or swelling is present
- new symptoms occur
- cough comes back or is accompanied by a fever, rash or persistent headache
- nervousness, dizziness, or sleeplessness occur.

These could be signs of a serious condition.

If pregnant or breast-feeding, ask a health professional before use.

Keep out of reach of children. In case of overdose, get medical help or contact a Poison Control Center right away. (1-800-222-1222) Prompt medical attention is critical

for adults as well as for children even if you do not notice any signs or symptoms.

Directions

adults and children 12 years and over	<ul style="list-style-type: none">▪ take 2 tablets with water every 6-8 hours as needed▪ do not take more than 8 tablets in 24 hours, or as directed by a doctor
children under 12 years of age	ask a doctor

Other information

- store at room temperature 59-86° F (15-30° C)

Inactive ingredients

maltodextrin, microcrystalline cellulose, povidone, silicon dioxide ¹, sodium starch glycolate, starch, stearic acid

¹ may contain

Questions or comments?

call toll-free **1-877-507-6516 (M-F 8AM-4:30PM CST)**

Proudly distributed by Circle K Stores Inc.

PRINCIPAL DISPLAY PANEL - 4 Tablet Pouch Carton

CIRCLE K ®

Cold
severe
congestion

Acetaminophen, Dextromethorphan HBr, Guaifenesin and Phenylephrine HCl
Pain Reliever/Fever Reducer, Cough Suppressant, Expectorant, Nasal Decongestant

ACTUAL SIZE

Non-Drowsy Relief of
- Nasal Congestion
- Sore Throat
- Cough, Impulse to Cough

4 Tablets

2 POUCHES, 2 TABLETS PER POUCH

CIRCLE K

Cold severe congestion

Acetaminophen, Dextromethorphan HBr, Guaifenesin and Phenylephrine HCl
Pain Reliever/Fever Reducer, Cough Suppressant, Expectorant, Nasal Decongestant



Non-Drowsy Relief of

- Nasal Congestion
- Sore Throat
- Cough, Impulse to Cough

4 Tablets
2 POUCHES, 2 TABLETS PER POUCH

severe congestion

4 Tablets

CIRCLE K

DO NOT USE IF INNER POUCH OR CARTON IS OPEN OR DAMAGED!

Drug Facts

Active ingredients (in each tablet)	Purpose
Acetaminophen 325 mg	Pain reliever/fever reducer
Dextromethorphan HBr 15 mg	Cough suppressant
Guaifenesin 200 mg	Expectorant
Phenylephrine HCl 5 mg	Nasal decongestant

Uses

- for the temporary relief of the following cold/flu symptoms:
 - minor aches and pains
 - headache
 - sore throat
 - nasal congestion
 - cough
 - impulse to cough
- helps loosen phlegm (mucus) and thin bronchial secretions to make coughs more productive
- temporarily reduces fever

Warnings

Liver warning: This product contains acetaminophen. Severe liver damage may occur if you take: ■ more than 4,000 mg of acetaminophen in 24 hours ■ with other drugs containing acetaminophen ■ 3 or more alcoholic drinks every day while using this product.

Allergy alert: Acetaminophen may cause severe allergic reactions. Symptoms may include: ■ skin reddening ■ blisters ■ rash. If a skin reaction occurs, stop use and seek medical help right away.

Sore throat warning: If sore throat is severe, persists for more than 2 days, is accompanied or followed by fever, headache, rash, nausea or vomiting, consult a doctor promptly.

A persistent cough may be a sign of a serious condition. If cough persists for more than 1 week, tends to recur, or is accompanied by fever, rash, or persistent headache, consult a doctor.

LIFT PANEL FOR CONTINUED DRUG FACTS

Drug Facts (continued)

Do not use

- with any other drug containing acetaminophen (prescription or nonprescription). If you are not sure whether a drug contains acetaminophen, ask a doctor or pharmacist.
- if you are now taking a prescription monoamine oxidase inhibitor (MAOI) (certain drugs for depression, psychiatric or emotional conditions, or Parkinson's disease), or for 2 weeks after stopping the MAOI drug. If you do not know if your prescription drug contains an MAOI, ask a doctor or pharmacist before taking this product.

Ask a doctor before use if you have

- liver disease
- heart disease
- high blood pressure
- diabetes
- thyroid disease
- trouble urinating due to an enlarged prostate gland
- cough that occurs with too much phlegm (mucus)
- persistent or chronic cough such as occurs with smoking, asthma, chronic bronchitis or emphysema

Ask a doctor or pharmacist before use if you are

- taking the blood thinning drug warfarin

When using this product

- do not exceed recommended dosage

Stop use and ask a doctor if

- pain, nasal congestion or cough gets worse or lasts more than 7 days
- fever gets worse or lasts more than 3 days
- redness or swelling is present
- new symptoms occur
- cough comes back or is accompanied by a fever, rash or persistent headache
- nervousness, dizziness, or sleeplessness occur.

These could be signs of a serious condition.

severe congestion

CIRCLE K

severe congestion

4 Tablets

Drug Facts (continued)

If pregnant or breast-feeding, ask a health professional before use. **Keep out of reach of children.** In case of overdose, get medical help or contact a Poison Control Center right away. (1-800-222-1222) Prompt medical attention is critical for adults as well as for children even if you do not notice any signs or symptoms.

Directions

adults and children 12 years and over	■ take 2 tablets with water every 6-8 hours as needed
children under 12 years of age	■ do not take more than 8 tablets in 24 hours, or as directed by a doctor
	ask a doctor

Other information

- store at room temperature 59-86° F (15-30° C)

Inactive ingredients

maltodextrin, microcrystalline cellulose, povidone, silicon dioxide*, sodium starch glycolate, starch, stearic acid
*may contain

Questions or comments?
call toll-free 1-877-507-6516 (M-F 8AM-4:30PM CST)

Product manufactured for: Lill Drug Store Products, Inc., 9300 Earhart Lane SW Cedar Rapids, IA 52404 www.lilldrugstore.com 57282C-US-10-18
Proudly distributed by Circle K Stores Inc.

EXP
LOT





Principal Display Panel CK Cold, 6ct

TEMPORARY RELIEVES COLD/FLU SYMPTOMS

Compare to the active ingredients in

Tylenol® Cold & Flu Severe**

CIRCLE K

severe congestion

cold relief

Acetaminophen, Dextromethorphan HBr, Guaifenesin and Phenylephrine HCl Pain Reliever/Fever Reducer, Cough Suppressant, Expectorant, Nasal Decongestant

actual size [tablet image]

6 tablets (2 per pouch)

Drug Facts (continued)

Directions
 adults and children 12 years and over ■ take 2 tablets with water every 6-8 hours as needed
 ■ do not take more than 8 tablets in 24 hours, or as directed by a doctor
 children under 12 years of age ■ ask a doctor

Other information
 ■ store at room temperature 59-86° F (15-30° C)

Inactive Ingredients
 malto-dextrin, microcrystalline cellulose, povidone, silicon dioxide*, sodium starch glycolate, starch, stearic acid *may contain

Questions or comments?
 call toll-free 1-877-507-6516 (M-F 8AM-4:30PM CST)

Product manufactured for: L1 Drug Store Products, Inc., 8000 Earhart Lane SW Cedar Rapids, IA 52404 www.l1drugstore.com 56200-US-10-18-3 Proudly distributed by Circle K Stores Inc.

Warnings
 Liver warning: This product contains acetaminophen. Severe liver damage may occur if you take:
 ■ more than 4,000 mg of acetaminophen in 24 hours
 ■ with other drugs containing acetaminophen
 ■ 3 or more alcoholic drinks every day while using this product
 Allergy alert: Acetaminophen may cause severe skin reactions. Symptoms may include:
 ■ skin reddening ■ blisters ■ rash
 If a skin reaction occurs, stop use and seek medical help right away.
 Sore throat warning: If sore throat is severe, persists for more than 2 days, is accompanied or followed by fever, headache, rash, nausea or vomiting, consult a doctor promptly.
 A persistent cough may be a sign of a serious condition. If cough persists for more than 1 week, tends to recur, or is accompanied by fever, rash, or persistent headache, consult a doctor.

Drug Facts (continued)

Do not use
 ■ with any other drug containing acetaminophen (prescription or nonprescription). If you are not sure whether a drug contains acetaminophen, ask a doctor or pharmacist.
 ■ if you are now taking a prescription monoamine oxidase inhibitor (MAOI) (certain drugs for depression, psychiatric or emotional conditions, or Parkinson's disease), or for 2 weeks after stopping the MAOI drug. If you do not know if your prescription drug contains an MAOI, ask a doctor or pharmacist before taking this product.

Ask a doctor before use if you have
 ■ liver disease ■ heart disease
 ■ diabetes ■ thyroid disease ■ high blood pressure
 ■ trouble urinating due to an enlarged prostate gland
 ■ cough that occurs with too much phlegm (mucus)
 ■ persistent or chronic cough such as occurs with smoking, asthma, chronic bronchitis or emphysema

Ask a doctor or pharmacist before use if you are
 ■ taking the blood thinning drug warfarin

When using this product
 ■ do not exceed recommended dosage

Stop use and ask a doctor if
 ■ pain, nasal congestion or cough gets worse or lasts more than 7 days
 ■ fever gets worse or lasts more than 3 days
 ■ redness or swelling is present
 ■ new symptoms occur
 ■ cough comes back or is accompanied by a fever, rash or persistent headache
 ■ nervousness, dizziness, or sleeplessness occur. These could be signs of a serious condition.

CIRCLE K COLD

acetaminophen, dextromethorphan hydrobromide, guaifenesin, and phenylephrine hydrochloride tablet

Product Information			
Product Type	HUMAN OTC DRUG	Item Code (Source)	NDC:66715-5728
Route of Administration	ORAL		

Active Ingredient/Active Moiety			
Ingredient Name	Basis of Strength	Strength	
ACETAMINOPHEN (UNII: 362O9ITL9D) (ACETAMINOPHEN - UNII:362O9ITL9D)	ACETAMINOPHEN	325 mg	
DEXTROMETHORPHAN HYDROBROMIDE (UNII: 9D2RTI9KYH) (DEXTROMETHORPHAN - UNII:7355X3ROTS)	DEXTROMETHORPHAN HYDROBROMIDE	15 mg	
GUAIFENESIN (UNII: 495W7451VQ) (GUAIFENESIN - UNII:495W7451VQ)	GUAIFENESIN	200 mg	
PHENYLEPHRINE HYDROCHLORIDE (UNII: 04JA59TNSJ) (PHENYLEPHRINE - UNII:1WS297W6MV)	PHENYLEPHRINE HYDROCHLORIDE	5 mg	

Inactive Ingredients

Ingredient Name	Strength
MALTODEXTRIN (UNII: 7CVR7L4A2D)	
MICROCRYSTALLINE CELLULOSE (UNII: OP1R32D61U)	
POVIDONE, UNSPECIFIED (UNII: FZ989GH94E)	
SILICON DIOXIDE (UNII: ETJ7Z6XBU4)	
SODIUM STARCH GLYCOLATE TYPE A CORN (UNII: AG9B65PV6B)	
STARCH, CORN (UNII: O8232NY3SJ)	
STEARIC ACID (UNII: 4ELV7Z65AP)	

Product Characteristics

Color	white	Score	no score
Shape	ROUND	Size	12mm
Flavor		Imprint Code	FR12
Contains			

Packaging

#	Item Code	Package Description	Marketing Start Date	Marketing End Date
1	NDC:66715-5728-2	2 in 1 CARTON	12/31/2018	11/15/2022
1		2 in 1 POUCH; Type 0: Not a Combination Product		
2	NDC:66715-5728-3	3 in 1 CARTON	10/15/2019	09/09/2024
2		2 in 1 POUCH; Type 0: Not a Combination Product		

Marketing Information

Marketing Category	Application Number or Monograph Citation	Marketing Start Date	Marketing End Date
OTC monograph final	part341	11/01/2014	09/09/2024

CIRCLE K COLD

acetaminophen, dextromethorphan hydrobromide, guaifenesin, and phenylephrine hydrochloride tablet

Product Information

Product Type	HUMAN OTC DRUG	Item Code (Source)	NDC:66715-5528
Route of Administration	ORAL		

Active Ingredient/Active Moiety

Ingredient Name	Basis of Strength	Strength
ACETAMINOPHEN (UNII: 362O9ITL9D) (ACETAMINOPHEN - UNII:362O9ITL9D)	ACETAMINOPHEN	325 mg
DEXTROMETHORPHAN HYDROBROMIDE (UNII: 9D2RTI9KYH) (DEXTROMETHORPHAN - UNII:7355X3ROTS)	DEXTROMETHORPHAN HYDROBROMIDE	15 mg
GUAIFENESIN (UNII: 495W7451VQ) (GUAIFENESIN - UNII:495W7451VQ)	GUAIFENESIN	200 mg
PHENYLEPHRINE HYDROCHLORIDE (UNII: 04JA59TNSJ) (PHENYLEPHRINE - UNII:1WS297W6MV)	PHENYLEPHRINE HYDROCHLORIDE	5 mg

Inactive Ingredients

Ingredient Name	Strength
MALTODEXTRIN (UNII: 7CVR7L4A2D)	
MICROCRYSTALLINE CELLULOSE (UNII: OP1R32D61U)	
POVIDONE, UNSPECIFIED (UNII: FZ989GH94E)	
SILICON DIOXIDE (UNII: ETJ7Z6XBU4)	
SODIUM STARCH GLYCOLATE TYPE A CORN (UNII: AG9B65PV6B)	
STARCH, CORN (UNII: O8232NY3SJ)	
STEARIC ACID (UNII: 4ELV7Z65AP)	

Product Characteristics

Color	white	Score	no score
Shape	ROUND	Size	12mm
Flavor		Imprint Code	FR;12
Contains			

Packaging

#	Item Code	Package Description	Marketing Start Date	Marketing End Date
1	NDC:66715-5528-3	3 in 1 CARTON	03/26/2021	03/03/2026
1		2 in 1 POUCH; Type 0: Not a Combination Product		

Marketing Information

Marketing Category	Application Number or Monograph Citation	Marketing Start Date	Marketing End Date
OTC monograph final	part341	03/26/2021	03/03/2026

Labeler - Lil' Drug Store Products, Inc (093103646)

Revised: 3/2023

Lil' Drug Store Products, Inc