

UDDERCARE PLUS PART B- stabilized sodium salt of chlorous acid solution

Aspen Veterinary Resources

Disclaimer: This drug has not been found by FDA to be safe and effective, and this labeling has not been approved by FDA. For further information about unapproved drugs, [click here](#).

FOR INDUSTRIAL USE ONLY

KEEP OUT OF REACH OF CHILDREN

NOT FOR HUMAN USE

DANGER

Causes severe skin burns and eye damage. Wash skin thoroughly after handling. Wear protective gloves/protective clothing/ eye protection/ face protection.

DIRECTIONS:

IMPORTANT: Do not mix Uddercare+ Part B with water, or any other products unless specified in the product use instructions. If transferred from this container to any other, make sure that the other container is thoroughly pre-cleaned and bears the proper container labeling for Uddercare+ Part B.

If product in dip cup becomes visibly dirty, discard contents and replenish with undiluted product. Do not reuse or return any unused product to the original container.

Mix Directions: Measure equal volumes of Uddercare+ Part A and Uddercare+ Part B into clean containers. Mix until uniform in color. Mix only the quantity needed for one milking shift and discard the remainder.

Pre-Milk Dipping: Fill teat dip cup with properly mixed Uddercare+ Part A and Uddercare+ Part B. Before each cow is milked, dip each teat full-length into the teat dip cup containing the product. Wipe teats dry after dipping using single-service towels to avoid contamination of milk.

Post-Milk Dipping: Immediately after each milking, use mixed Uddercare+ Part A and Uddercare+ Part B at full strength. Dip each teat full-length into the teat dip cup containing mixed product. Allow to air dry. Do not wipe. Always use fresh full strength, mixed product. Do not turn cows out in freezing weather until product is completely dry.

PRECAUTION: Uddercare+ Part B is not intended to cure or help the healing of chapped or irritated teats. As with any germicide, irritation or sensitization may occur in sensitive animals. In case of teat irritation or chapping, have the condition examined and, if necessary, treated by a veterinarian.

FIRST AID

IN CASE OF EYE CONTACT: Rinse immediately with plenty of water, also under the eyelids, for at least 15 minutes. Remove contact lenses, if present and easy to do. Continue rinsing. Get medical attention immediately.

IN CASE OF SKIN CONTACT: Wash off immediately with plenty of water for at least 15 minutes. Use a mild soap if available. Wash clothing before reuse. Thoroughly clean shoes before reuse. Get medical attention immediately.

IF SWALLOWED: Rinse mouth with water. Do NOT induce vomiting. Never give anything by mouth to an unconscious person. Get medical attention immediately.

IF INHALED: Remove to fresh air. Treat symptomatically. Get medical attention if symptoms occur.

READ SAFETY DATA SHEET (SDS) BEFORE USING THIS PRODUCT

EMERGENCY HEALTH INFORMATION: Contact local poison control.

aspen DIARY SOLUTIONS

UDDERCARE+ PART B

ACTIVE INGREDIENT:

Stabilized Sodium Salt of Chlorous Acid.....0.54%

INERT INGREDIENTS:..... 99.46%

TOTAL:.....100.00%

Sanitizing Pre/Post Teat Dip

Uddercare+ Part B is formulated to be used as a companion product for Uddercare+ Part A.

An aid in reducing the spread of organisms which may cause mastitis

NET CONTENTS: 208 L (55 US GAL)

6301575

Manufactured for:

Aspen Veterinary Resources®, Ltd.

3155 Heartland Drive, Liberty, MO 64068

1-800-203-5620 761372/5300/0315



UDDERCARE+ PART B™



ACTIVE INGREDIENT:
Stabilized Sodium Salt of Chlorous Acid0.54%
INERT INGREDIENTS:..... 99.46%
TOTAL:.....100.00%

Sanitizing Pre/Post Teat Dip

Uddercare+ Part B is formulated to be used as a companion product for Uddercare+ Part A.
• An aid in reducing the spread of organisms which may cause mastitis

FOR INDUSTRIAL USE ONLY

KEEP OUT OF REACH OF CHILDREN

NOT FOR HUMAN USE

DIRECTIONS:

IMPORTANT: Do not mix Uddercare+ Part B with water, or any other products unless specified in the product use instructions. If transferred from this container to any other, make sure that the other container is thoroughly pre-cleaned and bears the proper container labeling for Uddercare+ Part B.

If product in dip cup becomes visibly dirty, discard contents and replenish with undiluted product. Do not reuse or return any unused product to the original container.

Mix Directions: Measure equal volumes of Uddercare+ Part A and Uddercare+ Part B into clean containers. Mix until uniform in color. Mix only the quantity needed for one milking shift and discard the remainder.

Pre-Milk Dipping: Fill teat dip cup with properly mixed Uddercare+ Part A and Uddercare+ Part B. Before each cow is milked, dip each teat full-length into the teat dip cup containing the product. Wipe teats dry after dipping using single-service towels to avoid contamination of milk.

Post-Milk Dipping: Immediately after each milking, use mixed Uddercare+ Part A and Uddercare+ Part B at full strength. Dip each teat full-length into the teat dip cup containing mixed product. Allow to air dry. Do not wipe. Always use fresh full strength, mixed product. Do not turn cows out in freezing weather until product is completely dry.

PRECAUTION: Uddercare+ Part B is not intended to cure or help the healing of chapped or irritated teats. As with any germicide, irritation or sensitization may occur in sensitive animals. In case of teat irritation or chapping, have the condition examined and, if necessary, treated by a veterinarian.

WARNING

Causes severe skin burns and eye damage. Wash skin thoroughly after handling. Wear protective gloves/ protective clothing/ eye protection/ face protection.

FIRST AID

IN CASE OF EYE CONTACT: Rinse immediately with plenty of water, also under the eyelids, for at least 15 minutes. Remove contact lenses, if present and easy to do. Continue rinsing. Get medical attention immediately.

IN CASE OF SKIN CONTACT: Wash off immediately with plenty of water for at least 15 minutes. Use a mild soap if available. Wash clothing before reuse. Thoroughly clean shoes before reuse. Get medical attention immediately.

IF SWALLOWED: Rinse mouth with water. Do NOT induce vomiting. Never give anything by mouth to an unconscious person. Get medical attention immediately.

IF INHALED: Remove to fresh air. Treat symptomatically. Get medical attention if symptoms occur.

READ SAFETY DATA SHEET (SDS) BEFORE USING THIS PRODUCT

EMERGENCY HEALTH INFORMATION: Contact local poison control.

Desinfectante antes y después sellador de pezones

Uddercare+ Part B está formulado para su uso como producto complementario de Soft Chlor Part A.
• Ayuda a evitar la proliferación de organismos que pueden causar mastitis

PARA USO INDUSTRIAL SOLAMENTE

MANTÉNGASE FUERA DEL ALCANCE DE LOS NIÑOS

NO ES PARA USO HUMANO

INSTRUCCIONES:

IMPORTANTE: No mezcle Uddercare+ Part B con agua ni con otros productos, a menos que se especifique en las instrucciones de uso. En caso de trasvase desde este recipiente a cualquier otro, este último debe limpiarse previamente a fondo y tener la correspondiente etiqueta de Uddercare+ Part B.

Si el producto en el aplicador presenta una suciedad notable, deseche el contenido y vuelva a llenarlo con producto sin diluir. No reutilice el producto ni vuelva a colocar en el recipiente original el producto que quede sin usar.

Instrucciones de mezclado: Mida volúmenes iguales de Uddercare+ Part A y Uddercare+ Part B en recipientes limpios. Mezcle hasta obtener un color uniforme. Mezcle sólo la cantidad necesaria para un turno de ordeño y deseche el resto.

Sellado previo al ordeño: Llene el aplicador con una mezcla correcta de Uddercare+ Part A y Uddercare+ Part B. Antes de cada ordeño, sumerja cada pezón por completo en el aplicador con producto. Posteriormente, seque los pezones con toallas descartables para evitar la contaminación de la leche.

Sellado posterior al ordeño: Inmediatamente después de cada ordeño, utilice una mezcla sin diluir de Uddercare+ Part A y Uddercare+ Part B. Sumerja cada pezón por completo en el aplicador con producto mezclado. Deje secar al aire. No limpie. Siempre utilice una nueva mezcla de producto sin diluir. No coloque las vacas en lugares con temperaturas bajo cero hasta que el producto esté completamente seco.

PRECAUCIÓN: Uddercare+ Part B no tiene por objeto curar ni ayudar a eliminar el agrietamiento y la irritación de los pezones. Al igual que con cualquier bactericida, puede producirse irritación o sensibilización en animales sensibles. En caso de irritación o agrietamiento de los pezones, consulte con un veterinario para su examen y, de ser necesario, su tratamiento.

PELIGRO

Provoca graves quemaduras en la piel y lesiones oculares. Lave la piel minuciosamente después del manejo. Usar guantes/ ropa de protección/ equipo de protección para los ojos/ la cara.

PRIMEROS AUXILIOS

CONTACTO CON LOS OJOS: Enjuague inmediatamente con abundante agua, también debajo de los párpados, por lo menos durante 15 minutos. Quitar las lentes de contacto, cuando estén presentes y pueda hacerse con facilidad. Proseguir con el lavado. Consultar inmediatamente un médico.

CONTACTO CON LA PIEL: Lave inmediatamente con mucha agua por lo menos durante 15 minutos. Utilice un jabón suave si es posible. Lavar la ropa antes de reutilizarla. Limpiar a fondo los zapatos antes de reutilizarlos. Consultar inmediatamente un médico.

INGESTIÓN: Enjuague la boca con agua. No provoque vómitos. Nunca debe administrarse nada por la boca a una persona inconsciente. Consultar inmediatamente un médico.

INHALACIÓN: Desplazar al aire libre. Trate sintomáticamente. Consultar un médico si los síntomas aparecen.

LEA LA FICHA DE DATOS DE SEGURIDAD ANTES DE USAR EL PRODUCTO

INFORMACIÓN DE EMERGENCIA SOBRE SALUD: Contacto local de veneno.

NET CONTENTS: 208 L (55 US GAL)

6301575



Manufactured for:
Aspen Veterinary Resources®, Ltd.
3155 Heartland Drive, Liberty, MO 64068
1-800-203-5620
761372/5300/0315

UDDERCARE PLUS PART B

stabilized sodium salt of chlorous acid solution

Product Information

Product Type	OTC ANIMAL DRUG	Item Code (Source)	NDC:46066-414
Route of Administration	TOPICAL		

Active Ingredient/Active Moiety

Ingredient Name	Basis of Strength	Strength
SODIUM CHLORITE (UNII: G538EBV4VF) (CHLORITE ION - UNII:Z63H374SB6)	SODIUM CHLORITE	5.4 mg in 1 mL

Inactive Ingredients

Ingredient Name	Strength
WATER (UNII: 059QF0KO0R)	

Packaging

#	Item Code	Package Description	Marketing Start Date	Marketing End Date
1	NDC:46066-414-17	56800 mL in 1 DRUM		
2	NDC:46066-414-21	208000 mL in 1 DRUM		

Marketing Information

Marketing Category	Application Number or Monograph Citation	Marketing Start Date	Marketing End Date
unapproved drug other		02/12/2015	

Labeler - Aspen Veterinary Resources (627265361)

Revised: 7/2019

Aspen Veterinary Resources