

PEPTO-BISMOL- bismuth subsalicylate suspension
The Procter & Gamble Manufacturing Company

Pepto Bismol®

Drug Facts

Active ingredient (in each 30 mL dose)

Bismuth subsalicylate 525 mg

Purpose

Upset stomach reliever and antidiarrheal

Uses

relieves

- traveler's diarrhea
- diarrhea
- upset stomach due to overindulgence in food and drink, including:
- heartburn
- indigestion
- nausea
- gas
- belching
- fullness

Warnings

Reye's syndrome:

Children and teenagers who have or are recovering from chicken pox or flu-like symptoms should not use this product. When using this product, if changes in behavior with nausea and vomiting occur, consult a doctor because these symptoms could be an early sign of Reye's syndrome, a rare but serious illness.

Allergy alert: Contains salicylate. Do not take if you are

- allergic to salicylates (including aspirin)
- taking other salicylate products

Do not use if you have

- an ulcer
- a bleeding problem
- bloody or black stool

Ask a doctor before use if you have

- fever
- mucus in the stool

Ask a doctor or pharmacist before use if you are taking any drug for

- anticoagulation (thinning the blood)
- diabetes
- gout
- arthritis

When using this product a temporary, but harmless, darkening of the stool and/or tongue may occur

Stop use and ask a doctor if

- symptoms get worse or last more than 2 days
- ringing in the ears or loss of hearing occurs
- diarrhea lasts more than 2 days

If pregnant or breast-feeding, ask a health professional before use.

Keep out of reach of children. In case of overdose, get medical help or contact a Poison Control Center right away.

Directions

- shake well before use
- only use dose cup provided
- adults and children 12 years and over:
- 30 mL (1 dose) every ½ hour or 60 mL (2 doses) every hour as needed for diarrhea/traveler's diarrhea
- 30mL (1 dose) every ½ hour as needed for overindulgence (upset stomach, heartburn, indigestion, nausea)
- do not exceed 8 doses (240 mL) in 24 hours
- use until diarrhea stops but not more than 2 days
- children under 12 years: ask a doctor
- drink plenty of clear fluids to help prevent dehydration caused by diarrhea

Other information

- **each 30 mL dose contains:** magnesium 3 mg, sodium 4 mg
 - salicylate 261 mg
 - low sodium
 - protect from freezing
 - avoid excessive heat (over 104°F or 40°C)
- **TAMPER EVIDENT:** Do not use if printed shrinkband is missing or broken.

Inactive ingredients

benzoic acid, D&C Red No. 22, D&C Red No. 28, flavor, gellan gum, magnesium aluminum silicate, methylcellulose, salicylic acid, sodium salicylate, sorbic acid, sucralose, water

Questions?

1-800-717-3786

DIST. BY PROCTER & GAMBLE,
CINCINNATI OH 45202

PRINCIPAL DISPLAY PANEL - 354 mL Bottle Label

Pepto

Bismol®

Bismuth Subsalicylate
Upset Stomach Reliever/
Antidiarrheal

5 SYMPTOM

RELIEF

NAUSEA
HEARTBURN
INDIGESTION
UPSET STOMACH
DIARRHEA

12 FL OZ (354 mL)

Drug Facts (continued)

If pregnant or breast feeding, ask a health professional before use.
Keep out of reach of children. In case of overdose, get medical help or contact a Poison Control Center right away.

Directions

- stable well before use
- only use dose cup provided
- adults and children 12 years and over:
• 30 mL (1 dose) every 1/2 hour or 60 mL (2 doses) every hour as needed for diarrhea/traveler's diarrhea
- 30 mL (1 dose) every 1/2 hour as needed for overindulgence (upset stomach, heartburn, indigestion, nausea)
- do not exceed 8 doses (240 mL) in 24 hours
- use until diarrhea stops but not more than 2 days
- children under 12 years: ask a doctor
- drink plenty of clear fluids to help prevent dehydration caused by diarrhea

Other information

- each 30 mL dose contains: magnesium 3 mg, sodium 4 mg
- salicylate 261 mg • low sodium • protect from freezing
- avoid excessive heat (over 104°F or 40°C)

Inactive ingredients benzoic acid, D&C Red No. 22, D&C Red No. 28, flavor, gelatin gum, magnesium aluminum silicate, methylcellulose, salicylic acid, sodium salicylate, sorbic acid, sucrose, water

Questions? 1-800-717-3786

DIST. BY PROCTER & GAMBLE,
CINCINNATI OH 45202

P&G
www.pg.com
Patents: www.pg.com/patents
www.pepto-bismol.com

01578504

**Pepto
Bismol®**

Bismuth Subsalicylate
Upset Stomach Reliever/
Antidiarrheal

**5 SYMPTOM
RELIEF**

NAUSEA
HEARTBURN
INDIGESTION
UPSET STOMACH
DIARRHEA

12 FL OZ (354 mL)

Drug Facts

Active ingredient (in each 30 mL dose)	Purpose
Bismuth subsalicylate 525 mg	Upset stomach, reliever and antidiarrheal

Uses relieves • travelers' diarrhea • diarrhea
• upset stomach due to overindulgence in food and drink, including:
• heartburn • indigestion • nausea • gas
• belching • fullness

Warnings
Reye's syndrome: Children and teenagers who have or are recovering from chicken pox or flu-like symptoms should not use this product. When using this product, if changes in behavior with nausea and vomiting occur, consult a doctor because these symptoms could be an early sign of Reye's syndrome, a rare but serious illness.
Allergy alert: Contains salicylate. Do not take if you are:
• allergic to salicylates (including aspirin)
• taking other salicylate products
Do not use if you have:
• an ulcer • a bleeding problem • bloody or black stool
Ask a doctor before use if you have:
• fever • mucus in the stool
Ask a doctor or pharmacist before use if you are taking any drug for:
• anticoagulation (thinning the blood) • diabetes • gout • arthritis
When using this product a temporary, but harmless, darkening of the stool and/or tongue may occur
Stop use and ask a doctor if:
• symptoms get worse or last more than 2 days
• ringing in the ears or loss of hearing occurs
• diarrhea lasts more than 2 days

PEPTO-BISMOL

bismuth subsalicylate suspension

Product Information				
Product Type	HUMAN OTC DRUG	Item Code (Source)	NDC:37000-438	
Route of Administration	ORAL			
Active Ingredient/Active Moiety				
Ingredient Name		Basis of Strength	Strength	
BISMUTH SUBSALICYLATE (UNII: 62TEY51RR1) (SALICYLIC ACID - UNII:O414PZ4LPZ, BISMUTH CATION - UNII:ZS9CD1I8YE)		BISMUTH SUBSALICYLATE	525 mg in 30 mL	
Inactive Ingredients				
Ingredient Name		Strength		
BENZOIC ACID (UNII: 8SKN0B0MIM)				
MAGNESIUM ALUMINUM SILICATE (UNII: 6M3P64V0NC)				
D&C RED NO. 22 (UNII: 1678RKX8RT)				
D&C RED NO. 28 (UNII: 767IP0Y5NH)				
SUCRALOSE (UNII: 96K6UQ3ZD4)				
SALICYLIC ACID (UNII: O414PZ4LPZ)				
SODIUM SALICYLATE (UNII: MQ1H85SYP)				
SORBIC ACID (UNII: X045WJ989B)				
WATER (UNII: 059QF0KO0R)				
METHYLCELLULOSE, UNSPECIFIED (UNII: Z944H5SN0H)				
Product Characteristics				
Color	pink	Score		
Shape		Size		
Flavor	WINTERGREEN	Imprint Code		
Contains				
Packaging				
#	Item Code	Package Description	Marketing Start Date	Marketing End Date
1	NDC:37000-438-04	118 mL in 1 BOTTLE, PLASTIC; Type 0: Not a Combination Product	03/01/2018	
2	NDC:37000-438-08	236 mL in 1 BOTTLE, PLASTIC; Type 0: Not a Combination Product	03/01/2018	
3	NDC:37000-438-12	354 mL in 1 BOTTLE, PLASTIC; Type 0: Not a Combination Product	03/01/2018	
4	NDC:37000-438-16	473 mL in 1 BOTTLE, PLASTIC; Type 0: Not a Combination Product	03/01/2018	
5	NDC:37000-438-24	2 in 1 CELLO PACK	03/01/2018	
5		354 mL in 1 BOTTLE, PLASTIC; Type 0: Not a Combination Product		
6	NDC:37000-438-32	2 in 1 CELLO PACK	03/01/2018	

6		473 mL in 1 BOTTLE, PLASTIC; Type 0: Not a Combination Product		
7	NDC:37000-438-34	100 mL in 1 BOTTLE, PLASTIC; Type 0: Not a Combination Product	03/01/2018	
Marketing Information				
Marketing Category		Application Number or Monograph Citation	Marketing Start Date	Marketing End Date
OTC Monograph Drug		M008	03/01/2018	

Labeler - The Procter & Gamble Manufacturing Company (004238200)

Revised: 10/2023

The Procter & Gamble Manufacturing Company