

MULTI-SYMPTOM COLD SEVERE, NON-DROWSY, DAYTIME- acetaminophen, dextromethorphan hbr, guaifenesin, phenylephrine hcl tablet, film coated
Valu Merchandisers Company

Best Choice 44-503A-Delisted

Active ingredients (in each caplet)

Acetaminophen 325 mg
Dextromethorphan HBr 10 mg
Guaifenesin 200 mg
Phenylephrine HCl 5 mg

Purpose

Pain reliever/fever reducer
Cough suppressant
Expectorant
Nasal decongestant

Uses

- temporarily relieves these common cold and flu symptoms:
 - sore throat
 - cough
 - nasal congestion
 - headache
 - minor aches and pains
- helps loosen phlegm (mucus) and thin bronchial secretions to make coughs more productive
- temporarily reduces fever

Warnings

Liver warning: This product contains acetaminophen. Severe liver damage may occur if you take

- more than 4,000 mg of acetaminophen in 24 hours
- with other drugs containing acetaminophen
- 3 or more alcoholic drinks every day while using this product

Allergy alert: Acetaminophen may cause severe skin reactions. Symptoms may include:

- skin reddening
- blisters
- rash

If a skin reaction occurs, stop use and seek medical help right away.

Sore throat warning: If sore throat is severe, persists for more than 2 days, is accompanied or followed by fever, headache, rash, nausea, or vomiting, consult a doctor promptly.

Do not use

- with any other drug containing acetaminophen (prescription or nonprescription). If you are not sure whether a drug contains acetaminophen, ask a doctor or pharmacist.
- if you are now taking a prescription monoamine oxidase inhibitor (MAOI) (certain drugs for depression, psychiatric or emotional conditions, or Parkinson's disease), or for 2 weeks after stopping the MAOI drug. If you do not know if your prescription drug contains an MAOI, ask a doctor or pharmacist before taking this product.
- if you have ever had an allergic reaction to this product or any of its ingredients

Ask a doctor before use if you have

- heart disease
- liver disease
- diabetes
- persistent or chronic cough such as occurs with smoking, asthma, chronic bronchitis, or emphysema
- thyroid disease
- cough that occurs with too much phlegm (mucus)
- difficulty in urination due to enlargement of the prostate gland
- high blood pressure

Ask a doctor or pharmacist before use if you are

taking the blood thinning drug warfarin.

When using this product

do not exceed recommended dosage.

Stop use and ask a doctor if

- nervousness, dizziness, or sleeplessness occur
- pain, nasal congestion, or cough gets worse or lasts more than 7 days
- new symptoms occur
- fever gets worse or lasts more than 3 days
- redness or swelling is present
- cough comes back or occurs with rash or headache that lasts. These could be signs of a serious condition.

If pregnant or breast-feeding,

ask a health professional before use.

Keep out of reach of children.

In case of accidental overdose, get medical help or contact a Poison Control Center (1-800-222-1222) right away. Prompt medical attention is critical for adults as well as for children even if you do not notice any signs or symptoms.

Directions

- **do not take more than directed**
- adults and children 12 years and over
 - take 2 caplets every 4 hours
 - swallow whole - do not crush, chew, or dissolve
 - do not take more than 10 caplets in 24 hours
- children under 12 years: ask a doctor

Other information

- **TAMPER EVIDENT: DO NOT USE IF OUTER PACKAGE IS OPENED OR BLISTER IS TORN OR BROKEN**
- store at 25°C (77°F); excursions permitted between 15°-30°C (59°-86°F)
- see end flap for expiration date and lot number

Inactive ingredients

corn starch, crospovidone, D&C yellow #10 aluminum lake, flavor, magnesium stearate, maltodextrin, microcrystalline cellulose, polyethylene glycol, polyvinyl alcohol, povidone, silicon dioxide, sodium starch glycolate, stearic acid, sucralose, talc, titanium dioxide

Questions or comments?

1-800-426-9391

Principal Display Panel

**Best
Choice®**

**Non-Drowsy
Severe Daytime
Multi-Symptom Cold**

ACETAMINOPHEN - Pain Reliever/Fever Reducer
Dextromethorphan HBr - Cough Suppressant
Guaifenesin - Expectorant
Phenylephrine HCl - Nasal Decongestant

Actual Size

Relieves Fever/Headache,
Sore Throat, Nasal Congestion,
Cough, Chest Congestion

Pseudoephedrine Free

**24 Cool Caplets
with Cool Blast Flavor**

**TAMPER EVIDENT: DO NOT USE IF PACKAGE IS OPENED OR IF BLISTER
UNIT IS TORN, BROKEN OR SHOWS ANY SIGNS OF TAMPERING**

50844 REV0718B50308

**PROUDLY DISTRIBUTED BY:
VALU MERCHANDISERS, CO.
5000 KANSAS AVE
KANSAS CITY, KS 66106**

**Best Choice®
100% Guaranteed**
www.bestchoicebrand.com

No print/No varnish
Lot & Exp date

PROUDLY DISTRIBUTED BY:
VALU MERCHANDISERS, CO.
5000 KANSAS AVE
KANSAS CITY, KS 66106
www.bestchoicebrand.com

Best Choice,
100% Guaranteed
Drug Facts (continued)
Questions or comments:

Non-Drowsy Severe Daytime Multi-Symptom Cold

Actual Size

ACETAMINOPHEN - Pain Reliever/Fever Reducer
Dextromethorphan HBr - Cough Suppressant
Guaifenesin - Expectorant
Phenylephrine HCl - Nasal Decongestant

Best
Choice®

Drug Facts

KEEP OUTER PACKAGE FOR
COMPLETE PRODUCT INFORMATION

Active ingredients (in each caplet) Purpose
Acetaminophen 325 mg.....Pain reliever/fever reducer
Dextromethorphan HBr 10 mg.....Cough suppressant
Guaifenesin 200 mg.....Expectorant
Phenylephrine HCl 5 mg.....Nasal decongestant

Drug Facts (continued)
Uses ■ temporarily relieves these common cold
and flu symptoms: ■ sore throat ■ cough
■ nasal congestion ■ headache
■ minor aches and pains
■ helps loosen phlegm (mucus) and thin bronchial
secretions to make coughs more productive

Drug Facts (continued)

■ temporarily reduces fever

Warnings

Liver warning: This product contains acetaminophen.
Severe liver damage may occur if you take
■ more than 4,000 mg of acetaminophen in 24 hours
■ with other drugs containing acetaminophen
■ 3 or more alcoholic drinks every day while using
this product
Allergy alert: Acetaminophen may cause severe skin
reactions. Symptoms may include:
■ skin redness ■ blisters ■ rash
If a skin reaction occurs, stop use and seek medical
help right away.
Sore throat warning: If sore throat is severe, persists
for more than 2 days, is accompanied or followed by
fever, headache, rash, nausea, or vomiting, consult a
doctor promptly.

Do not use

■ with any other drug containing acetaminophen
(prescription or nonprescription). If you are not
sure whether a drug contains acetaminophen, ask a
doctor or pharmacist.
■ if you are now taking a prescription monoamine
oxidase inhibitor (MAOI) (certain drugs for
depression, psychiatric or emotional conditions, or
Parkinson's disease), or for 2 weeks after stopping
the MAOI drug. If you do not know if your
prescription drug contains an MAOI, ask a doctor
or pharmacist before taking this product.
■ if you have ever had an allergic reaction to this
product or any of its ingredients

Ask a doctor before use if you have

■ heart disease ■ liver disease ■ diabetes

Drug Facts (continued)

■ persistent or chronic cough such as occurs with
smoking, asthma, chronic bronchitis, or
emphysema ■ thyroid disease
■ cough that occurs with too much phlegm (mucus)
■ difficulty in urination due to enlargement of the
prostate gland ■ high blood pressure
Ask a doctor or pharmacist before use if you are
taking the blood thinning drug warfarin.
When using this product do not exceed
recommended dosage.
Stop use and ask a doctor if
■ nervousness, dizziness, or sleeplessness occur
■ pain, nasal congestion, or cough gets worse or
lasts more than 7 days ■ new symptoms occur
■ fever gets worse or lasts more than 3 days
■ redness or swelling is present
■ cough comes back or occurs with rash or
headache that lasts. These could be signs of a
serious condition.

If pregnant or breast-feeding, ask a health

professional before use.
Keep out of reach of children. In case of accidental
overdose, get medical help or contact a Poison
Control Center (1-800-222-1222) right away. Prompt
medical attention is critical for adults as well as for
children even if you do not notice any signs or
symptoms.

Directions

■ do not take more than directed
■ adults and children 12 years and over
■ take 2 caplets every 4 hours
■ swallow whole - do not crush, chew, or dissolve
■ do not take more than 10 caplets in 24 hours
■ children under 12 years: ask a doctor

B-2115-503A-08CMSR
REV0718B50308

TAMPER EVIDENT: DO NOT USE IF PACKAGE IS OPENED OR IF BLISTER
UNIT IS TORN, BROKEN OR SHOWS ANY SIGNS OF TAMPERING

50844 REV0718B50308

DATE





Best Choice 44-503A

MULTI-SYMPTOM COLD SEVERE, NON-DROWSY, DAYTIME

acetaminophen, dextromethorphan hbr, guaifenesin, phenylephrine hcl tablet, film coated

Product Information

Product Type	HUMAN OTC DRUG	Item Code (Source)	NDC:63941-503
Route of Administration	ORAL		

Active Ingredient/Active Moiety

Ingredient Name	Basis of Strength	Strength
ACETAMINOPHEN (UNII: 362O9ITL9D) (ACETAMINOPHEN - UNII:362O9ITL9D)	ACETAMINOPHEN	325 mg
DEXTROMETHORPHAN HYDROBROMIDE (UNII: 9D2RTI9KYH) (DEXTROMETHORPHAN - UNII:7355X3ROTS)	DEXTROMETHORPHAN HYDROBROMIDE	10 mg
GUAIFENESIN (UNII: 495W7451VQ) (GUAIFENESIN - UNII:495W7451VQ)	GUAIFENESIN	200 mg
PHENYLEPHRINE HYDROCHLORIDE (UNII: 04JA59TNSJ) (PHENYLEPHRINE - UNII:1WS297W6MV)	PHENYLEPHRINE HYDROCHLORIDE	5 mg

Inactive Ingredients

Ingredient Name	Strength
STARCH, CORN (UNII: O8232NY3SJ)	
CROSPVIDONE (UNII: 2S7830E561)	
D&C YELLOW NO. 10 ALUMINUM LAKE (UNII: CQ3XH3DET6)	
MAGNESIUM STEARATE (UNII: 70097M6I30)	
MALTODEXTRIN (UNII: 7CVR7L4A2D)	
MICROCRYSTALLINE CELLULOSE (UNII: OP1R32D61U)	
POLYETHYLENE GLYCOL, UNSPECIFIED (UNII: 3WJQ0SDW1A)	
POLYVINYL ALCOHOL, UNSPECIFIED (UNII: 532B59J990)	
POVIDONE, UNSPECIFIED (UNII: FZ989GH94E)	
SILICON DIOXIDE (UNII: ETJ7Z6XBU4)	
SODIUM STARCH GLYCOLATE TYPE A POTATO (UNII: 5856J3G2A2)	
STEARIC ACID (UNII: 4ELV7Z65AP)	
SUCRALOSE (UNII: 96K6UQ3ZD4)	

TALC (UNII: 7SEV7J4R1U)

TITANIUM DIOXIDE (UNII: 15FIX9V2JP)

Product Characteristics

Color	yellow	Score	no score
Shape	OVAL	Size	19mm
Flavor	MENTHOL	Imprint Code	44;503
Contains			

Packaging

#	Item Code	Package Description	Marketing Start Date	Marketing End Date
1	NDC:63941-503-08	2 in 1 CARTON	08/04/2005	08/04/2025
1		12 in 1 BLISTER PACK; Type 0: Not a Combination Product		

Marketing Information

Marketing Category	Application Number or Monograph Citation	Marketing Start Date	Marketing End Date
OTC Monograph Drug	M012	08/04/2005	08/04/2025

Labeler - Valu Merchandisers Company (868703513)

Establishment

Name	Address	ID/FEI	Business Operations
LNK International, Inc.		832867837	manufacture(63941-503) , pack(63941-503)

Establishment

Name	Address	ID/FEI	Business Operations
LNK International, Inc.		832867894	manufacture(63941-503)

Establishment

Name	Address	ID/FEI	Business Operations
LNK International, Inc.		117025878	manufacture(63941-503)

Revised: 12/2023

Valu Merchandisers Company