ACETAMINOPHEN EXTRA STRENGTH- acetaminophen tablet, coated Target Corporation

Target 44-519

Active ingredient (in each gelcap)

Acetaminophen 500 mg

Purpose

Pain reliever/fever reducer

Uses

- temporarily relieves minor aches and pains due to:
 - headache
 - the common cold
 - backache
 - minor pain of arthritis
 - toothache
 - muscular aches
 - premenstrual and menstrual cramps
- temporarily reduces fever

Warnings

Liver warning: This product contains acetaminophen. Severe liver damage may occur if you take

- more than 4,000 mg of acetaminophen in 24 hours
- with other drugs containing acetaminophen
- 3 or more alcoholic drinks every day while using this product

Allergy alert: Acetaminophen may cause severe skin reactions. Symptoms may include:

- skin reddening
- blisters
- rash

If a skin reaction occurs, stop use and seek medical help right away.

Do not use

- with any other drug containing acetaminophen (prescription or nonprescription). If you are not sure whether a drug contains acetaminophen, ask a doctor or pharmacist.
- if you are allergic to acetaminophen or any of the inactive ingredients in this product

Ask a doctor before use if you have

liver disease.

Ask a doctor or pharmacist before use if you are

taking the blood thinning drug warfarin.

Stop use and ask a doctor if

- pain gets worse or lasts more than 10 days
- new symptoms occur
- fever gets worse or lasts more than 3 days
- redness or swelling is present

These could be signs of a serious condition.

If pregnant or breast-feeding,

ask a health professional before use.

Keep out of reach of children.

In case of overdose, get medical help or contact a Poison Control Center right away. Prompt medical attention is critical for adults as well as for children even if you do not notice any signs or symptoms.

Directions

- do not take more than directed
- adults and children 12 years and over
 - take 2 gelcaps every 6 hours while symptoms last
 - do not take more than 6 gelcaps in 24 hours, unless directed by a doctor
 - do not take for more than 10 days unless directed by a doctor
- children under 12 years: ask a doctor

Other information

- store at 25°C (77°F); excursions permitted between 15°-30°C (59°-86°F)
- avoid high humidity
- use by expiration date on package

Inactive ingredients

croscarmellose sodium, D&C red #33, FD&C blue #1, FD&C red #40, gelatin, hydroxypropyl cellulose, hypromellose, iron oxide black, iron oxide red, iron oxide yellow, polyethylene glycol, povidone, pregelatinized starch, propylene glycol, shellac glaze, stearic acid, titanium dioxide

Call 1-800-910-6874

Principal Display Panel

NDC 11673-519-15

Compare to active ingredient in Extra Strength Tylenol® Rapid Release Gels*

extra strength acetaminophen

gelcaps, 500 mg

pain reliever/fever reducer


up&up_™

50 GELCAPS

TAMPER EVIDENT: DO NOT USE IF IMPRINTED SAFETY SEAL UNDER CAP IS BROKEN OR MISSING

*This product is not manufactured or distributed by Johnson & Johnson Corporation, owner of the registered trademark Extra Strength Tylenol® Rapid Release Gels. 50844 REV0322C51915

094 01 0548 R05 ID225450 Distributed by Target Corporation Minneapolis, MN 55403 TM & ©2022 Target Brands, Inc.



Drug Facts (continued)

Narnings

acetaminophen. Severe liver damage may iver warning: This product contains occur if you take

more than 4,000 mg of acetaminophen in

24 hours

■with other drugs containing acetaminophen ■3 or more alcoholic drinks every day while

Allergy alert: Acetaminophen may cause using this product

severe skin reactions. Symptoms may skin reddening blisters include

Poison Control Center right away. Prompt medical attention is critical for adults as well

Keep out of reach of children. In case of overdose, get medical help or contact a

Drug Facts (continued)

STOP PEELING

as for children even if you do not notice any

signs or symptoms.

If a skin reaction occurs, stop use and seek ■ rash medical help right away

whether a drug contains acetaminophen nonprescription). If you are not sure acetaminophen (prescription or ■ with any other drug containing Do not use

ask a doctor or pharmacist

if you are allergic to acetaminophen or any

of the inactive ingredients in this product Ask a doctor before use if you have liver Ask a doctor or pharmacist before use i

warfarin.

store at 25°C (77°F); excursions permitted between 15°-30°C (59°-86°F)

/ou are taking the blood thinning drug

■ pain gets worse or lasts more than 10 Stop use and ask a doctor if

■fever gets worse or lasts more than 3 days new symptoms occur redness or swelling is present

nactive ingredients croscarmellose

use by expiration date on package

avoid high humidity

sodium, D&C red #33, FD&C blue #1, FD&C

red #40, gelatin, hydroxypropyl cellulose,

rėd, iron oxidė yellow, polyethylene glycol, povidone, pregelatinized starch, propylene wpromellose, iron oxide black, iron oxide

These could be signs of a serious condition.

if pregnant or breast-feeding, ask a health

professional before use.

Call 1-800-910-6874 **Questions?**

alycol, shellac glaze, stearic acid, titanium dioxide

Target 44-519

24 hours, unless directed by a doctor

do not take for more than 10 days

children under 12 years: ask a doctor

Other information

unless directed by a doctor

do not take more than 6 gelcaps in

symptoms last

adults and children 12 years and over take 2 gelcaps every 6 hours while

do not take more than directed

Directions

EXTRA STRENGTH ACETAMINOPHEN

acetaminophen tablet, coated

Product Information			
Product Type	HUMAN OTC DRUG	Item Code (Source)	NDC:11673-519
Route of Administration	ORAL		

Active Ingredient/Active Moiety				
Ingredient Name	Basis of Strength	Strength		
ACETAMINOPHEN (UNII: 36209ITL9D) (ACETAMINOPHEN - UNII: 36209ITL9D)	ACETAMINOPHEN	500 mg		

Inactive Ingredients	
Ingredient Name	Strength
CROSCARMELLOSE SODIUM (UNII: M280L1HH48)	
D&C RED NO. 33 (UNII: 9DBA0SBB0L)	
FD&C BLUE NO. 1 (UNII: H3R47K3TBD)	
FD&C RED NO. 40 (UNII: WZB9127XOA)	
GELATIN, UNSPECIFIED (UNII: 2G86QN327L)	
HYDROXYPROPYL CELLULOSE, UNSPECIFIED (UNII: 9XZ8H6N6OH)	
HYPROMELLOSE, UNSPECIFIED (UNII: 3NXW29V3WO)	
FERROSOFERRIC OXIDE (UNII: XM0M87F357)	
FERRIC OXIDE RED (UNII: 1K09F3G675)	
FERRIC OXIDE YELLOW (UNII: EX43802MRT)	
POLYETHYLENE GLYCOL, UNSPECIFIED (UNII: 3WJQ0SDW1A)	
POVIDONE, UNSPECIFIED (UNII: FZ989GH94E)	
STARCH, CORN (UNII: O8232NY3SJ)	
PROPYLENE GLYCOL (UNII: 6DC9Q167V3)	
SHELLAC (UNII: 46N107B710)	
STEARIC ACID (UNII: 4ELV7Z65AP)	
TITANIUM DIOXIDE (UNII: 15FIX9V2JP)	

Product Characteristics			
Color	red, blue	Score	no score
Shape	OVAL	Size	19mm
Flavor		Imprint Code	L;5
Contains			

P	Packaging				
#	Item Code	Package Description	Marketing Start Date	Marketing End Date	
1	NDC:11673- 519-15	50 in 1 BOTTLE, PLASTIC; Type 0: Not a Combination Product	05/10/2004		
2	NDC:11673- 519-12	100 in 1 BOTTLE, PLASTIC; Type 0: Not a Combination Product	05/10/2004		
3	NDC:11673- 519-20	225 in 1 BOTTLE, PLASTIC; Type 0: Not a Combination Product	05/10/2004	02/22/2023	

Marketing Information				
Marketing Category	Application Number or Monograph Citation	Marketing Start Date	Marketing End Date	
OTC Monograph Drug	M013	05/10/2004		

Labeler - Target Corporation (006961700)

EstablishmentNameAddressID/FEIBusiness OperationsLNK International, Inc.038154464manufacture(11673-519), pack(11673-519)

Establishment			
Name	Address	ID/FEI	Business Operations
LNK International, Inc.		832867837	manufacture(11673-519)

Establishment			
Name	Address	ID/FEI	Business Operations
LNK International, Inc.		832867894	manufacture(11673-519)

Establishment			
Name	Address	ID/FEI	Business Operations
LNK International, Inc.		868734088	manufacture(11673-519)

Establishment			
Name	Address	ID/FEI	Business Operations
LNK International, Inc.		967626305	pack(11673-519)

Revised: 2/2024 Target Corporation