ANTI DIARRHEAL- loperamide hcl tablet, film coated CVS Pharmacy

CVS Pharmacy, Inc. Anti-Diarrheal Drug Facts

Active ingredient (in each caplet)

Loperamide HCl 2 mg

Purpose

Anti-diarrheal

Use

controls symptoms of diarrhea, including Travelers' Diarrhea

Warnings

Allergy alert: Do not use if you have ever had a rash or other allergic reaction to loperamide HCl

Heart alert: Taking more than directed can cause serious heart problems or death

Do not use

if you have bloody or black stool

Ask a doctor before use if you have

- fever
- mucus in the stool
- a history of liver disease
- a history of abnormal heart rhythm

Ask a doctor or pharmacist before use if you are

taking a prescription drug. Loperamide may interact with certain prescription drugs.

When using this product

tiredness, drowsiness or dizziness may occur. Be careful when driving or operating machinery.

Stop use and ask a doctor if

- symptoms get worse
- diarrhea lasts for more than 2 days
- you get abdominal swelling or bulging. These may be signs of a serious condition.

If pregnant or breast-feeding,

ask a health professional before use.

Keep out of reach of children.

In case of overdose, get medical help or contact a Poison Control Center right away (1-800-222-1222).

Directions

- drink plenty of clear fluids to help prevent dehydration caused by diarrhea
- find right dose on chart. If possible, use weight to dose; otherwise, use age.

adults and children 12 years and over	2 caplets after the first loose stool; 1 caplet after each subsequent loose stool; but no more than 4 caplets in 24 hours
children 9-11 years (60-95 lbs)	1 caplet after the first loose stool; 1/2 caplet after each subsequent loose stool; but no more than 3 caplets in 24 hours
children 6-8 years (48-59 lbs)	1 caplet after the first loose stool; 1/2 caplet after each subsequent loose stool; but no more than 2 caplets in 24 hours
children 2-5 years (34 to 47 lbs)	ask a doctor
children under 2 years (up to 33 lbs)	do not use

Other information

- store at 20°-25°C (68°-77°F)
- see end panel for lot number and expiration date

Inactive ingredients

anhydrous lactose, carnauba wax, D&C yellow no. 10 aluminum lake, FD&C blue no. 1 aluminum lake, hypromellose, magnesium stearate, microcrystalline cellulose, polyethylene glycol, pregelatinized starch

Questions or comments?

1-800-719-9260

Principal Display Panel

Compare to the active ingredient in Imodium® A-D

Anti-Diarrheal

Loperamide Hydrochloride Tablets, 2 mg

Anti-Diarrheal

Controls the symptoms of diarrhea

THIS PACKAGE FOR HOUSEHOLDS WITHOUT YOUNG CHILDREN

Actual Size

SEE NEW WARNINGS

24 CAPLETS*

*Capsule-Shaped Tablets



OPEN OTHER END

96062 17 CB

Drug Facts

Active in gredient (in each ca plet) Lo pera mide HCl 2 mg..

....Anti-diarrheal

Use controls symptoms of darrhea, including Travelers' Diarrhe a.

Allergy alert: Do not use if you have ever had a rash or other allergis reaction to loperamide HCI He art all ert: Taking more than directed can cause serious heart problems on death

Do not use if you have bloody or black sholl

As k a doctor before use if you have

- ■fever mucus in the stool
- au history of liver disease
- a. history of ab normal heart rhythm

As k a dector or pharmacist before use if you are taking a prescription drug. Lopera mide may interact with certain pre-scription drugs.

When using this product tirechess, dro weiness or dizzin ess may occur. Be careful

Drug Facts (continued)

Keepout of reach of children. In case of overabse, get me dical help or contact a Poison Control Cente rright away (1-80 0-2 22 -12 22).

- drink plenty of clear fluids to help prevent de hydration cause d by dianr he a
- find right dose on chart. If possible, use

weightto obse; otherwise, use age.	
a dults and chiblen 12 years and over	2 cap lets after the first loose stool; 1 caplet after each subsequent loose stool; but no more than 4 cap lets in 2 4 hours
childen 9-11years (60-95 lbs)	1 cap let after the first loose stool; 1.2 caplet after each subsequent loose stool; but no more than 3 cap lets in 2.4 hours
chiblen 6-8 years (48-59 lbs)	1 cap let after the first loose stool; 1.2 caplet after each subsequent loose stool; but no more than 2 cap lets in 2.4 hours
chibben 2-5 years (34 to 47 lbs)	ask a doct or



ANTI DIARRHEAL

loperamide hcl tablet, film coated

Product Information

Product Type HUMAN OTC DRUG Item Code (Source) NDC:59779-960

Route of Administration ORAL

Active Ingredient/Active Moiety

Ingredient Name Basis of Strength Strength

LOPERAMIDE HYDROCHLORIDE (UNII: 77TI35393C) (LOPERAMIDE - LOPERAMIDE - HYDROCHLORIDE 2 mg

Inactive Ingredients

Ingredient Name	Strength
ANHYDROUS LACTOSE (UNII: 3SY5LH9PMK)	
CARNAUBA WAX (UNII: R12CBM0EIZ)	

D&C YELLOW NO. 10 (UNII: 35SW5USQ3G)

FD&C BLUE NO. 1 (UNII: H3R47K3TBD)

HYPROMELLOSE, UNSPECIFIED (UNII: 3NXW29V3WO)

MAGNESIUM STEARATE (UNII: 70097M6I30)

MICROCRYSTALLINE CELLULOSE (UNII: OP1R32D61U)

POLYETHYLENE GLYCOL, UNSPECIFIED (UNII: 3WJQ0SDW1A)

Product Characteristics

Color	GREEN	Score	2 pieces
Shape	OVAL	Size	9mm
Flavor		Imprint Code	L2
Contains			

Packaging				
#	Item Code	Package Description	Marketing Start Date	Marketing End Date
1	NDC:59779-960- 62	24 in 1 CARTON	04/21/2003	
1		1 in 1 BLISTER PACK; Type 0: Not a Combination Product		

Marketing Information			
Marketing Category	Application Number or Monograph Citation	Marketing Start Date	Marketing End Date
ANDA	ANDA075232	04/21/2003	

Labeler - CVS Pharmacy (062312574)

Revised: 10/2022 CVS Pharmacy