# BENADRYL ALLERGY PLUS CONGESTION- diphenhydramine hydrochloride and phenylephrine hydrochloride tablet, film coated Johnson & Johnson Consumer Inc.

-----

### **Benadryl Allergy Plus Congestion**

# **Drug Facts**

Active ingredients (in each tablet)	Purposes
Diphenhydramine HCl 25 mg	Antihistamine
Phenylephrine HCl 10 mg	Nasal decongestant

#### Uses

- temporarily relieves these symptoms due to hay fever or other upper respiratory allergies:
  - runny nose
  - sneezing
  - itchy, watery eyes
  - itching of the nose or throat
  - nasal congestion
- temporarily relieves these symptoms due to the common cold:
  - runny nose
  - sneezing
  - nasal congestion
- temporarily relieves sinus congestion and pressure

# **Warnings**

#### Do not use

- to make a child sleepy
- with any other product containing diphenhydramine, even one used on skin
- if you are now taking a prescription monoamine oxidase inhibitor (MAOI) (certain drugs for depression, psychiatric or emotional conditions, or Parkinson's disease), or for 2 weeks after stopping the MAOI drug. If you do not know if your prescription drug contains an MAOI, ask a doctor or pharmacist before taking this product.

# Ask a doctor before use if you have

- heart disease
- high blood pressure
- thyroid disease
- diabetes
- trouble urinating due to an enlarged prostate gland
- a breathing problem such as emphysema or chronic bronchitis
- glaucoma

### Ask a doctor or pharmacist before use if you are taking sedatives or tranquilizers

#### When using this product

- do not exceed recommended dose
- marked drowsiness may occur
- avoid alcoholic drinks
- alcohol, sedatives, and tranquilizers may increase drowsiness
- be careful when driving a motor vehicle or operating machinery
- excitability may occur, especially in children

### Stop use and ask a doctor if

- nervousness, dizziness, or sleeplessness occur
- symptoms do not improve within 7 days or occur with a fever

**If pregnant or breast-feeding,** ask a health professional before use.

### Keep out of reach of children.

In case of overdose, get medical help or contact a Poison Control Center right away. (1-800-222-1222)

#### **Directions**

adults and children 12 years and over	<ul> <li>take 1 tablet every 4 hours</li> <li>do not take more than 6 tablets in 24 hours</li> </ul>
children under 12 years	ask a doctor

#### Other information

- store between 20-25°C (68-77°F).
- do not use if carton is opened or if blister unit is broken

# Inactive ingredients

carnauba wax, FD&C blue no. 1 aluminum lake, magnesium stearate, microcrystalline cellulose, modified starch, polyethylene glycol, polyvinyl alcohol, powdered cellulose, pregelatinized starch, sodium starch glycolate, talc, titanium dioxide

#### Questions or comments?

call **1-888-217-2117** (toll-free) or **215-273-8755** (collect)

#### PRINCIPAL DISPLAY PANEL

NDC 50580-415-01

# Benadryl<sup>®</sup>

# ALLERGY PLUS CONGESTION

# Diphenhydramine HCl 25mg | Antihistamine Phenylephrine HCl 10mg | Nasal Decongestant

- ✓ Sneezing
- ✓ Runny Nose
- ✓ Sinus Congestion
- ✓ Itchy, Watery Eyes
- ✓ Itchy Throat
- ✓ Sinus Pressure

## **ULTRATABS** ®\*

\*small tablet size

actual size

#### **24 TABLETS**



# **ALLERGY PLUS**

#### **EFFECTIVE ALLERGY RELIEF WHEN YOU NEED IT!®**

3 0045-0557-26

CSI[1-888-717-7117 (IDIFIIBB) OF 215-275-8755 (COLIBCT) Questions or comments?

starch glycolate, talc, titarium doxide Bitco, polyviny ialcohol, powdered cellulose, pregetalarzed starch, sodium magnesium stearate, microcrystalline cellulose, modified starch, polyethylene Inactive ingredients camauba wax, FD&C blue no. 1 aluminum lake,

> ■ do not use if carton is opened or if blister unit is broken ■ store between 20-25°C (68-77°F) Other information

lzyears children under	ask a doctor
JZ y eens and ower adults and children	■ do not take more fran 6 tablets in 24 hours

#### DIRECTIONS

(1-800-222-1222) In case of overdose, get medical help or contact a Polson Control Center right lf pregnant or breast leeding, ask aheath professional before use. Keep out of reach of children.

■ symptoms do not improve within 7 days or occur with a fever ■ Demouspess, dizziress, of sieepiess ress occul Stop use and ask a doctor if

Drug Facts (confinued)

exatability may occur, especially in children

- be careful when driving a motor vehicle or operating machinery sseusworb esearoni yem sreziliupnet bne , sevisbes , bricote ≡
  - - SANIA SICOTORIC ATINIS

■ marked drowsiness may occur When using this product **m** do not exceed recommended dose

Ask a doctor or pharmacist before use if you are taking sedatives or

- spreaguitó brogem andras emplykema or dronic pronquiga
- m trouble uninating due to an enlarged prostate gland eunsseud poolq q6µ = safedsib ... eseesib bior yith

eseesp µeeu ■ Ask a doctor before use if you have pharmadat before taking this product.

not know if your prescription drug contains an MAOI, ask a doctor or Parkinsori's disease), orfor 2 weeks after stopping the MAOI drug ffyou do (certain drugs for depression, psychiatric or emotional conditions, or

■ If you are now taking a prescription monoamine oxidase inhibitor (MAOI) ■ with any other product containing dehening remine, even one used on skin ■ to make a child sleepy

osn ton ou Warnings

- pubocsuj\u00e3 jenes ajuns cauβesgou sug buesarue
- uouseguco reseum gueseus escu /uuru m
- inconomino entro d'en bandem y sesent seveller (il moorauli).
  - mitching of the nose or throat messi congestion se/e Augum /Augum Guizeeus m esou Auuru m
    - respiratory altergles:
- temporarily relieves these symptoms due to hay fever or other upper SASU

mataegnooebleasM. Phanylephrine HCl 10 mg. enims aritinA Diphenhydramine HCl 25 mg. Active ingredients (in each tablet) sasod und

Drug Facts

Benadry

ALLERGY <del>PLUS</del>

EFFECTIVE ALLERGY RELIEF WHEN YOU NEED IT!®

NDC 50580-415-01

6



# ALLERGY <del>P</del>LUS CONGESTION

Diphenhydramine HCl 25mg | Antihistamine Phenylephrine HCl 10mg | Nasal Decongestant

- Sneezing
- Runny Nose Sinus Congestion
- 🗸 Itchy, Watery Eyes 💆 Itchy Throat 💆 Sinus Pressure


McNeil Consumer Healthcare Division Fort Washington, PA 19034 USA ©J&JCI 2016

ALLERGY

stributed by: J&ICI

98 67





# **BENADRYL ALLERGY PLUS CONGESTION**

diphenhydramine hydrochloride and phenylephrine hydrochloride tablet, film coated

Product Information			
Product Type	HUMAN OTC DRUG	Item Code (Source)	NDC:50580-415
Route of Administration	ORAL		

Active Ingredient/Active Moiety			
Ingredient Name	Basis of Strength	Strength	
<b>DIPHENHYDRAMINE HYDROCHLORIDE</b> (UNII: TC2D6JAD40) (DIPHENHYDRAMINE - UNII:8GTS82S83M)	DIPHENHYDRAMINE HYDROCHLORIDE	25 mg	
PHENYLEPHRINE HYDROCHLORIDE (UNII: 04JA59TNSJ) (PHENYLEPHRINE - UNII: 1W5297W6MV)	PHENYLEPHRINE HYDROCHLORIDE	10 mg	

Inactive Ingredients			
Ingredient Name	Strength		
CARNAUBA WAX (UNII: R12CBM0EIZ)			
FD&C BLUE NO. 1 ALUMINUM LAKE (UNII: J9EQA3S2JM)			
MAGNESIUM STEARATE (UNII: 70097M6I30)			
MICROCRYSTALLINE CELLULOSE (UNII: OP1R32D61U)			
POLYETHYLENE GLYCOL, UNSPECIFIED (UNII: 3WJQ0SDW1A)			
POLYVINYL ALCOHOL, UNSPECIFIED (UNII: 532B59J990)			
POWDERED CELLULOSE (UNII: SMD1X3XO9M)			
SODIUM STARCH GLYCOLATE TYPE A (UNII: H8AV0SQX4D)			
TALC (UNII: 7SEV7J4R1U)			
TITANIUM DIOXIDE (UNII: 15FIX9V2JP)			

Product Characteristics			
Color	blue	Score	no score
Shape	OVAL	Size	13mm
Flavor		Imprint Code	PE;WL95
Contains			

Packaging			
# Item Co	ode Package Description	Marketing Start Date	Marketing End Date
1 NDC:50580	2 in 1 CARTON	06/19/2017	
1	12 in 1 BLISTER PACK; Type 0: Not a Com Product	bination	

Marketing Information			
Marketing Category	Application Number or Monograph Citation	Marketing Start Date	Marketing End Date
OTC Monograph Drug	M012	06/19/2017	

# Labeler - Johnson & Johnson Consumer Inc. (878046358)

Revised: 1/2024 Johnson & Johnson Consumer Inc.