

IBUPROFEN - ibuprofen capsule, liquid filled
Strides Pharma Inc

DRUG FACTS

READ AND KEEP CARTON FOR COMPLETE WARNINGS AND INFORMATION

Active ingredient (in each capsule)

Solubilized ibuprofen equal to 200 mg ibuprofen (NSAID)*

(Present as the free acid and potassium salt)

*nonsteroidal anti-inflammatory drug

Purpose

Pain reliever/Fever reducer

Uses

- temporarily relieves minor aches and pains due to:
 - headache
 - toothache
 - backache
 - menstrual cramps
 - the common cold
 - muscular aches
 - minor pain of arthritis
- temporarily reduces fever

Warnings

Allergy alert: Ibuprofen may cause a severe allergic reaction, especially in people allergic to aspirin.

Symptoms may include:

- hives
- facial swelling
- asthma (wheezing)
- shock
- skin reddening
- rash
- blisters

If an allergic reaction occurs, stop use and seek medical help right away.

Stomach bleeding warning: This product contains an NSAID, which may cause severe stomach bleeding. The chance is higher if you

- are aged 60 or older
- have had stomach ulcers or bleeding problems
- take a blood thinning (anticoagulant) or steroid drug
- take other drugs containing prescription or nonprescription NSAIDs [aspirin, ibuprofen, naproxen, or others]
- have 3 or more alcoholic drinks every day while using this product
- take more or for a longer time than directed

Heart attack and stroke warning: NSAIDs, except aspirin, increase the risk of heart attack, heart failure, and stroke. These can be fatal. The risk is higher if you use more than directed or for longer than directed.

Do not use

- if you have ever had an allergic reaction to any other pain reliever/fever reducer
- right before or after heart surgery

Ask a doctor before use if

- stomach bleeding warning applies to you
- you have problems or serious side effects from taking pain relievers or fever reducers
- you have a history of stomach problems, such as heartburn
- you have high blood pressure, heart disease, liver cirrhosis, kidney disease, asthma, or had a stroke
- you are taking a diuretic

Ask a doctor or pharmacist before use if you are

- under a doctor's care for any serious condition
- taking aspirin for heart attack or stroke, because ibuprofen may decrease this benefit of aspirin
- taking any other drug

When using this product

- take with food or milk if stomach upset occurs

Stop use and ask a doctor if

- you experience any of the following signs of stomach bleeding:
 - feel faint
 - vomit blood
 - have bloody or black stools
 - have stomach pain that does not get better
- you have symptoms of heart problems or stroke:
 - chest pain
 - trouble breathing
 - weakness in one part or side of body
 - slurred speech
 - leg swelling
- pain gets worse or lasts more than 10 days
- fever gets worse or lasts more than 3 days
- redness or swelling is present in the painful area

- any new symptoms appear

If pregnant or breast-feeding, ask a health professional before use. It is especially important not to use ibuprofen at 20 weeks or later in pregnancy unless definitely directed to do so by a doctor because it may cause problems in the unborn child or complications during delivery.

Keep out of reach of children. In case of overdose, get medical help or contact a Poison Control Center right away.

Directions

- **do not take more than directed**
- **the smallest effective dose should be used**
- adults and children 12 years and over: take 1 capsule every 4 to 6 hours while symptoms persist
- if pain or fever does not respond to 1 capsule, 2 capsules may be used
- do not exceed 6 capsules in 24 hours, unless directed by a doctor
- children under 12 years: ask a doctor

Other information

- each capsule contains: **potassium 17.840 mg**
- read all warnings and directions before use. Keep carton.
- store at 20-25°C (68-77°F)
- avoid excessive heat above 40°C (104°F)

Inactive ingredients

Ibuprofen Capsules

FD&C blue 1, gelatin, gelatin (hydrolyzed), medium chain triglyceride, PEG 600, potassium hydroxide, purified water, sorbital and sorbitan solution.

Printing ink: propylene glycol, shellac resins, sodium lauryl sulphate, titanium dioxide.

Ibuprofen Capsules - Minis

FD&C green no.3, gelatin, gelatin (hydrolyzed), lecithin (Soya Lecithin), medium chain triglyceride, polyethylene Glycol 600, potassium hydroxide, purified water, sorbital and sorbitan solution.

Printing ink: propylene glycol, shellac resins, sodium lauryl sulphate, titanium dioxide.

Questions or comments?

call toll free **1-877-244-9825**

Do Not Use if seal under bottle cap imprinted with "SEALED for YOUR PROTECTION" is broken or missing.

For most recent product information, visit us at www.strides.com

Revised: 05/2023

Manufactured by:

Strides Pharma Science Limited

Bengaluru - 562106, India

Distributed by:

Strides Pharma Inc.

East Brunswick, NJ 08816

PRINCIPAL DISPLAY PANELS

Ibuprofen Capsules, 200 mg Pain Reliever/Fever Reducer (NSAID)

Liquid Filled Capsules

NDC 59556-764-48

READ AND KEEP CARTON FOR COMPLETE WARNINGS AND INFORMATION

Ibuprofen Liquid Gels

200 mg

Solubilized Ibuprofen Capsules, 200 mg Pain Reliever/Fever Reducer (NSAID)

160 Liquid Filled Capsules

Strides Pharma Inc.

Do Not Use if seal under bottle cap imprinted with "SEALED for YOUR PROTECTION" is broken or missing.

Uses

- temporarily relieves minor aches and pains due to: headache, muscular aches, menstrual cramps, the common cold, backache, toothache, minor pain of arthritis.
- temporarily reduces fever

Warnings

- Ask your doctor before use if you are pregnant, under a doctor's care for a serious condition, age 60 or over, taking any other drug or have stomach problems.
- This product may cause a severe allergic reaction, especially in people allergic to aspirin. Symptoms may include: hives, facial swelling, asthma (wheezing), shock, skin reddening, rash, blisters. If an allergic reaction occurs, stop use and seek medical help right away. Do not use this product if you have ever had an allergic reaction to any pain reliever/fever reducer.

Directions

- do not take more than directed
- the smallest effective dose should be used
- adults and children 12 years and over: take

Stomach bleeding warning: This product contains a nonsteroidal anti-inflammatory drug (NSAID), which may cause severe stomach bleeding. The chance is higher if you

- are age 60 or older
- have had stomach ulcers or bleeding problems
- take a blood thinning (anticoagulant) or steroid drug
- take other drugs containing prescription or nonprescription NSAIDs (aspirin, ibuprofen, naproxen, or others)
- have 3 or more alcoholic drinks every day while using this product
- take more or for a longer time than directed.

Heart attack and stroke warning: NSAIDs, except aspirin, increase the risk of heart attack, heart failure, and stroke. These can be fatal. The risk is higher if you use more than directed or for longer than directed.

1 capsule every 4 to 6 hours while symptoms persist

- if pain or fever does not respond to 1 capsule, 2 capsules may be used
- do not exceed 6 capsules in 24 hours, unless directed by a doctor
- children under 12 years: ask a doctor.

Each capsule contains: **potassium 17.840 mg**

Questions or comments?
call toll free 1-877-244-9825

For most recent product information, visit www.strides.com

Manufactured by:
Strides Pharma Science Ltd.
Bengaluru - 562106, India.

Distributed by:
Strides Pharma Inc.
East Brunswick,
NJ 08816

LOT: _____
Exp: _____

NO VARNISH ZONE

Revised: 03/2023
Code No: K220106K1705416/01

F.P.O


Ibuprofen
Liquid Gels
200 mg
 Solubilized Ibuprofen Capsules, 200 mg
 Pain Reliever/Fever Reducer (NSAID)

1X0000X

READ AND KEEP CARTON FOR COMPLETE WARNINGS AND INFORMATION

Drug Facts

Active ingredient (in each capsule)	Purpose
Solubilized ibuprofen equal to 200 mg ibuprofen (NSAID)* (present as the free acid and potassium salt) *nonsteroidal anti-inflammatory drug	Pain reliever/ Fever reducer

Uses

- temporarily relieves minor aches and pains due to:
 - headache
 - backache
 - the common cold
 - minor pain of arthritis
 - temporarily reduces fever
 - toothache
 - menstrual cramps
 - muscular aches

Warnings

Allergy alert: Ibuprofen may cause a severe allergic reaction, especially in people allergic to aspirin.

Symptoms may include:

- hives
- asthma (wheezing)
- skin redness
- blisters
- facial swelling
- shock
- itch

If an allergic reaction occurs, stop use and seek medical help right away.

Stomach bleeding warning: This product contains an NSAID, which may cause severe stomach bleeding. The chance is higher if you

- are age 60 or older
- have had stomach ulcers or bleeding problems
- take a blood thinning (anticoagulant) or steroid drug
- take other drugs containing prescription or non-prescription NSAIDs (aspirin, ibuprofen, naproxen, or others)
- have 3 or more alcoholic drinks every day while using this product
- take more or for a longer time than directed

Heart attack and stroke warning: NSAIDs, except aspirin, increase the risk of heart attack, heart failure, and stroke. These can be fatal. The risk is higher if you use more than directed or for longer than directed.

Drug Facts (continued)

Do not use

- if you have ever had an allergic reaction to any other pain reliever/fever reducer
- right before or after heart surgery

Ask a doctor before use if

- stomach bleeding warning applies to you
- you have problems or serious side effects from taking pain relievers or fever reducers
- you have a history of stomach problems, such as heartburn
- you have high blood pressure, heart disease, liver cirrhosis, kidney disease, asthma, or had a stroke
- you are taking a diuretic

Ask a doctor or pharmacist before use if you are

- under a doctor's care for any serious condition
- taking aspirin for heart attack or stroke, because ibuprofen may decrease the benefit of aspirin
- taking any other drug

When using this product

- take with food or milk if stomach upset occurs

Stop use and ask a doctor if

- you experience any of the following signs of stomach bleeding:
 - feel faint
 - have bloody or black stools
 - have stomach pain that does not get better
- you have symptoms of heart problems or stroke:
 - chest pain
 - weakness in one part or side of body
 - slurred speech
 - leg swelling
- pain gets worse or lasts more than 10 days
- fever gets worse or lasts more than 3 days
- redness or swelling is present in the painful area
- any new symptoms appear


 N 3 15 9 5 5 6 1 7 6 4 1 3 4

Drug Facts (continued)

If pregnant or breast-feeding, ask a health professional before use. It is especially important not to use ibuprofen at 20 weeks or later in pregnancy unless definitely directed to do so by a doctor because it may cause problems in the unborn child or complications during delivery.

Keep out of reach of children. In case of overdose, get medical help or contact a Poison Control Center right away.

Directions

- do not take more than directed
- the smallest effective dose should be used
- adults and children 12 years and over: take 1 capsule every 4 to 6 hours while symptoms persist
- if pain or fever does not respond to 1 capsule, 2 capsules may be used
- do not exceed 6 capsules in 24 hours, unless directed by a doctor
- children under 12 years: ask a doctor

Other information

- each capsule contains potassium 17.840 mg
- read all warnings and directions before use. Keep carton.
- store at 20-25°C (68-77°F)
- avoid excessive heat above 40°C (104°F)

Inactive ingredients

FD&C blue 1, gelatin, gelatin (hydrolyzed), medium chain triglyceride, PEG 600, potassium hydroxide, purified water, sorbitol and sorbitan solution, printing ink: propylene glycol, shellac resins, sodium lauryl sulphate, titanium dioxide.

Questions or comments? call toll free 1-877-244-9825

Do Not Use if seal under bottle cap imprinted with "SEALED for YOUR PROTECTION" is broken or missing.

For most recent product information
 Visit us at www.strides.com

Code No.: KR/DRUGS/KTK/25/415/98
 Revised: 05/2023
 Manufactured by: Strides Pharma Science Ltd. Bengaluru - 562106, India.
 Distributed by: Strides Pharma Inc, East Brunswick, NJ 08816

NDC 59556-764-13

Ibuprofen Liquid Gels

200 mg

Solubilized Ibuprofen Capsules, 200 mg
Pain Reliever/Fever Reducer (NSAID)

80 Liquid Filled Capsules

 Strides Pharma Inc.

NDC 59556-764-13

Ibuprofen Liquid Gels

200 mg

Solubilized Ibuprofen Capsules, 200 mg
Pain Reliever/Fever Reducer (NSAID)

80 Liquid Filled Capsules

 Strides Pharma Inc.

READ AND KEEP CARTON FOR COMPLETE WARNINGS AND INFORMATION

Do Not Use if seal under bottle cap imprinted with "SEALED for YOUR PROTECTION" is broken or missing.

Uses

- temporarily relieves minor aches and pains due to: headache, muscular aches, menstrual cramps, the common cold, backache, toothache, minor pain of arthritis.
- temporarily reduces fever

Warnings

- Ask your doctor before use if you are pregnant, under a doctor's care for a serious

F.P.O

Code No.: KR/DRUGS/KTK/25/415/98

LOT: _____

Exp.: _____

NO VARNISH ZONE

PEEL HERE

Ibuprofen Capsules, 200 mg Pain Reliever/Fever Reducer (NSAID)
Liquid Filled Capsules - Minis



Solubilized Ibuprofen Capsules, 200 mg
Pain Reliever/Fever Reducer (NSAID)

Ibuprofen Liquid Gels

100000X

READ AND KEEP CARTON FOR COMPLETE WARNINGS AND INFORMATION

Drug Facts

Active ingredient (in each capsule) Solubilized ibuprofen equal to 200 mg ibuprofen (NSAID)*
Purpose Pain reliever/ Fever reducer (present as the free acid and potassium salt) *nonsteroidal anti-inflammatory drug

Uses

- temporarily relieves minor aches and pains due to:
 - headache
 - toothache
 - backache
 - menstrual cramps
 - the common cold
 - muscular aches
 - minor pain of arthritis
 - temporarily reduces fever

Warnings

Allergy alert: Ibuprofen may cause a severe allergic reaction, especially in people allergic to aspirin.

Symptoms may include:

- hives
- facial swelling
- asthma (wheezing)
- shock
- skin reddening
- rash
- blisters

If an allergic reaction occurs, stop use and seek medical help right away.

Stomach bleeding warning: This product contains an NSAID, which may cause severe stomach bleeding. The chance is higher if you

- are age 60 or older
- have had stomach ulcers or bleeding problems
- take a blood thinning (anticoagulant) or steroid drug
- take other drugs containing prescription or nonprescription NSAIDs [aspirin, (ibuprofen, naproxen, or others)]
- have 3 or more alcoholic drinks every day while using this product
- take more or for a longer time than directed

Heart attack and stroke warning: NSAIDs, except aspirin, increase the risk of heart attack, heart failure, and stroke. These can be fatal. The risk is higher if you use more than directed or for longer than directed.

Drug Facts (continued)

Do not use

- if you have ever had an allergic reaction to any other pain reliever/fever reducer
- right before or after heart surgery

Ask a doctor before use if

- stomach bleeding warning applies to you
- you have problems or serious side effects from taking pain relievers or fever reducers
- you have a history of stomach problems, such as heartburn
- you have high blood pressure, heart disease, liver cirrhosis, kidney disease, asthma, or had a stroke
- you are taking a diuretic

Ask a doctor or pharmacist before use if you are

- under a doctor's care for any serious condition
- taking aspirin for heart attack or stroke, because ibuprofen may decrease this benefit of aspirin
- taking any other drug

When using this product

- take with food or milk if stomach upset occurs

Stop use and ask a doctor if

- you experience any of the following signs of stomach bleeding:
 - feel faint
 - vomit blood
 - have bloody or black stools
 - have stomach pain that does not get better
- you have symptoms of heart problems or stroke:
 - chest pain
 - trouble breathing
 - weakness in one part or side of body
 - slurred speech
 - leg swelling
- pain gets worse or lasts more than 10 days
- fever gets worse or lasts more than 3 days
- redness or swelling is present in the painful area
- any new symptoms appear



N 3159556176413 4

Drug Facts (continued)

If pregnant or breast-feeding, ask a health professional before use. It is especially important not to use ibuprofen at 20 weeks or later in pregnancy unless definitely directed to do so by a doctor because it may cause problems in the unborn child or complications during delivery.

Keep out of reach of children. In case of overdose, get medical help or contact a Poison Control Center right away.

Directions

- do not take more than directed
- the smallest effective dose should be used
- adults and children 12 years and over: take 1 capsule every 4 to 6 hours while symptoms persist
- if pain or fever does not respond to 1 capsule, 2 capsules may be used
- do not exceed 6 capsules in 24 hours, unless directed by a doctor
- children under 12 years: ask a doctor

Other information

- each capsule contains: potassium 17.840 mg
- read all warnings and directions before use. Keep carton store at 20-25°C (68-77°F)
- avoid excessive heat above 40°C (104°F)

Inactive ingredients

FD&C blue 1, gelatin, gelatin (hydrolyzed), medium chain triglyceride, PEG 600, potassium hydroxide, purified water, sorbitol and sorbitan solution, Printing ink: propylene glycol, shellac resins, sodium lauryl sulphate, titanium dioxide.

Questions or comments? call toll free 1-877-244-9825

Do Not Use if seal under bottle cap imprinted with "SEALED for YOUR PROTECTION" is broken or missing.

For most recent product information Visit us at www.strides.com

Code No.: KR/DRUGS/KTK/25/415/98

Manufactured by: Strides Pharma Science Ltd. Bengaluru - 562106, India.

Revised: 05/2023

Distributed by: Strides Pharma Inc, East Brunswick, NJ 08816

NDC 59556-764-13

Ibuprofen Liquid Gels

200 mg

Solubilized Ibuprofen Capsules, 200 mg
Pain Reliever/Fever Reducer (NSAID)

80 Liquid Filled Capsules

Strides Pharma Inc.



READ AND KEEP CARTON FOR COMPLETE WARNINGS AND INFORMATION

Do Not Use if seal under bottle cap imprinted with "SEALED for YOUR PROTECTION" is broken or missing.

Uses

- temporarily relieves minor aches and pains due to: headache, muscular aches, menstrual cramps, the common cold, backache, toothache, minor pain of arthritis.
- temporarily reduces fever

Warnings

- Ask your doctor before use if you are pregnant, under a doctor's care for a serious condition,

Code No.: KR/DRUGS/KTK/25/415/98

LOT:

Exp.:

NO VARNISH ZONE

PEEL HERE

NEW STRIDES
SINCE 1970

NDC 59556-895-13



Ibuprofen Liquid Gels

200 mg

Solubilized Ibuprofen Capsules, 200 mg
Pain Reliever/Fever Reducer (NSAID)

80 Liquid Filled Capsules

Strides Pharma Inc.

NEW product from STRIDES PHARMA
NDC 59556-895-48
READ AND KEEP CARTON FOR COMPLETE WARNINGS AND INFORMATION

Ibuprofen Liquid Gels

200 mg
 Solubilized Ibuprofen Capsules, 200 mg
 Pain Reliever/Fever Reducer (NSAID)

160 Liquid Filled Capsules
 Strides Pharma Inc.

Do Not Use if seal under bottle cap imprinted with "SEALED for YOUR PROTECTION" is broken or missing.

Uses
 ■ temporarily relieves minor aches and pains due to: headache, muscular aches, menstrual cramps, the common cold, backache, toothache, minor pain of arthritis. ■ temporarily reduces fever

Warnings
 ■ Ask your doctor before use if you are pregnant, under a doctor's care for a serious condition, age 60 or over, taking any other drug or have stomach problems.
 ■ This product may cause a severe allergic reaction, especially in people allergic to aspirin. Symptoms may include: hives, facial swelling, asthma (wheezing), shock, skin reddening, rash, blisters. If an allergic reaction occurs, stop use and seek medical help right away. Do not use this product if you have ever had an allergic reaction to any pain reliever/fever reducer.

Stomach bleeding warning: This product contains a nonsteroidal anti-inflammatory drug (NSAID), which may cause severe stomach bleeding. The chance is higher if you ■ are age 60 or older ■ have had stomach ulcers or bleeding problems ■ take a blood thinning (anticoagulant) or steroid drug ■ take other drugs containing prescription or nonprescription NSAIDs [aspirin, ibuprofen, naproxen, or others] ■ have 3 or more alcoholic drinks every day while using this product ■ take more or for a longer time than directed.

Heart attack and stroke warning: NSAIDs, except aspirin, increase the risk of heart attack, heart failure, and stroke. These can be fatal. The risk is higher if you use more than directed or for longer than directed.

Directions
 ■ do not take more than directed ■ the smallest effective dose should be used ■ adults and children 12 years and over: take 1 capsule every 4 to 6 hours while symptoms persist ■ if pain or fever does not respond to 1 capsule, 2 capsules may be used ■ do not exceed 6 capsules in 24 hours, unless directed by a doctor ■ children under 12 years: ask a doctor.

Each capsule contains: potassium 17.840 mg

Questions or comments?
 call toll free 1-877-244-9825
 For most recent product information, visit www.strides.com

Manufactured by:
Strides Pharma Science Ltd.
 Bengaluru - 562106, India.

Distributed by:
Strides Pharma Inc.
 East Brunswick,
 NJ 08816

LOT:
 Exp:

NO VARNISH ZONE

Revised: 03/2023
 Code No.: NPURUSKTKZ646588

IBUPROFEN

ibuprofen capsule, liquid filled

Product Information

Product Type	HUMAN OTC DRUG	Item Code (Source)	NDC:59556-764
Route of Administration	ORAL		

Active Ingredient/Active Moiety

Ingredient Name	Basis of Strength	Strength
IBUPROFEN (UNII: WK2XYI10QM) (IBUPROFEN - UNII:WK2XYI10QM)	IBUPROFEN	200 mg

Inactive Ingredients

Ingredient Name	Strength
POTASSIUM HYDROXIDE (UNII: WZ H3C48M4T)	
POLYETHYLENE GLYCOL 600 (UNII: NL4J9F21N9)	
WATER (UNII: 059QF0KO0R)	
GELATIN (UNII: 2G86QN327L)	
SORBITOL (UNII: 506T60A25R)	
SORBITAN (UNII: 6O92ICV9RU)	
FD&C BLUE NO. 1 (UNII: H3R47K3TBD)	
MEDIUM-CHAIN TRIGLYCERIDES (UNII: C9H2L21V7U)	
PROPYLENE GLYCOL (UNII: 6DC9Q167V3)	
SHELLAC (UNII: 46N107B71O)	
SODIUM LAURYL SULFATE (UNII: 368GB5141J)	
TITANIUM DIOXIDE (UNII: 15FIX9V2JP)	

Product Characteristics

Color	GREEN (transparent green to bluish green)	Score	no score
Shape	OVAL (oblong shaped)	Size	8mm
Flavor		Imprint Code	
Contains			

Packaging

#	Item Code	Package Description	Marketing Start Date	Marketing End Date
1	NDC:59556-764-29	1 in 1 CARTON	11/16/2022	
1		20 in 1 BOTTLE; Type 0: Not a Combination Product		
2	NDC:59556-764-12	1 in 1 CARTON	11/16/2022	
2		40 in 1 BOTTLE; Type 0: Not a Combination Product		
3	NDC:59556-764-13	1 in 1 CARTON	11/16/2022	
3		80 in 1 BOTTLE; Type 0: Not a Combination Product		
4	NDC:59556-764-11	1 in 1 CARTON	11/16/2022	
4		120 in 1 BOTTLE; Type 0: Not a Combination Product		
5	NDC:59556-764-48	1 in 1 CARTON	11/16/2022	
5		160 in 1 BOTTLE; Type 0: Not a Combination Product		
6	NDC:59556-764-49	1 in 1 CARTON	11/16/2022	
6		200 in 1 BOTTLE; Type 0: Not a Combination Product		
7	NDC:59556-764-50	1 in 1 CARTON	11/16/2022	
7		300 in 1 BOTTLE; Type 0: Not a Combination Product		
8	NDC:59556-764-18	1 in 1 CARTON	11/16/2022	
8		180 in 1 BOTTLE; Type 0: Not a Combination Product		
9	NDC:59556-764-92	3000 in 1 BAG; Type 0: Not a Combination Product	11/16/2022	

Marketing Information

Marketing Category	Application Number or Monograph Citation	Marketing Start Date	Marketing End Date
ANDA	ANDA204469	11/16/2022	

IBUPROFEN

ibuprofen capsule, liquid filled

Product Information

Product Type	HUMAN OTC DRUG	Item Code (Source)	NDC:59556-895
Route of Administration	ORAL		

Active Ingredient/Active Moiety

Ingredient Name	Basis of Strength	Strength
IBUPROFEN (UNII: WK2XYI10QM) (IBUPROFEN - UNII:WK2XYI10QM)	IBUPROFEN	200 mg

Inactive Ingredients

Ingredient Name	Strength
POTASSIUM HYDROXIDE (UNII: WZH3C48M4T)	
POLYETHYLENE GLYCOL 600 (UNII: NL4J9F21N9)	
WATER (UNII: 059QF0KO0R)	
GELATIN (UNII: 2G86QN327L)	
SORBITOL (UNII: 506T60A25R)	
MEDIUM-CHAIN TRIGLYCERIDES (UNII: C9H2L21V7U)	
PROPYLENE GLYCOL (UNII: 6DC9Q167V3)	
SHELLAC (UNII: 46N107B71O)	
SODIUM LAURYL SULFATE (UNII: 368GB5141J)	
TITANIUM DIOXIDE (UNII: 15FIX9V2JP)	
FD&C GREEN NO. 3 (UNII: 3P3ONR6O1S)	
LECITHIN, SOYBEAN (UNII: 1DI56QDM62)	
SORBITAN (UNII: 6O92ICV9RU)	

Product Characteristics

Color	GREEN (transparent green to bluish green)	Score	no score
Shape	OVAL	Size	8mm
Flavor		Imprint Code	469
Contains			

Packaging

#	Item Code	Package Description	Marketing Start Date	Marketing End Date
1	NDC:59556-895-29	1 in 1 CARTON	11/16/2022	
1		20 in 1 BOTTLE; Type 0: Not a Combination Product		
2	NDC:59556-895-13	1 in 1 CARTON	11/16/2022	
2		80 in 1 BOTTLE; Type 0: Not a Combination Product		
3	NDC:59556-895-48	1 in 1 CARTON	11/16/2022	
3		160 in 1 BOTTLE; Type 0: Not a Combination Product		
4	NDC:59556-895-50	1 in 1 CARTON	11/16/2022	
4		300 in 1 BOTTLE; Type 0: Not a Combination Product		
5	NDC:59556-895-79	7000 in 1 BAG; Type 0: Not a Combination Product	11/16/2022	

6	NDC:59556-895-85	1 in 1 CARTON	11/16/2022	
6		80 in 1 BOTTLE; Type 0: Not a Combination Product		
7	NDC:59556-895-86	1 in 1 CARTON	11/16/2022	
7		120 in 1 BOTTLE; Type 0: Not a Combination Product		
8	NDC:59556-895-87	1 in 1 CARTON	11/16/2022	
8		160 in 1 BOTTLE; Type 0: Not a Combination Product		
9	NDC:59556-895-39	1 in 1 CARTON	11/16/2022	
9		200 in 1 BOTTLE; Type 0: Not a Combination Product		
10	NDC:59556-895-88	1 in 1 CARTON	11/16/2022	
10		300 in 1 BOTTLE; Type 0: Not a Combination Product		

Marketing Information

Marketing Category	Application Number or Monograph Citation	Marketing Start Date	Marketing End Date
ANDA	ANDA204469	11/16/2022	

Labeler - Strides Pharma Inc (078868278)

Revised: 4/2024

Strides Pharma Inc