

**PERDIEM- sennosides, stimulant laxative tablet**  
**Haleon US Holdings LLC**

-----  
**Drug Facts**

**Active Ingredient**

Sennosides 15 mg

**Purpose**

Stimulant Laxative

**Uses**

- relieves occasional constipation (irregularity)
- generally produces bowel movement in 6 to 12 hours

**Warnings**

**Do not use laxative product when abdominal pain, nausea or vomiting are present unless directed by a doctor**

**Ask a doctor or pharmacist before use if you**

- have noticed a sudden change in bowel habits that persists over a period of 2 weeks
- are taking any other drug. Take this product 2 or more hours before or after other drugs. Laxatives may affect how other drugs work.

**When using this product**

- do not use for a period longer than 1 week

**Stop use and ask a doctor if**

- rectal bleeding or failure to have a bowel movement occur after use of a laxative. These may be signs of a serious condition.

**If pregnant or breast-feeding, ask a health professional before use.**

**Keep out of reach of children. In case of overdose, get medical help or contact a Poison Control Center right away.**

**Directions**

- swallow tablet with a glass of water

- swallow tablet whole, do not crush, break or chew

Adults and children 12 years of age and older	2 tablets once or twice daily
Children 6 to under 12 years of age	1 tablet once or twice daily
Children under 6 years of age	ask a doctor

### **Other Information**

- **each tablet contains:**calcium 45mg, magnesium 5 mg
- store at controlled room temperature 20-25°C (68-77°F).

### **Inactive Ingredients**

acacia, calcium carbonate, carnuba wax, corn starch, dibasic calcium phosphate, iron oxide black, iron oxide red, iron oxide yellow, magnesium stearate, methylparaben, microcrystalline cellulose, polyethylene glycol, polyvinyl alcohol, povidone, propylene glycol, propylparaben, shellac, silicon dioxide, sodium benzoate, sodium lauryl sulfate, sucrose, talc, titanium dioxide

### **Questions?**

call **1-855-297-3036**

### **Package/Label Principal Display Panel**

OVERNIGHT RELIEF

# Perdiem®

Sennosides 15 mg,  
stimulant laxative

OVERNIGHT RELIEF

# Perdiem®

Sennosides 15 mg,  
stimulant laxative

A NATURAL  
VEGETABLE INGREDIENT  
No Mixing No Mess

60 Tablets



FORMULATED  
FOR DEPENDABLE OVERNIGHT RELIEF

Perdiem  
Perdie.

NDC 0057-8143-01

OVERNIGHT RELIEF

# Perdiem®

Sennosides 15 mg,  
stimulant laxative

A NATURAL  
VEGETABLE INGREDIENT  
No Mixing No Mess

60 Tablets



FORMULATED  
FOR DEPENDABLE OVERNIGHT RELIEF

Perdiem  
Perdie.

gsk

TAMPER-EVIDENT Feature. Do Not Use if  
inner seal imprinted with "SEALER FOR  
YOUR PROTECTION" is broken or missing.

Distributed by:  
GSK Consumer Healthcare, Warren, NJ 07059  
©2017 GSK group of companies or its licensor.  
All rights reserved. 12958

Trademarks are owned by or licensed to the GSK group of companies.

1-855-297-3036

### Drug Facts (cont. tued)

**Inactive ingredients** acetate, calcium carbonate, carnuba wax, corn starch, dibasic calcium

phosphate, iron oxide black, iron oxide red, iron oxide yellow, magnesium aspartate, methylparaben,

microcrystalline cellulose, polyethylene glycol, polyvinyl alcohol, polydioxane, propylene glycol, propylparaben,

stibac, silicon dioxide, sodium benzoate, sodium lauryl sulfate, sucrose, talc, titanium dioxide

**Questions or comments?** call 1-855-297-3036

### Other information

• each tablet contains calcium 45 mg, magnesium 5 m

• store at controlled room temperature 20-25°C (68-77°F)

### Directions

• swallow tablet(s) with a glass of water

• adults and children 12 years of age and older

2 tablets once or twice daily

children 6 to under 12 years of age

1 tablet once or twice daily

ask a doctor

in pregnant or breast-feeding, ask a health professional before use.  
Keep out of reach of children. In case of overdose, get medical help or contact a Poison Control Center right away.

**Drug Facts**

**Active Ingredient (in each tablet)** Sennosides 15 mg

**Purpose** Stimulant laxative

**Uses**

- relieves occasional constipation (irregularity)
- generally produces bowel movement in 6 to 12 hours

**Warnings**

Do not use laxative products when abdominal pain, nausea or vomiting are present unless directed by a doctor.

Ask a doctor before use if you have noticed a sudden change in bowel habits that persists over a period of 2 weeks.

Ask a doctor or pharmacist before use if you are taking any other drug. Laxatives may affect how other drugs work. Take this product 2 or more hours before or after other drugs.

When using this product do not use for a period longer than 1 week.

Stop use and ask a doctor if rectal bleeding or failure to have a bowel movement occur after use of a laxative. These may be signs of a serious condition.

## Primary Label

**Drug Facts (continued)**

If pregnant or breast-feeding, ask a health professional before use. Keep out of reach of children. In case of overdose, get medical help or contact a Poison Control Center right away.

**Directions**

- swallow tablet(s) with a glass of water
- swallow tablet(s) whole, do not crush, break or chew

adults and children 12 years of age and older 2 tablets once or twice daily

children 6 to under 12 years of age 1 tablet once or twice daily

children under 6 years of age ask a doctor

**Other information**

- each tablet contains: calcium 45 mg, magnesium 5 mg
- store at controlled room temperature 20-25°C (68-77°F)

**Inactive ingredients** acacia, calcium carbonate, carnauba wax, corn starch, dibasic calcium phosphate, iron oxide black, iron oxide red, iron oxide yellow, magnesium stearate, methylparaben, microcrystalline cellulose, polyethylene glycol, polyvinyl alcohol, povidone, propylene glycol, propylparaben, triethylaluminum dioxide, sodium benzoate, sodium lauryl sulfate, sucrose, talc, titanium dioxide

**Questions or comments?**  
Call 1-855-297-3036

1-855-297-3036

Distributed by:  
GSK Consumer Healthcare  
Warren, NJ 07059  
©2017 GSK group of companies or its licensor. All rights reserved.

## Primary Label

NDC 0087-8143-01

OVERNIGHT RELIEF

**Perdiem**<sup>®</sup>

Sennosides 15 mg, stimulant laxative

A NATURAL VEGETABLE INGREDIENT

No Mixing No Mess

60 Tablets

FORMULATED FOR PAINKIND OVERNIGHT RELIEF

Do not detach this front label. It contains important information.

**Tamper-Evident Feature:** Protective printed liner seal over each cap. If missing or damaged, do not use contents.

**The Perdiem<sup>®</sup> Guarantee:** When taken at bedtime, Perdiem<sup>®</sup> is guaranteed to gently, effectively or your money back. Return product to GSK, Attention Consumer Affairs, for full refund.

**Drug Facts**

**Active ingredient (in each tablet)** Sennosides 15 mg

**Purpose** Stimulant laxative

**Uses**

- relieves occasional constipation (irregularity)
- generally produces bowel movement in 6 to 12 hours

**Warnings**

Do not use laxative products when abdominal pain, nausea or vomiting are present unless directed by a doctor

Ask a doctor before use if you have noticed a sudden change in bowel habits that persists over a period of 2 weeks

Ask a doctor or pharmacist: before use if you are taking any other drug. Laxatives may affect how other drugs work. Take this product 2 or more hours before or after other drugs.

When using this product do not use for a period longer than 1 week

Stop use and ask a doctor if rectal bleeding or failure to have a bowel movement occur after use of a laxative. These may be signs of a serious condition.

12957 0630-506-11

PEEL HERE FOR MORE DRUG FACTS

## Primary Label



## PERDIEM

sennosides, stimulant laxative tablet

### Product Information

<b>Product Type</b>	HUMAN OTC DRUG	<b>Item Code (Source)</b>	NDC:0067-8143
<b>Route of Administration</b>	ORAL		

### Active Ingredient/Active Moiety

Ingredient Name	Basis of Strength	Strength
<b>SENNOSIDES</b> (UNII: 3FYP5M0IJX) (SENNOSIDES - UNII:3FYP5M0IJX)	SENNOSIDES	15 mg

### Inactive Ingredients

Ingredient Name	Strength
<b>ACACIA</b> (UNII: 5C5403N26O)	
<b>CALCIUM CARBONATE</b> (UNII: H0G9379FGK)	
<b>CARNAUBA WAX</b> (UNII: R12CBM0EIZ)	
<b>STARCH, CORN</b> (UNII: O8232NY3SJ)	
<b>ANHYDROUS DIBASIC CALCIUM PHOSPHATE</b> (UNII: L11K75P92J)	
<b>FERROSFERRIC OXIDE</b> (UNII: XM0M87F357)	
<b>FERRIC OXIDE RED</b> (UNII: 1K09F3G675)	
<b>FERRIC OXIDE YELLOW</b> (UNII: EX438O2MRT)	
<b>MAGNESIUM STEARATE</b> (UNII: 70097M6I30)	
<b>METHYLPARABEN</b> (UNII: A2I8C7HI9T)	
<b>MICROCRYSTALLINE CELLULOSE</b> (UNII: OP1R32D61U)	
<b>POLYETHYLENE GLYCOL 3350</b> (UNII: G2M7P15E5P)	
<b>POLYVINYL ALCOHOL, UNSPECIFIED</b> (UNII: 532B59J990)	
<b>POVIDONE, UNSPECIFIED</b> (UNII: FZ989GH94E)	
<b>PROPYLENE GLYCOL</b> (UNII: 6DC9Q167V3)	
<b>PROPYLPARABEN</b> (UNII: Z8IX2SC1OH)	
<b>SHELLAC</b> (UNII: 46N107B71O)	
<b>SILICON DIOXIDE</b> (UNII: ETJ7Z6XBU4)	
<b>SODIUM BENZOATE</b> (UNII: OJ245FE5EU)	
<b>SODIUM LAURYL SULFATE</b> (UNII: 368GB5141J)	
<b>SUCROSE</b> (UNII: C151H8M554)	
<b>TALC</b> (UNII: 7SEV7J4R1U)	
<b>TITANIUM DIOXIDE</b> (UNII: 15FIX9V2JP)	

## Product Characteristics

<b>Color</b>	brown (Tan)	<b>Score</b>	no score
<b>Shape</b>	ROUND	<b>Size</b>	10mm
<b>Flavor</b>		<b>Imprint Code</b>	Perdiem
<b>Contains</b>			

## Packaging

#	Item Code	Package Description	Marketing Start Date	Marketing End Date
1	NDC:0067-8143-01	1 in 1 CARTON	07/01/2017	
1		60 in 1 BOTTLE; Type 0: Not a Combination Product		

## Marketing Information

Marketing Category	Application Number or Monograph Citation	Marketing Start Date	Marketing End Date
OTC Monograph Drug	M007	07/01/2017	

**Labeler** - Haleon US Holdings LLC (079944263)

Revised: 2/2024

Haleon US Holdings LLC