

**COLD AND SINUS MAXIMUM STRENGTH- acetaminophen, guaifenesin,
phenylephrine hydrochloride liquid
DeMoulas Market Basket**

Cold & Sinus 515

Drug Facts

**Active ingredients
(in each 20 mL)**

Acetaminophen 650 mg
Guaifenesin 400 mg
Phenylephrine HCL 10 mg

Purpose

Pain reliever/Fever reducer
Expectorant
Nasal decongestant

Uses

temporarily relieves these common cold and flu symptoms:

- nasal congestion
- sinus congestion and pressure
- minor aches and pains
- sore throat
- headache
- fever
- helps loosen phlegm (mucus) and thin bronchial secretions to drain bronchial tubes

Warnings

Liver warning This product contains acetaminophen. Severe liver damage may occur if you take:

- more than 4,000 mg of acetaminophen in 24 hours
- with other drugs containing acetaminophen
- 3 or more alcoholic drinks every day while using this product

Allergy alert: Acetaminophen may cause severe skin reactions, Symptoms may include:

- skin reddening
- blisters
- rash

If a skin reaction occurs, stop use and seek medical help right away.

Sore throat warning If sore throat is severe, persists for more than 2 days, is accompanied or followed by fever, headache, rash, nausea or vomiting, consult a doctor promptly.

Do not use

- with any other drug containing acetaminophen (prescription or nonprescription). If you are not sure whether a drug contains acetaminophen, ask a doctor or pharmacist.
- if you are now taking a prescription monoamine oxidase inhibitor (MAOI) (certain drugs for depression, psychiatric or emotional conditions, or Parkinson's disease), or for 2 weeks after stopping the MAOI drug. If you do not know if your prescription drug contains an MAOI, ask a doctor or pharmacist before taking this product.
- for children under 12 years of age

Ask a doctor before use if you have

- liver disease
- heart disease
- thyroid disease
- high blood pressure
- diabetes
- trouble urinating due to an enlarged prostate gland
- a breathing problem or chronic cough that lasts or as occurs with smoking, asthma or emphysema
- cough that occurs with too much phlegm (mucus)

Ask a doctor or pharmacist before use if you are

- taking the blood thinning product warfarin.

When using this product

- **do not use more that directed**

stop use and ask a doctor if

- nervousness, dizziness or sleeplessness occur
- redness or swelling is present
- symptoms do not get better within 7 days or are accompanied by a fever
- fever gets worse or lasts more than 3 days
- new symptoms occur
- cough lasts more than 7 days, comes back, or occurs with fever, rash or persistent headache. These could be signs of a serious condition.

If pregnant or breast feeding,

ask a health professional before use.

Keep out of reach of children.

Keep out of reach of children.

In case of overdose, get medical help or contact a Poison Control Center right away.

Overdose warning

Taking more than the recommended dose (overdose) may cause liver damage. In case of overdose, get medical help or contact a Poison Control Center right away. Quick medical attention is critical for adults as well as for children even if you do not notice any signs or symptoms.

Directions

- take only as recommended (see Overdose warning)
- use dosage cup
- mL = milliliter
- do not take more than 6 doses in any 24-hour period

Age	Dose
Adults & children 12 years and older	20 mL every 4 hours
Children under 12 years of age	Do not use

Other information

- each 20 mL contains: sodium 10 mg
- dosage cup provided
- store between 15-30 C (59-86 F)
- do not refrigerate

Inactive ingredients

anhydrous citric acid, edetate disodium, FD&C Blue #1, FD&C Red #40, flavors, glycerin, propylene glycol, propyl gallate, purified water, sodium benzoate, sorbitol, sucralose, Xanthan gum

Questions?

Call weekdays from 9:30 AM to 4:30 PM EST 1-877-798-5944

Cold and Sinus product label

Market Basket®

"MORE FOR YOUR MONEY"

* COMPARE TO

the active ingredients found in
MUCINEX® FAST-MAX™
COLD & SINUS

**Cold & Sinus
MAXIUN STRENGTH**

Acetaminophen

(Pain Reliever/Fever Reducer)

Guiafenesisin (Expectorant)

**Phenylephrine HCL
(Nasal Decongestant)**

**Relieves Fever & Body Aches
Thins & Loosens Mucus
Relieves Nasal & Chest Congestion**

Adults For Ages 12 & Over

LF-009 Rev 01

6 FL OZ (177 mL)


Peel Corner to Read Completed Drug Facts and information

DO NOT USE IF PRINTED SEAL UNDER CAP IS TORN OR MISSING

Failure to follow these warnings could result in serious consequences.

DISTRIBUTED BY
DEMOULAS SUPERMARKETS INC.
875 East Street
TEWKSBURY, MA 01876

* This product is not manufactured or distributed by Reckitt Benckiser Inc. distributor of Mucinex® FAST-MAX™ COLD & SINUS



Peel Corner to Read Complete
Drug Facts and Information



**DO NOT USE IF PRINTED SEAL
UNDER CAP IS TORN OR MISSING**
Failure to follow these warnings
could result in serious
consequences

DISTRIBUTED BY
DEMOULAS SUPERMARKETS INC.
876 East Street
TEWKSBURY, MA 01876
LB-009 Rev 01



Lot:
Exp:

Drug Facts

Active ingredients (in each 20 mL)

Acetaminophen 650 mg..... Pain reliever/Fever reducer
 Guaifenesin 400 mg..... Expectorant
 Phenylephrine HCl 10 mg..... Nasal decongestant

Uses

temporarily relieves these common cold and flu symptoms:
 ■ nasal congestion
 ■ sinus congestion and pressure
 ■ minor aches and pains
 ■ sore throat ■ headache ■ fever
 ■ helps loosen phlegm (mucus) and thin bronchial secretions to drain bronchial tubes

Warnings

Liver warning This product contains acetaminophen. Severe liver damage may occur if you take:
 ■ more than 4,000 mg of acetaminophen in 24 hours
 ■ with other drugs containing acetaminophen
 ■ 3 or more alcoholic drinks every day while using this product
Allergy alert: Acetaminophen may cause severe skin reactions. Symptoms may include:
 ■ skin reddening
 ■ blisters
 ■ rash

Drug Facts (continued)

If a skin reaction occurs, stop use and seek medical help right away.

Sore throat warning If sore throat is severe, persists for more than 2 days, is accompanied or followed by fever, headache, rash, nausea or vomiting, consult a doctor promptly.

Do not use ■ with any other drug containing acetaminophen (prescription or nonprescription). If you are not sure whether a drug contains acetaminophen, ask a doctor or pharmacist.

■ if you are now taking a prescription monoamine oxidase inhibitor (MAOI) (certain drugs for depression, psychiatric, or emotional conditions, or Parkinson's disease), or for 2 weeks after stopping the MAOI drug. If you do not know if your prescription drug contains an MAOI, ask a doctor or pharmacist before taking this product.

■ for children under 12 years of age

Ask a doctor before use if you have ■ liver disease
 ■ heart disease
 ■ thyroid disease
 ■ high blood pressure
 ■ diabetes

Drug Facts (continued)

- trouble urinating due to an enlarged prostate gland
- a breathing problem or chronic cough that lasts or as occurs with smoking, asthma or emphysema
- cough that occurs with too much phlegm (mucus)

Ask a doctor or pharmacist before use if you are ■ taking the blood thinning drug warfarin.

When using this product ■ do not use more than directed

Stop use and ask a doctor if
 ■ nervousness, dizziness or sleeplessness occur
 ■ redness or swelling is present
 ■ symptoms do not get better within 7 days or are accompanied by a fever
 ■ fever gets worse or lasts more than 3 days
 ■ new symptoms occur
 ■ cough lasts more than 7 days, comes back, or occurs with fever, rash or persistent headache. These could be signs of a serious condition.

If pregnant or breast feeding, ask a health professional before use.

Keep out of reach of children. In case of overdose, get medical help or contact a Poison Control Center right away.

Overdose warning Taking more than the recommended dose (overdose) may cause liver damage.

Drug Facts (continued)

In case of overdose, get medical help or contact a Poison Control Center right away. Quick medical attention is critical for adults as well as for children even if you do not notice any signs or symptoms.

Directions

■ take only as recommended (see Overdose warning) ■ use dosage cup
 ■ mL = milliliter ■ do not take more than 6 doses in any 24-hour period

Age	Dose
Adults & children 12 years and older	20 mL every 4 hours
Children under 12 years of age	Do not use

Other information

■ each 20 mL contains: sodium 10 mg
 ■ dosage cup provided
 ■ store between 15-30°C (59-86°F)
 ■ do not refrigerate

Inactive ingredients

anhydrous citric acid, edetate disodium, FD&C Blue #1, FD&C Red #40, flavors, glycerin, propylene glycol, propyl gallate, purified water, sodium benzoate, sorbitol, sucralose, Xanthan gum

Questions?

Call weekdays from 9:30 AM to 4:30 PM EST 1-877-798-5944

*This product is not manufactured or distributed by Reckitt Benckiser Inc., distributor of Mucinex® FAST-MAX™ Cold & Sinus.

COLD AND SINUS MAXIUM STRENGTH

acetaminophen, guaifenesin, phenylephrine hydrochloride liquid

Product Information

Product Type

HUMAN OTC DRUG

Item Code (Source)

NDC:53942-515

Route of Administration ORAL

Active Ingredient/Active Moiety

Ingredient Name	Basis of Strength	Strength
ACETAMINOPHEN (UNII: 362O9ITL9D) (ACETAMINOPHEN - UNII:362O9ITL9D)	ACETAMINOPHEN	650 mg in 20 mL
GUAIFENESIN (UNII: 495W7451VQ) (GUAIFENESIN - UNII:495W7451VQ)	GUAIFENESIN	400 mg in 20 mL
PHENYLEPHRINE HYDROCHLORIDE (UNII: 04JA59TNSJ) (PHENYLEPHRINE - UNII:1WS297W6MV)	PHENYLEPHRINE	10 mg in 20 mL

Inactive Ingredients

Ingredient Name	Strength
ANHYDROUS CITRIC ACID (UNII: XF417D3PSL)	
EDETATE DISODIUM (UNII: 7FLD91C86K)	
FD&C BLUE NO. 1 (UNII: H3R47K3TBD)	
FD&C RED NO. 40 (UNII: WZB9127XOA)	
GLYCERIN (UNII: PDC6A3C0OX)	
PROPYLENE GLYCOL (UNII: 6DC9Q167V3)	
PROPYL GALLATE (UNII: 8D4SNN7V92)	
WATER (UNII: 059QF0KO0R)	
SODIUM BENZOATE (UNII: OJ245FE5EU)	
SORBITOL (UNII: 506T60A25R)	
SUCRALOSE (UNII: 96K6UQ3ZD4)	
XANTHAN GUM (UNII: TTV12P4NEE)	

Packaging

#	Item Code	Package Description	Marketing Start Date	Marketing End Date
1	NDC:53942-515-25	177 mL in 1 BOTTLE; Type 0: Not a Combination Product	09/19/2013	

Marketing Information

Marketing Category	Application Number or Monograph Citation	Marketing Start Date	Marketing End Date
OTC Monograph Drug	M012	09/19/2013	

Labeler - DeMoulas Market Basket (007869647)

Registrant - AptaPharma Inc. (790523323)

Establishment

Name	Address	ID/FEI	Business Operations
------	---------	--------	---------------------

AptaPharma Inc.

790523323

manufacture(53942-515)

Revised: 12/2023

DeMoulas Market Basket