

ANTACID- aluminum hydroxide, magnesium hydroxide, simethicone suspension

Meijer Distribution Inc

Disclaimer: Most OTC drugs are not reviewed and approved by FDA, however they may be marketed if they comply with applicable regulations and policies. FDA has not evaluated whether this product complies.

Meijer Distribution, Inc. Antacid Drug Facts

Active ingredients (in each 10 mL)

Aluminum hydroxide (equiv. to dried gel, USP) 800 mg

Magnesium hydroxide 800 mg

Simethicone 80 mg

Purposes

Antacid

Antigas

Uses

relieves

- heartburn
- acid indigestion
- sour stomach
- upset stomach due to these symptoms
- pressure and bloating commonly referred to as gas

Warnings

Ask a doctor before use if you have

- kidney disease
- a magnesium-restricted diet

Ask a doctor or pharmacist before use if you are

presently taking a prescription drug. Antacids may interact with certain prescription drugs.

When using this product

do not exceed 60 mL in a 24-hour period, or use the maximum dosage for more than 2 weeks

Keep out of reach of children.

In case of overdose, get medical help or contact a Poison Control Center right away (1-800-222-1222).

Directions

- shake well before use

- measure with dosing cup provided
- adults and children 12 years and over: 10 mL – 20 mL between meals, at bedtime or as directed by a doctor
- do not take more than 60 mL in any 24 hour period
- do not use the maximum dosage for more than 2 weeks
- children under 12 years: ask a doctor
- mL = milliliter

Other information

- **each 10 mL contains:** magnesium 350 mg, potassium 20 mg and sodium 3 mg
- **does not meet USP requirements for preservative effectiveness**
- store at 20-25°C (68-77°F), do not freeze

Inactive ingredients

butylparaben, flavor, hypromellose, microcrystalline cellulose and carboxymethylcellulose sodium, potassium citrate, propylparaben, purified water, simethicone emulsion, sorbitol, sorbitol solution

Questions or comments?

1-800-719-9260

Principal Display Panel

Compare to Mylanta® Maximum Strength active ingredients

MAXIMUM STRENGTH

antacid

Antacid | Anti-Gas

RELIEF OF:

Heartburn, Indigestion, Sour Stomach

Classic Flavor

Works on Contact

Tough on Heartburn

12 FL OZ (355 mL)

meijer[®]

NDC 41250-340-40

Compare to Mylanta[®]
Maximum Strength
active ingredients*

MAXIMUM STRENGTH
antacid

Antacid | Anti-Gas

RELIEF OF:
Heartburn, Indigestion, Sour Stomach

Classic Flavor



Works on Contact
Tough on Heartburn

12 FL OZ (355 mL)

: 34040 6E F11

**DO NOT USE IF
PRINTED NECKBAND
IS BROKEN OR
MISSING**

*This product is not manufactured or distributed by McNeil Consumer Pharmaceuticals Co., owner of the registered trademark Mylanta®.



PID 463704

0 41250 52115 0

Drug Facts

Active ingredients (in each 10 mL)	Purposes
Aluminum hydroxide (equiv. to dried gel, USP) 800 mg.....	Antacid
Magnesium hydroxide 800 mg.....	Antacid
Simethicone 80 mg.....	Antigas

Uses relieves ■ heartburn ■ acid indigestion ■ sour stomach
 ■ upset stomach due to these symptoms
 ■ pressure and bloating commonly referred to as gas

Warnings
Ask a doctor before use if you have ■ kidney disease ■ a magnesium-restricted diet
Ask a doctor or pharmacist before use if you are presently taking a prescription drug. Antacids may interact with certain prescription drugs.

When using this product do not exceed 60 mL in a 24-hour period, or use the maximum dosage for more than 2 weeks
Keep out of reach of children. In case of overdose, get medical help or contact a Poison Control Center right away (1-800-222-1222).

Directions ■ shake well before use ■ measure with dosing cup provided
 ■ **adults and children 12 years and over:** 10 mL – 20 mL between meals, at bedtime or as directed by a doctor
 ■ do not take more than 60 mL in any 24 hour period
 ■ do not use the maximum dosage for more than 2 weeks
 ■ **children under 12 years:** ask a doctor ■ mL = milliliter

Other information
 ■ each 10 mL contains: magnesium 350 mg, potassium 20 mg and sodium 3 mg
 ■ does not meet USP requirements for preservative effectiveness
 ■ store at 20-25°C (68-77°F), do not freeze

Inactive ingredients butylparaben, flavor, hypromellose, microcrystalline cellulose and carboxymethylcellulose sodium, potassium citrate, propylparaben, purified water, simethicone emulsion, sorbitol, sorbitol solution ▶

Drug Facts
(continued)
Questions or comments?
 1-800-719-9260

DIST. BY MEIJER DISTRIBUTION, INC.
 GRAND RAPIDS, MI 49544
 www.meijer.com



: 34040 6E B12

ANTACID

aluminum hydroxide, magnesium hydroxide, simethicone suspension

Product Information

Product Type	HUMAN OTC DRUG	Item Code (Source)	NDC:41250-340
Route of Administration	ORAL		

Active Ingredient/Active Moiety

Ingredient Name	Basis of Strength	Strength
ALUMINUM HYDRO XIDE (UNII: 5QB0T2IUN0) (ALUMINUM HYDRO XIDE - UNII:5QB0T2IUN0)	ALUMINUM HYDRO XIDE	800 mg in 10 mL
MAGNESIUM HYDRO XIDE (UNII: NBZ3QY004S) (MAGNESIUM CATION - UNII:T6V3LHY838, HYDRO XIDE ION - UNII:9159UV381P)	MAGNESIUM HYDRO XIDE	800 mg in 10 mL
DIMETHICONE (UNII: 92RU3N3Y1O) (DIMETHICONE - UNII:92RU3N3Y1O)	DIMETHICONE	80 mg in 10 mL

Inactive Ingredients

Ingredient Name	Strength
SILICON DIOXIDE (UNII: ETJ7Z6XBU4)	
BUTYLPARABEN (UNII: 3QPIIU3FV8)	
HYPROMELLOSES (UNII: 3NXW29V3WO)	
CELLULOSE, MICROCRYSTALLINE (UNII: OP1R32D61U)	
CARBOXYMETHYLCELLULOSE SODIUM (UNII: K679OBS311)	
POTASSIUM CITRATE (UNII: EE90ONI6FF)	
PROPYLPARABEN (UNII: Z8IX2SC1OH)	
WATER (UNII: 059QF0KO0R)	
SORBITOL (UNII: 506T60A25R)	

Product Characteristics

Color	WHITE (opaque)	Score	
Shape		Size	
Flavor	LEMON	Imprint Code	
Contains			

Packaging

#	Item Code	Package Description	Marketing Start Date	Marketing End Date
1	NDC:41250-340-40	355 mL in 1 BOTTLE; Type 0: Not a Combination Product	08/15/1990	

Marketing Information

Marketing Category	Application Number or Monograph Citation	Marketing Start Date	Marketing End Date
OTC monograph final	part331	08/15/1990	

Labeler - Meijer Distribution Inc (006959555)

Revised: 11/2018

Meijer Distribution Inc