

ORAL-B MINUTE-FOAM GRAPE PUNCH - acidulated phosphate fluoride aerosol

Oral-B Laboratories

Disclaimer: This drug has not been found by FDA to be safe and effective, and this labeling has not been approved by FDA. For further information about unapproved drugs, click here.

Package Label Principal Display Panel

MINUTE-FOAM

Topical Fluoride Foaming Solution

(Acidulated Phosphate Fluoride Containing 1.23 % w/w Fluoride Ion)

FOR PROFESSIONAL USE ONLY

Active Ingredient: Acidulated Phosphate Fluoride (Sodium Fluoride and Hydrofluoric Acid) 1.23 % w/w fluoride ion.

DESCRIPTION: Oral-B Minute Foam is a flavored aqueous foaming solution of acidulated phosphate sodium fluoride with pH of 3.5 and a fluoride ion concentration of 1.23% w/w.

Inactive Ingredients: Purified water, poloxamer 407, isobutane, poloxamer 234, phosphoric acid, sodium saccharin, flavor.

CLINICAL PHARMACOLOGY: Acidulated Phosphate Fluoride inhibits caries formation by reducing enamel solubility and enhancing remineralization.

INDICATION AND USE: A topically-applied foaming solution to aid in the prevention of dental caries.

CONTRAINDICATIONS: Do not use in patients with hypersensitivity to fluoride.
Do not use in patients with dysphagia.

WARNINGS: DO NOT SWALLOW

Accidental ingestion of the usual treatment dose (approx 12.4 mg of fluoride) is not harmful. In the event more than the treatment dose is swallowed, administer calcium (e.g. milk) and get medical or contact a Poison control Center right away. One bottle of Minute Foam contained 1.95 grams of fluoride ion which could be lethal for children adults.

Keep out of the reach of infants and children under 12 years.

Pediatric patients under age 12 should be supervised during use of this product.

Avoid spraying towards open flame.

Contents under pressure. Do not puncture or incinerate. Do not expose to heat or store at temperatures over 120 F (49 C).

PRECAUTIONS: FOR PROFESSIONAL USE ONLY

Laboratory studies have indicated that repeated use of Acidulated Phosphate Fluoride products may dull porcelain and composite restorations.

Safety and effectiveness below age 6 have not been established. There have been no long term studies with this product to evaluate carcinogenic, mutagenic or impairment of fertility potential.

Carcinogenesis, Mutagenesis, Impairment of Fertility:

No evidence of carcinogenicity was observed in female and male mice at doses ranging from 2.4 to 18.8 mg/kg sodium fluoride of body weight (3,4). Equivocal evidence of carcinogenicity was reported in male rats at doses ranging from 2.5 to 4.1 mg/kg fluoride, but no evidence of carcinogenicity was observed in female rates (3,4). In another study, no carcinogenicity was observed in rats treated with

fluoride up to 25 mg/kg of body weight (5). Overall, epidemiological studies do not show an association between fluoridated drinking water and increased cancer risks in humans (7).

Fluoride ion is not mutagenic in standard bacterial systems but has been associated with genetic aberrations in cultured human cells at doses much higher than expected for human exposure (6,8). Some in vivo studies report chromosomal aberrations in rodents while other studies using similar protocols report negative results (7).

Potential adverse reproductive effects of fluoride exposure in humans have not been adequately evaluated. Adverse reproductive effects of fluoride have been reported in animal studies, but at high concentrations sufficient to produce other manifestations of toxicity (9).

Pregnancy:Teratogenic Effects: Pregnancy Category B. Fluoride readily crosses the placenta (7,9). Animal studies (rats and rabbits) have shown that fluoride is not a teratogen (10,12,13). Maternal exposure to 18 mg Fluoride/kg of body weight did not affect maternal body weight, litter size or fetal weight and did not increase frequency of skeletal or visceral malformations (10). There are no adequate and well-controlled studies in pregnant women. Several epidemiological studies show no increase in birth defects in areas with fluoridated water compared to areas with low fluoridated water (7). However, caution should be exercised when fluoride is administered to pregnant women.

Nursing Mothers: Due to the relative insensitivity of human milk fluoride levels to changes in maternal fluoride intake, and due to the low concentrations of fluoride in human milk, fluoride supplementation during lactation would not be expected to significantly affect fluoride intake by the nursing infant (11). However, caution should be exercised when fluoride is administered to nursing women.

Pediatric Use: The use of fluoride solutions, gels, and foams containing up to 1.23 % fluoride ion as caries preventives in pediatric patients aged 6 to 16 years is supported by clinical studies in students aged 6 to 12 years (1,2). Safety and effectiveness in pediatric patients below the age of 6 years has not been established. Please refer to CONTRAINDICATIONS and WARNINGS sections.

Geriatric Use: No overall differences in safety or effectiveness have been observed between geriatric and younger patients. This drug is known to be substantially excreted by the kidney, therefore the risk of toxic reactions to this drug may be greater in patients with impaired renal function. Because elderly patients are more likely to have decreased renal function, care should be taken in dose selection, and it may be useful to monitor renal function.

ADVERSE REACTIONS: Developing teeth of children under age 6 may become permanently discolored if excessive amounts are repeatedly swallowed. The following adverse reactions are possible in individuals hypersensitive to fluoride: eczema, atopic dermatitis, urticaria, gastric distress, headache and weakness

OVERDOSAGE: Accidental ingestion of large amounts of fluoride can result in acute irritation of the mouth and gastrointestinal symptoms such as nausea, vomiting, diarrhea, hematemesis epigastric cramping and abdominal pain. If large amounts of fluoride equal to or greater than 5mg fluoride/kg body weight (or 2.3 mg fluoride/lb body weight) are ingested, give calcium (e.g. milk, 5% calcium gluconate or calcium lactate solution) orally to relieve gastrointestinal symptoms and admit immediately to a hospital facility.

A standard treatment dose of Minute-Foam contains approximately 12.4 mg fluoride. One 165 g (5.8oz) bottle contains approximately 1.95 g of fluoride.

DOSAGE AND ADMINISTRATION: Adults and Children 6 years and over: Use foam 2 times a year. This initial time you dispense from a new bottle, gently lift upward on the nozzle to break the protective seal (thin plastic located adjacent to trigger). If this seal is broken do not use the product.

Shake bottle vigorously for 3-4 seconds prior to dispensing.

Invert bottle 180 degree with nozzle tip pointed downward into tray.

Place nozzle tip close to the tray floor and at one end of the tray arch. Moving from one end of the tray to the other in one fluid motion, slowly press down on the trigger.

Dispense foam into applicator tray. (Note: foam will expand slightly to fill the tray.)

Immediately place trays in mouth. (Do not fill trays with foam too far in advance of treatment as the foam will collapse and not be as effective.)

Have patient bite down on the trays lightly but firmly for 4 minutes.

Remove trays and have patient expectorate excess.

Instruct patient not to eat, drink, or rinse for at least 30 minutes.

For optimal tooth coverage, use a fluoride applicator tray which is deep enough to reach the entire vertical height of all teeth (even the molars)

HOW SUPPLIED: Minute-Foam is available in a 165 g (5.8 oz) plastic aerosol container in six different flavors; NDC 0041-0340-06 Banana Splitz, NDC 0041-03141-06 Mellow Mint, NDC 0041-0342-06 Orang-a-Tangy, NDC 0041-0345-06 Bubblegum, NDC 0041-0346-06 Strawberry, NDC 0041-0347-06 Grape Punch.

REFERENCES:

(1.) Wellock, W.D. and Brudevold, F.: Arch. Oral Biol., 10, 453-460 (1965) (2.) Jiang, H et al.:J. Dent, 33, 469-473 (2005) (3.) National Toxicology Program: NTP TR 393, NIH Publication 91-2842, (1990) (4.) Bucker, J.R et al.L Int. J. Cancer 48, 1118-1126 (1990) (6.) Martin, G.R. et al.: Mutat. Res. 66, 159-167 (1979) (7.) Agency for toxic Substances and Disease Registry: Toxicological Profile for Fluorides (2003) (8.) Aardema, M.J. et al.: Mutat. Res. 331 (1), 171-172 (9.) National Research Council: Fluoride in Drinking Water (2006) (10.) Heindel, J.J. et al.: Fundam Appl Toxicol, 30, 162-177 (1996) (11.) Institute of Medicine, Food and Nutrition Board: Dietary Reference Intakes (1997) (12.) Collins, T.F et al.: Food Chem. Toxicol. 33 (11), 951-960 (13.) Collins, T.F. et al.: Food Chem Toxicol. 39 (8), 867-876

Minute Foam Label

Oral-B Minute-Foam® est une solution moussante aqueuse aromatisée de fluorure de phosphate acidulé ayant un pH de 3,5 et une concentration en ions de fluorure de 1,23 % p/p.

INGRÉDIENT ACTIF/MÉDICAMENTEUX : Fluorure de phosphate acidulé (fluorure de sodium et acide fluorhydrique) à 1,23 % p/p d'ions de fluorure.

INGRÉDIENTS INACTIFS/NON-MÉDICAMENTEUX : eau purifiée, poloxamère 407, isobutane, poloxamère 234, acide phosphorique, saccharine sodique, arôme de raisins.

INDICATIONS ET MODE D'EMPLOI : Appliquer la solution moussante pour aider à prévenir la carie dentaire.

AVERTISSEMENTS : NE PAS AVALER. Tenir hors de la portée des enfants. En cas d'ingestion d'une quantité supérieure à la quantité recommandée, donner du calcium (du lait, par exemple) et consulter sans délai un médecin ou un centre antipoison. Éviter de vaporiser le produit en direction d'une flamme nue. Contenu sous pression. Le contenant ne doit pas être percé ni jeté au feu. Le contenant ne doit pas être exposé à la chaleur ni entreposé à une température excédant 49 °C (120 °F).

POSOLOGIE ET ADMINISTRATION : Le temps d'application est de 4 minutes. POUR PLUS DE DÉTAILS, CONSULTER LA NOTICE D'ACCOMPAGNEMENT DU PRODUIT.

U.S.A./É.U. : Manufactured for/Fabriqué pour Oral-B Laboratories, a Division of/une division de Procter & Gamble, Cincinnati OH 45202

Canada : Distr. by/par : Oral-B Laboratories, a Division of/une division de Procter & Gamble Inc., Toronto, ON M5W 1C5

Questions?
1-800-566-7252

www.dentalcare.com

NDC 0041-0347-06

NPN 02048787

Oral-B® Minute-Foam®

**Topical Fluoride
Foaming Solution*
Solution moussante
topique au fluorure ***

*Acidulated Phosphate Fluoride
Containing 1.23% w/w Fluoride Ion
*Fluorure de phosphate acidulé
Contient 1,23 % p/p d'ions de fluorure

Rx only (in U.S.A.)

Sur ordonnance seulement (aux É.-U.)

For professional use only.
Shake vigorously before each use.
Protect from freezing.

Pour usage professionnel seulement.
Secouer la bouteille vigoureusement
avant d'administrer le produit.
Garder à l'abri du gel.

**Grape Punch
Punch aux raisins**

165 g (5.8 oz)

DESCRIPTION: Oral-B Minute-Foam® is a flavored aqueous foaming solution of acidulated phosphate sodium fluoride with a pH of 3.5 and a fluoride ion concentration of 1.23% w/w.

ACTIVE/MEDICINAL INGREDIENT: Acidulated Phosphate Fluoride (Sodium Fluoride and Hydrofluoric Acid) 1.23% w/w fluoride ion.

INACTIVE/NON-MEDICINAL INGREDIENTS: Purified water, poloxamer 407, isobutane, poloxamer 234, phosphoric acid, sodium saccharin, grape flavor.

INDICATIONS AND USE: A topically-applied foaming solution to aid in the prevention of dental caries.

WARNINGS: DO NOT SWALLOW. Keep out of reach of children. In the event more than the treatment dose is swallowed, administer calcium (e.g. milk) and get medical help or contact a Poison Control Center right away. Avoid spraying towards open flame. Contents under pressure. Do not puncture or incinerate. Do not expose to heat or store at temperatures over 120°F (49°C).

DOSAGE AND ADMINISTRATION: Requires a 4 minute application time. SEE PRODUCT INFORMATION LEAFLET FOR COMPLETE DETAILS.

98989321

ORAL-B MINUTE-FOAM GRAPE PUNCH

acidulated phosphate fluoride aerosol

Product Information

Product Type	HUMAN PRESCRIPTION DRUG	Item Code (Source)	NDC:0041-0347
Route of Administration	DENTAL		

Active Ingredient/Active Moiety

Ingredient Name	Basis of Strength	Strength
Sodium Fluoride (UNII: 8ZYQ1474W7) (Fluoride ion - UNII:Q80VPU408O)	Fluoride ion	10 mg in 1 g

Product Characteristics

Color		Score	
Shape		Size	
Flavor	GRAPE (Grape Flavor)	Imprint Code	
Contains			

Packaging

#	Item Code	Package Description	Marketing Start Date	Marketing End Date
1	NDC:0041-0347-06	165 g in 1 BOTTLE		

Marketing Information

Marketing Category	Application Number or Monograph Citation	Marketing Start Date	Marketing End Date
unapproved drug other		05/04/2000	

Labeler - Oral-B Laboratories (183102243)

Revised: 8/2010

Oral-B Laboratories