TUSSLIN PEDIATRIC- dextromethorphan hbr, guiafenes in, phenylephrine hcl solution/ drops Kramer Novis

Disclaimer: Most OTC drugs are not reviewed and approved by FDA, however they may be marketed if they comply with applicable regulations and policies. FDA has not evaluated whether this product complies.

Drug Facts

Active Ingredients (in each 1 mL)	Purpose
Dextromethorphan HBr, 7.5 mg	Antitussive
Guaifenesin, 88 mg	Expectorant
Phenylephrine HCl, 2.5 mg	Nasal decongestant

Uses

- helps loosen phlegm (mucus) and thin bronchial secretions to drain bronchial tubes and make coughs more productive
- temporarily relieves:
- nasal congestion due to the common cold
- cough due to minor throat and bronchial irritation as may occur with the common cold

Warnings

Do not use

in child who is taking a prescription monoamine oxidase inhibitor (MAOI) (certain drugs for depression, psychiatric, or emotional conditions, or Parkinson's disease), or for 2 weeks after stopping the MAOI drug. If you do not know if your child's prescription drug contains a MAOI, ask a doctor or pharmacist before giving this product.

Ask a doctor before use if your child has

- heart disease
- thyroid disease
- high blood pressure
- diabetes
- a cough that occurs with too much phlegm (mucus)
- cough that lasts or is chronic such as occurs with asthma

When using this product do not exceed recommended dosage

Stop use and ask a doctor if

- new symptoms occur
- your child gets nervous, dizzy or sleepless
- symptoms do not get better within 7 days or are accompanied by fever
- cough last more than 7 days, comes back, or is accompanied by fever, rash or persistent headache. These could be a signs of a serious condition.

If pregnant or breast-feeding, ask a health professional before use.

Keep out of reach of children. In case of accidental overdose, get medical help or contact a Poison Control Center right away.

Directions

do not use more than 4 doses in any 24-hour period

Children 6 to under 12 years of age

• 2 mL every 6 hours, do not exceed 4 doses.

Children 2 to under 6 years of age

• 1 mL every 6 hours, do not exceed 4 doses, or as directed by a doctor.

Children under 2, consult a doctor

• measure with the dosage device provided. Do not use any other device.

Other information

Tamper Evident Feature: Do not use if cap seal is torn, broken or missing.

Store at controlled room temperature 15-30°C (59-86°F).

Avoid excessive heat and humidity.

Inactive ingredients Citric acid, flavors, menthol, methyl paraben, monoammonium glycyrrhizinate, polyethylene glycol, propylene glycol, propyl paraben, purified water, sorbitol and sucralose.

Manufactured in the USA for Kramer-Novis, San Juan, PR 00917.

Tel: (787) 767-2072 www.kramernovis.com

Principal Display Panel

TUSSLIN®

PEDIATRIC DROPS

Contains the same active ingredients as Glituss® Pediatric

Antitus sive

Expectorant

Nasal Decongestant

Sugar, Alcohol & Dye Free

NET CONTENT: 30 mL (1 Fl. Oz.)

KRAMER-NOVIS

Antitussive • Expectorant • Masal Decongestant

TUSSLIN®

NDC 95083-953-01

Drug Facts

Active Ingredients (in each 1 mL) Purpose
Dextromethorphan HBr,7.5 mg........Antitussive
Guaifenesin, 88 mg.......Expectorant
Phenylephrine HCl, 2.5 mg.....Nasal Decongestant

Uses

- helps loosen phlegm (mucus) and thin bronchial secretions to drain bronchial tubes and make coughs more productive
- temporarily relieves:
 - nasal congestion due to hay fever or other upper respiratory allergies (allergic rhinitis)
 - cough due to minor throat and bronchial imitation as may occur with the common cold

Warnings

Do not use

in a child who is taking a prescription monoamine oxidase inhibitor (MAOI) (certain drugs for depression, psychiatric, or emotional conditions, or Parkinson's disease), or for 2 weeks after stopping the MAOI drug. If you do not know if your child's prescription drug contains an MAOI, ask a doctor or pharmacist before giving this product.

Ask a doctor before use if the child has

- · heart disease · thyroid disease · high blood pressure
- diabetes
- cough that occurs with too much phiegm (mucus)
 cough that lasts or is chronic such as occurs with asthma.

When using this product do not exceed recommended dosage.

Stop use and ask a doctor if

new symptoms occur · your child gets nervous, fizzy, or sleepless

RETAIN CARTON FOR FULL PRESCRIBING INFORMATION

Drug Facts (continued)

Stop use and ask a doctor if • symptoms do not get better within 7 days or are accompanied by fever

- coughs lasts for more than 7 days, comes back, or is accompanied by fever, rash, or persistent headache. These could be signs of a serious condition.
- If pregnant or breastfeeding, ask a health professional before use. Keep out of reach of children. In case of accidental overdose, get medical help or contact a Poison Control Center right away.

Directions

do not take more than 4 doses in any 24-hour period Children 6 to under 12 years of age • 2 mL every 6 hours, do not

exceed 4 doses. Children 2 to under 6 years

Children 2 to under 6 years of age

 1 mL every 6 hours, do not exceed 4 doses, or as directed by a doctor.

Children under 2, consult a

Children under 2, consult doctor.

 measure with the dosage device provided. Do not use any other device. NDC 52083-623-01

TUSSLIN

PEDIATRIC DROPS

Contains the same active ingredients as Giltuss® Pediatric*

Antitussive Expectorant Nasal Decongestant

Sugar, Alcohol & Dye Free

NET CONTENT: 30 mL (1 FL Oz.)



Drug Facts (continued)

Other information

Tamper Evident Feature;
 Do not use if cap seal is torn, broken or missing.
 Store at controlled room temperature 15°-30°C (59°-86°F).

 Avoid excessive heat and humidity.

Inactive ingredients

Citric acid, flavors, menthol, methyl paraben, monoammonium glycormizinate, pollyethylene glycol, propylene glycol, propyl paraben, purified water, sorbitol and sucralose.

Manufactured in the USA for Kramer-Novis, San Juan, PR 00917. Tel: (787) 767-2072.

Tel: (787) 767-2072. www.kramernovis.com

* Giltuss® Pediatric is a registered trademark of Gil Pharmaceutical Corp. This product is not manufactured, distributed or marketed by Gil Pharmaceutical Corp.



Lot No.

Exp. Date

Non Coating Area

RETAIN CARTON FOR FULL PRESCRIBING INFORMATION

TUSSLIN PEDIATRIC

dextromethorphan hbr, guiafenesin, phenylephrine hcl solution/ drops

Product Information

Product Type HUMAN OTC DRUG Item Code (Source) NDC:52083-623

Route of Administration (

ORAL

Active Ingredient/Active Moiety				
Ingredient Name	Basis of Strength	Strength		
DEXTROMETHORPHAN HYDROBROMIDE (UNII: 9 D2RTI9 KYH) (DEXTROMETHORPHAN - UNII:7355X3ROTS)	DEXTROMETHORPHAN HYDROBROMIDE	7.5 mg in 1 mL		
GUAIFENESIN (UNII: 495W7451VQ) (GUAIFENESIN - UNII:495W7451VQ)	GUAIFENESIN	88 mg in 1 mL		
PHENYLEPHRINE HYDRO CHLO RIDE (UNII: 04JA59TNSJ) (PHENYLEPHRINE - UNII:1WS297W6MV)	PHENYLEPHRINE HYDROCHLORIDE	2.5 mg in 1 mL		

Inactive Ingredients		
Ingredient Name	Strength	
CITRIC ACID MONO HYDRATE (UNII: 2968 PHW8 QP)		
METHYLPARABEN (UNII: A2I8C7HI9T)		
AMMO NIUM GLYCYRRHIZATE (UNII: 3VRD35U26C)		
POLYETHYLENE GLYCOL, UNSPECIFIED (UNII: 3WJQ0SDW1A)		
PROPYLENE GLYCOL (UNII: 6DC9Q167V3)		
PROPYLPARABEN (UNII: Z8IX2SC1OH)		
WATER (UNII: 059QF0KO0R)		
SORBITOL (UNII: 506T60A25R)		
SUCRALOSE (UNII: 96K6UQ3ZD4)		

Packaging				
#	Item Code	Package Description	Marketing Start Date	Marketing End Date
1	NDC:52083-623- 01	30 mL in 1 BOTTLE, DROPPER; Type 0: Not a Combination Product	0 1/27/20 14	

Marketing Information					
Marketing Category	Application Number or Monograph Citation	Marketing Start Date	Marketing End Date		
OTC monograph final	part341	0 1/27/20 14			

Labeler - Kramer Novis (090158395)

Revised: 9/2019 Kramer Novis