

CIMETIDINE ACID REDUCER- cimetidine tablet, film coated
Walgreen Company

Walgreen Co. Cimetidine 200 Drug Facts

Active ingredient (in each tablet)

Cimetidine 200 mg

Purpose

Acid reducer

Uses

- **relieves** heartburn associated with acid indigestion and sour stomach
- **prevents** heartburn associated with acid indigestion and sour stomach brought on by eating or drinking certain foods and beverages

Warnings

Allergy alert: Do not use if you are allergic to cimetidine or other acid reducers

Do not use

- if you have trouble or pain swallowing food, vomiting with blood, or bloody or black stools. These may be signs of a serious condition. See your doctor.
- with other acid reducers

Ask a doctor before use if you have

- frequent **chest pain**
- frequent wheezing, particularly with heartburn
- unexplained weight loss
- nausea or vomiting
- stomach pain
- had heartburn over 3 months. This may be a sign of a more serious condition.
- heartburn with **lightheadedness, sweating or dizziness**
- chest pain or shoulder pain with shortness of breath; sweating; pain spreading to arms, neck or shoulders; or lightheadedness
- kidney disease
- liver disease

Ask a doctor or pharmacist before use if you are taking

- a prescription drug. Acid reducers may interact with certain prescription drugs.

Stop use and ask a doctor if

- you need to take this product for more than 14 days
- stomach pain continues
- your heartburn continues or worsens

If pregnant or breast-feeding,

ask a health professional before use.

Keep out of reach of children.

In case of overdose, get medical help or contact a Poison Control Center right away (1-800-222-1222).

Directions

- adults and children 12 years and over:
- to **relieve** symptoms, swallow 1 tablet with a glass of water
- to **prevent** symptoms, swallow 1 tablet with a glass of water **right before or any time up to 30 minutes before** eating food or drinking beverages that cause heartburn
- do not take more than 2 tablets in 24 hours
- children under 12 years: ask a doctor

Other information

- store at 20°-25°C (68°-77°F)

Inactive ingredients

hypromellose, magnesium stearate, microcrystalline cellulose, polyethylene glycol, polysorbate 80, povidone, pregelatinized starch, sodium lauryl sulfate, sodium starch glycolate, titanium dioxide

Questions or comments?

1-800-719-9260

Principal Display Panel

Compare to Tagamet HB® 200 active ingredient

Cimetidine 200

CIMETIDINE TABLETS USP, 200 mg / ACID REDUCER

Just one tablet relieves & prevents heartburn & acid indigestion

Take before, during or after a meal

60 TABLETS (60 DOSES)

ACTUAL SIZE



02272 94 05



Gluten Free
 Walgreens Pharmacist Survey
 If this product is not manufactured or
 distributed by Medtech Products Inc.,
 distributor of Tagamet® HB200.

Drug Facts**Active ingredient (in each tablet)**

Cimetidine 200 mg.....

Purpose

Acid reducer

Uses ■ relieves heartburn associated with acid indigestion and sour stomach
 ■ prevents heartburn associated with acid indigestion and sour stomach brought on by eating or drinking certain foods and beverages

Warnings

Allergy alert: Do not use if you are allergic to cimetidine or other acid reducers

Do not use ■ if you have trouble or pain swallowing food, vomiting with blood, or bloody or black stools. These may be signs of a serious condition. See your doctor.
 ■ with other acid reducers

Ask a doctor before use if you have ■ frequent chest pain ■ frequent wheezing, particularly with heartburn
 ■ unexplained weight loss ■ nausea or vomiting ■ stomach pain
 ■ had heartburn over 3 months. This may be a sign of a more serious condition.
 ■ heartburn with lightheadedness, sweating or dizziness
 ■ chest pain or shoulder pain with shortness of breath; sweating; pain spreading to arms, neck or shoulders; or lightheadedness
 ■ kidney disease ■ liver disease

Ask a doctor or pharmacist before use if you are taking ■ a prescription drug. Acid reducers may interact with certain prescription drugs.

Stop use and ask a doctor if ■ you need to take this product for more than 14 days
 ■ stomach pain continues ■ your heartburn continues or worsens

If pregnant or breast-feeding, ask a health professional before use.

Keep out of reach of children. In case of overdose, get medical help or contact a Poison Control Center right away (1-800-222-1222).

Directions

■ adults and children 12 years and over:
 ■ to relieve symptoms, swallow 1 tablet with a glass of water
 ■ to prevent symptoms, swallow 1 tablet with a glass of water **right before or anytime up to 30 minutes before** eating food or drinking beverages that cause heartburn
 ■ do not take more than 2 tablets in 24 hours
 ■ children under 12 years: ask a doctor

PHARMACIST

Walgreens

RECOMMENDED

Health expert is
 you rely on,
 quality you trust.

DISTRIBUTED BY: WALGREEN CO.
 200 WILMOT RD., DEERFIELD, IL 60015
 Walgreens
**100% SATISFACTION
 GUARANTEED**
 walgreens.com ©2017 Walgreen Co.

**Drug Facts (continued)****Other information**


■ store at 20°-25°C (68°-77°F)

Inactive ingredients hypromellose, magnesium stearate, microcrystalline cellulose, polyethylene glycol, polyorbate 80, povidone, pregelatinized starch, sodium lauryl sulfate, sodium starch glycolate, titanium dioxide

Questions or comments? 1-800-719-9260

Tips for Managing Heartburn

- Do not lie flat or bend over soon after eating.
- Do not eat late at night, or just before bedtime.
- Certain foods or drinks are more likely to cause heartburn, such as rich, spicy, fatty, and fried foods, chocolate, caffeine, alcohol, even some fruits and vegetables.
- Eat slowly and do not eat big meals.
- If you are overweight, lose weight.
- If you smoke, quit smoking.
- Raise the head of your bed.
- Wear loose fitting clothing around your stomach.

**CIMETIDINE ACID REDUCER**

cimetidine tablet, film coated

Product Information

Product Type	HUMAN OTC DRUG	Item Code (Source)	NDC:0363-0022
Route of Administration	ORAL		

Active Ingredient/Active Moiety

Ingredient Name	Basis of Strength	Strength
CIMETIDINE (UNII: 80061L1WGD) (CIMETIDINE - UNII:80061L1WGD)	CIMETIDINE	200 mg

Inactive Ingredients

Ingredient Name	Strength
HYPROMELLOSE, UNSPECIFIED (UNII: 3NXW29V3WO)	
MAGNESIUM STEARATE (UNII: 70097M6130)	
MICROCRYSTALLINE CELLULOSE (UNII: OP1R32D61U)	
POLYETHYLENE GLYCOL, UNSPECIFIED (UNII: 3WJQ0SDW1A)	

POLYSORBATE 80 (UNII: 6OZP39ZG8H)	
POVIDONE, UNSPECIFIED (UNII: FZ989GH94E)	
SODIUM LAURYL SULFATE (UNII: 368GB5141J)	
TITANIUM DIOXIDE (UNII: 15FIX9V2JP)	

Product Characteristics

Color	WHITE	Score	no score
Shape	OVAL	Size	13mm
Flavor		Imprint Code	L022
Contains			

Packaging

#	Item Code	Package Description	Marketing Start Date	Marketing End Date
1	NDC:0363-0022-72	1 in 1 CARTON	04/14/2010	
1		60 in 1 BOTTLE; Type 0: Not a Combination Product		

Marketing Information

Marketing Category	Application Number or Monograph Citation	Marketing Start Date	Marketing End Date
ANDA	ANDA075285	03/27/2001	

Labeler - Walgreen Company (008965063)

Revised: 5/2021

Walgreen Company