SOMXL- salicylic acid solution WR Group Skin Care Ltd

SOMXL

Drug Facts

Active ingredient

Salicylic acid 17% (w/w)

Purpose

Wart remover

Use

For the removal of common warts. The common wart is easily recognized by the rough "cauliflower-like" appearace on the surface.

Warnings

For external use only

If product gets into eyes, flush with water for 15-minutes. Avoid inhaling vapors.

Do not use

- on irritated skin
- on any area that is infected or reddened
- if you are diabetic or if you have poor blood circulation
- piles
- birthmarks
- warts with hair growing from them
- on moles
- warts on the face or mucous membreanes.

Stop use and ask a doctor if

discomfort persists

If pregnant or breast-feeding,

ask a health professional before use.

Keep out of reach of children.

If swallowed, get medical help or contact a Poison Control Center right away.

Directions

- Wash affected area (May soak wart in warm water for 15-minutes if desired).
- Dry area throughly.
- Apply small amount with applicator to cover each wart. Let dry.
- Repeat this procedure once of twice daily as needed for up to 12-weeks.

Inactive ingredients

castor oil, ethyl lactate, flexible collodion, polybutene, (Contains alcohol 18% v/v and ether 53% v/v from flexible collodion.)

Package Labeling:

_®TXWOS

Drug Facts

Active ingredient Salicylic acid 17% (w/w) ...

Use For the removal of common warts. The common wart is easily recognized by the rough "cauliflower-like" appearance on the surface.

Warnings

For external use only If product gets into eyes, flush with water for 15-minutes. Avoid inhaling vapors.

- Do not use on irritated skin on any area that is infected or reddened
- if you are diabetic or if you have poor blood
- circulation piles birthmarks
 warts with hair growing from them on moles
- warts on the face or mucous membranes

Stop use and ask a doctor if discomfort persists

If pregnant or breast-feeding, ask a health professional before use.

Keep out of reach of children. If swallowed, get medical help or contact a Poison Control Center right away.

Directions

- Wash affected area (May soak wart in warm water for 15-minutes if desired).
- Dry area throughly.
- Apply small amount with applicator to cover each wart. Let dry.
- Repeat this procedure once or twice daily as needed for up to 12-weeks.

Inactive ingredients

castor oil, ethyl lactate, flexible collodion, polybutene. (Contains alcohol 18% v/v and ether 53% v/v from flexible collodion.)

DISTRIBUTED BY WR GROUP SKIN CARE LTD PO BOX 13867, CHRISTCHURCH, NEW ZEALAND



Wart Remove







Net wt. 0.5 oz (15 g)

SOMXL

salicylic acid solution

Product Information

HUMAN OTC DRUG **Product Type** Item Code (Source) NDC:70463-000

Route of Administration TOPICAL

Active Ingredient/Active Moiety					
Ingredient Name	Basis of Strength	Strength			
SALICYLIC ACID (UNII: 0414P74LP7) (SALICYLIC ACID - UNII:0414P74LP7)	SALICYLIC ACID	170 mg in 1 g			

Inactive Ingredients				
Ingredient Name	Strength			
CASTOR OIL (UNII: D5340Y2I9G)				
ETHYL LACTATE (UNII: F3P750VW8I)				

P	Packaging					
#	Item Code	Package Description	Marketing Start Date	Marketing End Date		
1	NDC:70463-000- 01	1 in 1 BOX	06/26/2023			
1		15 g in 1 TUBE; Type 0: Not a Combination Product				

Marketing Information			
Marketing Category	Application Number or Monograph Citation	Marketing Start Date	Marketing End Date
OTC Monograph Drug	M028	06/26/2023	

Labeler - WR Group Skin Care Ltd (592719822)

Revised: 11/2023 WR Group Skin Care Ltd