

ACETAMINOPHEN- acetaminophen tablet, film coated, extended release
Amerisource Bergen

Drug Facts

ACTIVE INGREDIENT (IN EACH CAPLET)

Acetaminophen USP, 650 mg

PURPOSE

Pain reliever/fever reducer

USES

- temporarily relieves minor aches and pains due to:
 - minor pain of arthritis
 - muscular aches
 - backache
 - headache
 - toothache
 - the common cold
 - premenstrual and menstrual cramps

- temporarily reduces fever

WARNINGS

Liver warning: This product contains acetaminophen. Severe liver damage may occur if you take

- more than 6 caplets in 24 hours, which is the maximum daily amount
- with other drugs containing acetaminophen
- 3 or more alcoholic drinks every day while using this product

Do not use

- with any other drug containing acetaminophen (prescription or nonprescription). If you are not sure whether a drug contains acetaminophen, ask a doctor or pharmacist.
- if you are allergic to acetaminophen or any of the inactive ingredients in this product.

Ask a doctor before use if you have

Liver disease.

Ask a doctor or pharmacist before use if you are

Taking the blood thinning drug warfarin.

Stop use and ask a doctor if

- pain gets worse or lasts more than 10 days
- fever gets worse or lasts more than 3 days
- new symptoms occur
- redness or swelling is present

These could be signs of a serious condition.

If pregnant or breast-feeding

Ask a health professional before use.

Keep out of reach of children.

Overdose warning: In case of overdose, get medical help or contact a Poison Control Center right away (1-800-222-1222). Quick medical attention is critical for adults as well as for children even if you do not notice any signs or symptoms.

DIRECTIONS

- **do not take more than directed (see overdose warning)**

adults	<ul style="list-style-type: none">▪ take 2 caplets every 8 hours with water▪ swallow whole - do not crush, chew, split or dissolve▪ do not take more than 6 caplets in 24 hours▪ do not use for more than 10 days unless directed by a doctor
under 18 years of age	<ul style="list-style-type: none">▪ ask a doctor

OTHER INFORMATION

- store at 20 - 25° C (68 - 77° F). Avoid excessive heat 40° C (104° F).
- **TAMPER EVIDENT: DO NOT USE IF IMPRINTED SEAL IS BROKEN OR MISSING FROM BOTTLE.**

INACTIVE INGREDIENTS

Croscarmellose sodium, hypromellose, magnesium stearate, microcrystalline cellulose, povidone, pregelatinized starch, propylene glycol, sodium lauryl sulfate, stearic acid, titanium dioxide

QUESTIONS?

Call **1-800-406-7984**

PRINCIPAL DISPLAY PANEL - 650 mg Caplet Bottle Label

†Compare to the active ingredient
in Tylenol® Arthritis Pain

GOOD
NEIGHBOR
PHARMACY®

NDC 46122-170-81

Last up to 8 HOURS

Use only as directed.

Arthritis Pain Relief
acetaminophen
extended-release tablets
USP, 650 mg

Pain Reliever/Fever Reducer

For the Temporary Relief of Minor Arthritis Pain

DO NOT USE WITH OTHER MEDICINES
CONTAINING ACETAMINOPHEN

200 caplets*

(*capsule-shaped tablets)



GOOD NEIGHBOR PHARMACY

*Compare to the active ingredient in Tylenol® Arthritis Pain
NDC 46122-170-81

Last up to **8 HOURS**

Use only as directed.

Arthritis Pain Relief

acetaminophen

extended-release tablets
USP, 650 mg

Pain Reliever/Fever Reducer

For the Temporary Relief of Minor Arthritis Pain

DO NOT USE WITH OTHER MEDICINES CONTAINING ACETAMINOPHEN

200 caplets*
(*capsule-shaped tablets)



Drug Facts

Active ingredient (in each caplet)
Acetaminophen USP, 650 mg. Pain reliever/fever reducer.

Uses ■ temporarily relieves minor aches and pains due to:
■ minor pain of arthritis ■ muscular aches ■ backache ■ premenstrual and menstrual cramps ■ the common cold ■ headache ■ toothache ■ temporarily reduces fever.

Warnings
Liver warning: This product contains acetaminophen. Severe liver damage may occur if you take ■ more than 6 caplets in 24 hours, which is the maximum daily amount ■ with other drugs containing acetaminophen ■ 3 or more alcoholic drinks every day while using this product.
Alert: Acetaminophen may cause severe skin reactions. Symptoms may include:
■ skin redness ■ blisters ■ rash ■ if a skin reaction occurs, stop use and see medical help right away.
Do not use ■ with any other drug containing acetaminophen (prescription or nonprescription). If you are not sure whether a drug contains acetaminophen, ask a doctor or pharmacist. ■ if you are allergic to acetaminophen or any of the inactive ingredients in this product.
Ask a doctor before use if you have liver disease.

Stop use and ask a doctor if ■ pain gets worse or lasts more than 10 days ■ fever gets worse or lasts more than 3 days ■ new symptoms occur ■ redness or swelling is present. These could be signs of a serious condition.

Keep out of reach of children.
Overdose warning: In case of overdose, get medical help or contact a Poison Control Center right away (1-800-222-1222). Quick medical attention is critical for adults as well as for children even if you do not notice any signs or symptoms.

Directions ■ do not take more than directed (see overdose warning)
adults ■ take 2 caplets every 8 hours with water ■ swallow whole; do not crush, chew, split or dissolve ■ do not take more than 6 caplets in 24 hours ■ do not use for more than 10 days unless directed by a doctor
under 18 years of age ■ ask a doctor

Other information ■ store at 20°-25° C (68°-77°F). Avoid excessive heat 40° C (104°F). ■ TAMPER EVIDENT: DO NOT USE IF IMPRINTED SEAL IS BROKEN OR MISSING FROM BOTTLE.

Inactive ingredients croscarmellose sodium, hypromellose, magnesium stearate, microcrystalline cellulose, povidone, prepolimerized starch, propylene glycol, sodium lauryl sulfate, stearic acid, titanium dioxide

Questions? call 1-400-406-7984
† All trademarks are property of their respective owners. This product is not affiliated with the manufacturer of Tylenol® Arthritis Pain.
Distributed by: Amneal Pharmaceuticals, Inc., 7300 Morris Drive, Basking Ridge, NJ 07005
Visit www.MylgHP.com for more information, questions or concerns?
R0319 Batch No. Expiration Date:



5196478



ABC# 10101657



0 877014 114916

non varnish area

ACETAMINOPHEN

acetaminophen tablet, film coated, extended release

Product Information

Product Type	HUMAN OTC DRUG	Item Code (Source)	NDC:46 122-170
Route of Administration	ORAL		

Active Ingredient/Active Moiety

Ingredient Name	Basis of Strength	Strength
ACETAMINOPHEN (UNII: 362O9ITL9D) (ACETAMINOPHEN - UNII:362O9ITL9D)	ACETAMINOPHEN	650 mg

Inactive Ingredients

Ingredient Name	Strength
CROSCARMELLOSE SODIUM (UNII: M28OL1HH48)	
HYPROMELLOSE, UNSPECIFIED (UNII: 3NXW29V3WO)	

MAGNESIUM STEARATE (UNII: 70097M6I30)	
MICROCRYSTALLINE CELLULOSE (UNII: OP1R32D61U)	
POVIDONE, UNSPECIFIED (UNII: FZ989GH94E)	
STARCH, CORN (UNII: O8232NY3SJ)	
PROPYLENE GLYCOL (UNII: 6DC9Q167V3)	
SODIUM LAURYL SULFATE (UNII: 368GB5141J)	
STEARIC ACID (UNII: 4ELV7Z65AP)	
TITANIUM DIOXIDE (UNII: 15FIX9V2JP)	

Product Characteristics

Color	white	Score	no score
Shape	OVAL (Capsule Shaped)	Size	19mm
Flavor		Imprint Code	cor116
Contains			

Packaging

#	Item Code	Package Description	Marketing Start Date	Marketing End Date
1	NDC:46122-170-81	200 in 1 BOTTLE; Type 0: Not a Combination Product	04/30/2002	

Marketing Information

Marketing Category	Application Number or Monograph Citation	Marketing Start Date	Marketing End Date
ANDA	ANDA076200	04/30/2002	

Labeler - Amerisource Bergen (007914906)

Registrant - Ranbaxy Pharmaceuticals Inc. (937890044)

Establishment

Name	Address	ID/FEI	Business Operations
Ohm Laboratories Inc.		184769029	manufacture(46122-170)

Revised: 4/2019

Amerisource Bergen