

PAIN RELIEVER EXTRA STRENGTH- acetaminophen tablet, coated
L.N.K. International, Inc.

Quality Plus 44-519

Active ingredient (in each gelcap)

Acetaminophen 500 mg

Purpose

Pain reliever/fever reducer

Uses

- temporarily relieves minor aches and pains due to:
 - headache
 - the common cold
 - backache
 - minor pain of arthritis
 - toothache
 - muscular aches
 - premenstrual and menstrual cramps
- temporarily reduces fever

Warnings

Liver warning: This product contains acetaminophen. Severe liver damage may occur if you take

- more than 4,000 mg of acetaminophen in 24 hours
- with other drugs containing acetaminophen
- 3 or more alcoholic drinks every day while using this product

Allergy alert: Acetaminophen may cause severe skin reactions. Symptoms may include:

- skin reddening
- blisters
- rash

If a skin reaction occurs, stop use and seek medical help right away.

Do not use

- if you are allergic to acetaminophen or any of the inactive ingredients in this product
- with any other drug containing acetaminophen (prescription or nonprescription). If you are not sure whether a drug contains acetaminophen, ask a doctor or pharmacist.

Ask a doctor before use if you have

liver disease.

Ask a doctor or pharmacist before use if you are

taking the blood thinning drug warfarin.

Stop use and ask a doctor if

- pain gets worse or lasts more than 10 days
- fever gets worse or lasts more than 3 days
- new symptoms occur
- redness or swelling is present

These could be signs of a serious condition.

If pregnant or breast-feeding,

ask a health professional before use.

Keep out of reach of children.

In case of overdose, get medical help or contact a Poison Control Center right away. Prompt medical attention is critical for adults as well as for children even if you do not notice any signs or symptoms.

Directions

- **do not take more than directed**
- adults and children 12 years and over
 - take 2 gelcaps every 6 hours while symptoms last
 - do not take more than 6 gelcaps in 24 hours, unless directed by a doctor
 - do not take for more than 10 days unless directed by a doctor
- children under 12 years: ask a doctor

Other information

- store at 25°C (77°F); excursions permitted between 15°C-30°C (59°F-86°F)
- avoid high humidity
- see end flap for expiration date and lot number

Inactive ingredients

croscarmellose sodium, D&C red #33, FD&C blue #1, FD&C red #40, gelatin, hydroxypropyl cellulose, hypromellose, iron oxide black, iron oxide red, iron oxide yellow, polyethylene glycol, povidone, pregelatinized starch, propylene glycol, shellac glaze, stearic acid, titanium dioxide

Questions or comments?

1-800-426-9391

Principal Display Panel

QUALITY

+PLUS

NDC 50844-519-21

*Compare to active ingredient
in Extra Strength Tylenol®
Rapid Release Gels

EXTRA STRENGTH

PAIN RELIEVER

Acetaminophen 500 mg

PAIN RELIEVER/FEVER REDUCER

16 Gelcaps

CONTAINS NO ASPIRIN

ACTUAL

SIZE

**TAMPER EVIDENT: DO NOT USE IF IMPRINTED
SAFETY SEAL UNDER CAP IS BROKEN OR MISSING**

*This product is not manufactured or distributed
by Johnson & Johnson Corporation, owner of the
registered trademark Extra Strength Tylenol®
Rapid Release Gels. 50844 ORG032251921

Distributed by **LNK INTERNATIONAL, INC.**

60 Arkay Drive

Hauppauge, NY 11788

USA

QUALITY PLUS

EXTRA STRENGTH

PAIN RELIEVER

Acetaminophen 500 mg

PAIN RELIEVER/FEVER REDUCER

16 Gelcaps

QUALITY PLUS

EXTRA STRENGTH

PAIN RELIEVER

Acetaminophen 500 mg

PAIN RELIEVER/FEVER REDUCER

16 Gelcaps

CONTAINS NO ASPIRIN

ACTUAL SIZE



NDC 50844-519-21

*Compare to active ingredient in Extra Strength Tylenol® Rapid Release Gels

TAMPER EVIDENT: DO NOT USE IF IMPRINTED SAFETY SEAL UNDER CAP IS BROKEN OR MISSING



0 50844 51921 3

No print/No varnish Lot & Exp date

Drug Facts (continued)

Do not use

- if you are allergic to acetaminophen or any of the inactive ingredients in this product
- with any other drug containing acetaminophen (prescription or nonprescription). If you are not sure whether a drug contains acetaminophen, ask a doctor or pharmacist.

Ask a doctor before use if you have liver disease.

Ask a doctor or pharmacist before use if you are taking the blood thinning drug warfarin.

Stop use and ask a doctor if

- pain gets worse or lasts more than 10 days
- fever gets worse or lasts more than 3 days
- new symptoms occur
- redness or swelling is present

These could be signs of a serious condition.

If pregnant or breast-feeding, ask a health professional before use.

Keep out of reach of children. In case of overdose, get medical help or contact a Poison Control Center right away. Prompt medical attention is critical for adults as well as for children even if you do not notice any signs or symptoms.

Directions

- do not take more than directed
- adults and children 12 years and over
- take 2 gelcaps every 6 hours while symptoms last

Drug Facts

Active ingredient
(in each gelcap)

Acetaminophen 500 mg

Purpose

Pain reliever/fever reducer

Uses

- temporarily relieves minor aches and pains due to:
 - the common cold
 - minor pain of arthritis
 - backache
 - muscular aches
 - premenstrual and menstrual cramps
- temporarily reduces fever

Warnings

Liver warning: This product contains acetaminophen. Severe liver damage may occur if you take

- more than 4,000 mg of acetaminophen in 24 hours
- with other drugs containing acetaminophen
- 3 or more alcoholic drinks every day while using this product

Allergy alert: Acetaminophen may cause severe skin reactions. Symptoms may include:

- skin redness
- hives
- itching
- blistering
- rash

If a skin reaction occurs, stop use and seek medical help right away.

Drug Facts (continued)

- do not take more than 6 gelcaps in 24 hours, unless directed by a doctor
- do not take for more than 10 days unless directed by a doctor
- children under 12 years: ask a doctor

Other information

- store at 25°C (77°F); excursions permitted between 15°-30°C (59°-86°F)
- avoid high humidity
- see end flap for expiration date and lot number

Inactive ingredients croscarmellose sodium, D&C red #33, FD&C blue #1, FD&C red #40, gelatin, hydroxypropyl cellulose, hypromellose, iron oxide black, iron oxide red, iron oxide yellow, polyethylene glycol, povidone, pregelatinized starch, propylene glycol, stearic acid, stearic acid, titanium dioxide

Questions or comments?

1-800-426-9391

*This product is not manufactured or distributed by Johnson & Johnson Corporation, owner of the registered trademark Extra Strength Tylenol® Rapid Release Gels. 50844 ORG032251921 Distributed by LNK INTERNATIONAL, INC. 60 Arkay Drive Hauppauge, NY 11788 USA

Quality Plus 44-519

PAIN RELIEVER EXTRA STRENGTH

acetaminophen tablet, coated

Product Information

Product Type	HUMAN OTC DRUG	Item Code (Source)	NDC:50844-519
Route of Administration	ORAL		

Active Ingredient/Active Moiety

Ingredient Name	Basis of Strength	Strength
ACETAMINOPHEN (UNII: 362O9ITL9D) (ACETAMINOPHEN - UNII:362O9ITL9D)	ACETAMINOPHEN	500 mg

Inactive Ingredients

Ingredient Name	Strength
CROSCARMELOSE SODIUM (UNII: M28OL1HH48)	
D&C RED NO. 33 (UNII: 9DBA0SBB0L)	
FD&C BLUE NO. 1 (UNII: H3R47K3TBD)	
FD&C RED NO. 40 (UNII: WZB9127XOA)	
GELATIN, UNSPECIFIED (UNII: 2G86QN327L)	
HYDROXYPROPYL CELLULOSE, UNSPECIFIED (UNII: 9XZ8H6N6OH)	
HYPROMELLOSE, UNSPECIFIED (UNII: 3NXW29V3WO)	
FERROSOFERRIC OXIDE (UNII: XM0M87F357)	
FERRIC OXIDE RED (UNII: 1K09F3G675)	
FERRIC OXIDE YELLOW (UNII: EX438O2MRT)	
POLYETHYLENE GLYCOL, UNSPECIFIED (UNII: 3WJQ0SDW1A)	
POVIDONE, UNSPECIFIED (UNII: FZ989GH94E)	
STARCH, CORN (UNII: O8232NY3SJ)	
PROPYLENE GLYCOL (UNII: 6DC9Q167V3)	
SHELLAC (UNII: 46N107B71O)	
STEARIC ACID (UNII: 4ELV7Z65AP)	
TITANIUM DIOXIDE (UNII: 15FIX9V2JP)	

Product Characteristics

Color	red, blue	Score	no score
Shape	OVAL	Size	19mm
Flavor		Imprint Code	L;5
Contains			

Packaging

#	Item Code	Package Description	Marketing Start Date	Marketing End Date
1	NDC:50844-519-02	1 in 1 CARTON	05/10/2004	
1		12 in 1 BOTTLE, PLASTIC; Type 0: Not a Combination Product		
2	NDC:50844-519-21	1 in 1 CARTON	05/10/2004	
2		16 in 1 BOTTLE, PLASTIC; Type 0: Not a Combination Product		
3	NDC:50844-519-12	100 in 1 BOTTLE, PLASTIC; Type 0: Not a Combination Product	11/14/2023	
4	NDC:50844-519-20	225 in 1 BOTTLE, PLASTIC; Type 0: Not a Combination Product	11/14/2023	
5	NDC:50844-519-15	1 in 1 CARTON	05/10/2004	03/21/2022
5		50 in 1 BOTTLE; Type 0: Not a Combination Product		

Marketing Information

Marketing Category	Application Number or Monograph Citation	Marketing Start Date	Marketing End Date
OTC Monograph Drug	M013	05/10/2004	

Labeler - L.N.K. International, Inc. (038154464)

Establishment

Name	Address	ID/FEI	Business Operations
LNK International, Inc.		038154464	manufacture(50844-519) , pack(50844-519)

Establishment

Name	Address	ID/FEI	Business Operations
LNK International, Inc.		832867837	manufacture(50844-519)

Establishment

Name	Address	ID/FEI	Business Operations
LNK International, Inc.		832867894	manufacture(50844-519)

Establishment

Name	Address	ID/FEI	Business Operations
LNK International, Inc.		868734088	manufacture(50844-519)

Establishment

Name	Address	ID/FEI	Business Operations
LNK International, Inc.		967626305	pack(50844-519)

Revised: 11/2023

L.N.K. International, Inc.