

ADAPALENE- adapalene gel
MEIJER, INC.

Adapalene Gel USP 0.1 percent

Active ingredient

Adapalene USP 0.1% (retinoid)*

*read consumer information leaflet

Purpose

Acne treatment

Use

- For the treatment of acne

Warnings

For external use only

Do not use

- on damaged skin (cuts, abrasions eczema, sunburn)
- if you are allergic to adapalene or any of the ingredients in this product

Pregnancy/Breastfeeding,

ask a doctor before use.

When using this product

- limit sun exposure, including light from tanning beds, and use sunscreen when going outdoors.
- do not wax to remove hair in areas where the product has been applied
- during the early weeks of use, your acne may appear to worsen before it improves (this is normal); continue using as directed, unless you get irritation that become severe
- irritation (redness, itching, dryness, burning) is more likely to occur:
 - in the first few weeks of use
 - if using more than one topical acne medication at a time
 - but irritation usually lessens with continued use of this product
- it may take up to 3 months of once daily use to see result
- avoid product contact eyes, lips, and mouth. If contact occurs immediately flush the area with water.
- wash hands after use

Stop use and ask a doctor if

- you become pregnant, or planning to become pregnant while using this product
- you have symptoms of an allergic reaction (such as itching, rash, hives, swelling of the lips, eyelids, and shortness of breath)
- irritation become severe
- you see no improvement after 3 months of once daily use

Keep out of reach of children.

If swallowed, get medical help or contact a Poison Control Center (1-800-222-1222) right away.

Directions

Adults and children 12 years of age and older:

- use once daily
- clean the skin gently and pat dry before applying the product
- cover the entire affected area with a thin layer. For example, if your acne is on the face, apply the product to the entire face.
- do not use more than one time a day. Applying more than directed will not provide faster or better results, but may worsen skin irritation.
- **Children under 12 years of age:** ask a doctor

Other information

- store at room temperature 68° to 77°F
- protect from freezing

Inactive ingredients

carbomer homopolymer, edetate disodium, methylparaben, poloxamer 182, propylene glycol, purified water, sodium hydroxide

Questions or comments?

Call **1-877-753-3935** Monday-Friday 9AM-5PM EST

Principal Display Panel

Compare to Differin® Gel Active Ingredient**

adapalene gel USP, 0.1%

ACNE TREATMENT

OIL FREE

FRAGRANCE FREE

PREVIOUSLY AVAILABLE

ONLY BY PRESCRIPTION

FDA-APPROVED

DERMATOLOGIST

DEVELOPED AND TESTED

ONCE DAILY TOPICAL RETINOID*

*Read consumer information leaflet before use

TAMPER EVIDENT: DO NOT USE IF SAFETY SEAL UNDER CAP IS BROKEN OR MISSING.

KEEP OUTER CARTON FOR COMPLETE WARNINGS AND PRODUCT INFORMATION.

**This product is not manufactured or distributed by Galderma Laboratories, L.P. distributor of Differin® Gel.

DIST.BY MEIJER

DISTRIBUTION, INC.

GRAND RAPIDS, MI 49544

www.meijer.com


Product Label

Drug Facts	
Active ingredient Adapalene USP 0.1% (retinoid)*	Purpose Acne treatment *read consumer information leaflet
Use ■ For the treatment of acne	
Warnings For external use only	
Do not use ■ on damaged skin (cuts, abrasions, eczema, sunburn) ■ if you are allergic to adapalene or any of the ingredients in this product	
If pregnant or breast-feeding, ask a doctor before use.	
When using this product ■ limit sun exposure, including light from tanning beds, and use sunscreen when going outdoors. ■ do not wax to remove hair in areas where the product has been applied ■ during the early weeks of use, your acne may appear to worsen before it improves (this is normal); continue using as directed, unless you get irritation that becomes severe ■ irritation (redness, itching, dryness, burning) is more likely to occur: ■ in the first few weeks of use ■ if using more than one topical acne medication at a time ■ but irritation usually lessens with continued use of this product ■ it may take up to 3 months of once daily use to see results ■ avoid product contact with eyes, lips, and mouth. If contact occurs, immediately flush the area with water. ■ wash hands after use	
Stop use and ask a doctor if ■ you become pregnant, or are planning to become pregnant while using the product ■ you have symptoms of an allergic reaction (such as itching, rash, hives, swelling of the lips, eyelids, and shortness of breath) ■ irritation becomes severe ■ you see no improvement after 3 months of once daily use	
Keep out of reach of children. If swallowed, get medical help or contact a Poison Control Center (1-800-222-1222) right away.	
Directions Adults and children 12 years of age and older: ■ use once daily ■ clean the skin gently and pat dry before applying the product ■ cover the entire affected area with a thin layer. For example, if your acne is on the face, apply the product to the entire face. ■ do not use more than one time a day. Applying more than directed will not provide faster or better results, but may worsen skin irritation. Children under 12 years of age: ask a doctor	
Other information ■ store at room temperature 68° to 77°F ■ protect from freezing	
Inactive ingredients carbomer homopolymer, edetate disodium, methylparaben, poloxamer 182, propylene glycol, purified water, sodium hydroxide	
Questions or comments? Call 1-877-753-3935 Monday-Friday 9AM-5PM EST	

**This product is not manufactured or distributed by Galderma Laboratories, L.P., distributor of Differin® Gel.

First FDA-Approved over-the-counter topical retinoid* for acne treatment

*Read carton and enclosed consumer information leaflet before using this product. Keep this carton and consumer information leaflet. They contain important information.



Actual Size

One Tube Inside

NDC 41250-288-05

meijer

**adapalene
gel USP, 0.1%**

ACNE TREATMENT

**OIL FREE
FRAGRANCE FREE**

**PREVIOUSLY AVAILABLE
ONLY BY PRESCRIPTION**

FDA-APPROVED

**DERMATOLOGIST
DEVELOPED AND TESTED**

ONCE DAILY TOPICAL RETINOID*

**Compare to Differin® Gel
Active Ingredient ****

*Read consumer information
leaflet before use

NET WT 0.5 OZ (15 g)

**TAMPER EVIDENT: DO NOT USE IF SAFETY
SEAL UNDER CAP IS BROKEN OR MISSING.**

**KEEP OUTER CARTON FOR COMPLETE
WARNINGS AND PRODUCT INFORMATION.**

**DIST. BY MEIJER
DISTRIBUTION, INC.
GRAND RAPIDS, MI 49544
www.meijer.com**
PRODUCT OF INDIA



WWW.MEIJER.COM/SATISFACTION

PID 4629990



7 19283 61507 7

PLD-C576A FC008724

Lot No.:

Exp. Date:

MEIJER Acne Treatment

ADAPALENE

adapalene gel

Product Information

Product Type	HUMAN OTC DRUG	Item Code (Source)	NDC:41250-288
Route of Administration	TOPICAL		

Active Ingredient/Active Moiety

Ingredient Name	Basis of Strength	Strength
ADAPALENE (UNII: 1L4806J2QF) (ADAPALENE - UNII:1L4806J2QF)	ADAPALENE	1 mg in 1 g

Inactive Ingredients

Ingredient Name	Strength
CARBOMER HOMOPOLYMER TYPE C (ALLYL PENTAERYTHRITOL CROSSLINKED) (UNII: 4Q93RCW27E)	
EDETATE DISODIUM (UNII: 7FLD91C86K)	
METHYLPARABEN (UNII: A2I8C7HI9T)	
POLOXAMER 182 (UNII: JX0HIX6OAG)	
PROPYLENE GLYCOL (UNII: 6DC9Q167V3)	
WATER (UNII: 059QF0KO0R)	
SODIUM HYDROXIDE (UNII: 55X04QC32I)	

Packaging

#	Item Code	Package Description	Marketing Start Date	Marketing End Date
1	NDC:41250-288-05	1 in 1 CARTON	11/29/2019	
1		15 g in 1 TUBE; Type 0: Not a Combination Product		

Marketing Information

Marketing Category	Application Number or Monograph Citation	Marketing Start Date	Marketing End Date
ANDA	ANDA091314	11/29/2019	

Labeler - MEIJER, INC. (006959555)

Revised: 7/2023

MEIJER, INC.