NIGHTTIME COUGH DM- dextromethorphan hbr, doxylamine succinate solution L.N.K. International. Inc.

Quality Plus 44-043

Active ingredients (in each 20 mL)

Dextromethorphan HBr 30 mg Doxylamine Succinate 12.5 mg

Purpose

Cough suppressant Antihistamine

Uses

- temporarily relieves cough due to minor throat and bronchial irritation as may occur with a cold
- temporarily relieves these symptoms due to hay fever or other upper respiratory allergies:
 - itching of the nose or throat
 - sneezing
 - itchy, watery eyes
 - runny nose
- controls the impulse to cough to help you sleep

Warnings

Do not use

if you are now taking a prescription monoamine oxidase inhibitor (MAOI) (certain drugs for depression, psychiatric or emotional conditions, or Parkinson's disease), or for 2 weeks after stopping the MAOI drug. If you do not know if your prescription drug contains an MAOI, ask a doctor or pharmacist before taking this product.

Ask a doctor before use if you have

- a cough that occurs with too much phlegm (mucus)
- glaucoma
- difficulty in urination due to enlargement of the prostate gland
- a breathing problem or persistent or chronic cough as occurs with smoking, asthma, chronic bronchitis, or emphysema

Ask a doctor or pharmacist before use if you are

taking sedatives or tranquilizers.

When using this product

- marked drowsiness may occur
- alcohol, sedatives, and tranquilizers may increase drowsiness
- avoid alcoholic drinks
- use caution when driving a motor vehicle or operating machinery
- excitability may occur, especially in children

Stop use and ask a doctor if

cough persists more than 7 days, tends to recur, or is accompanied by a fever, rash, or persistent headache. These could be signs of a serious condition.

If pregnant or breast-feeding,

ask a health professional before use.

Keep out of reach of children.

In case of overdose, get medical help or contact a Poison Control Center right away.

Directions

- do not take more than directed
- do not take more than 4 doses in any 24-hour period
- mL = milliliter
- only use the dose cup provided
- adults and children 12 years and over: 20 mL in dosing cup provided every 6 hours
- children under 12 years: do not use

Other information

- each 20 mL contains: sodium 14 mg
- store at 25°C (77°F); excursions permitted between 15°-30°C (59°-86°F)
- see end flap for expiration date and lot number

Inactive ingredients

anhydrous citric acid, FD&C blue #1, FD&C red #40, flavors, glycerin, high fructose corn syrup, polyethylene glycol, propylene glycol, purified water, sodium benzoate, sodium chloride, sodium citrate dihydrate, sucralose, sugar, xanthan gum

Questions or comments?

1-800-426-9391

Principal display panel

NDC 50844-043-36

Quality

Plus

Compare to the active ingredients in Robitussin® MAXIMUM STRENGTH Nighttime Cough DM*

MAXIMUM STRENGTH

NIGHTTIME COUGH DM

DEXTROMETHORPHAN HBr Cough suppressant DOXYLAMINE SUCCINATE Antihistamine

- Controls cough
- Relieves runny nose & sneezing

Menthol-Berry Flavor

Ages 12 Years and Over

4 FL OZ (118 mL)


Dosage cup included

*This product is not manufactured or distributed by PF Consumer Healthcare 1 LLC, owner of the registered trademark Robitussin® MAXIMUM STRENGTH Nighttime Cough DM. 50844 REV0123A04336

Distributed by LNK INTERNATIONAL, INC. 60 Arkay Drive Hauppauge, NY 11788 USA

TAMPER EVIDENT: DO NOT USE IF IMPRINTED SAFETY SEAL UNDER CAP IS BROKEN OR MISSING

TAMPER EVIDENT: DO NOT USE IF PRINTED NECK WRAP IS BROKEN OR MISSING





4 FL OZ (118 mL)

KEEP OUTER PACKAGE FOR COMPLETE PRODUCT INFORMATION

Drug Facts

Active ingredients (in each 20 mL)

Dextromethorphan HBr 30 mg...Cough suppressant Doxylamine succinate 12.5 mg... Antihistamine

Uses

- Uses

 Itemporarily relieves cough due to minor throat
 and bronchial irritation as may occur with a cold
 Itemporarily relieves these symptoms due to hay
 fever or other upper respiratory allergies.
 It itching of the nose or throat
 Intervention of the income of the control of the cold of the cold

Warnings
Do not use if you are now taking a prescription
monoamine oxidase inhibitor (MAOI) (certain drugs
for depression, psychiatric or emotional conditions,
or Parkinson's disease), or for 2 weeks after
stopping the MAOI drug. If you do not know if your
prescription drug contains an MAOI, ask a doctor
or pharmacist before taking this product.

- Ask a doctor before use if you have
 a cough that occurs with too much phlegm
 (mucus) glaucoma
 difficulty in urination due to enlargement of the
- prostate gland
- a breathing problem or persistent or chronic cough as occurs with smoking, asthma, chronic bronchitis, or emphysema

Ask a doctor or pharmacist before use if you are taking sedatives or tranquilizers.

- When using this product
 marked drowsiness may occur
 alcohol, sedatives, and tranquilizers may increase drowsiness
- avoid alcoholic drinks
 use caution when driving a motor vehicle or operating machinery

■ excitability may occur, especially in children Stop use and ask a doctor if cough persists more than 7 days, tends to recur, or is accompanied by a fever, rash, or persistent headache. These could

be signs of a serious condition If pregnant or breast-feeding, ask a health professional before use.

Drug Facts (continued)

Keep out of reach of children. In case of overdose. edical help or contact a Poison Control Center

Directions ■ do not take more than directed

- do not take more than 4 doses in any 24-hou period ■ mL = milliliter

- only use the dose cup provided
 adults and children 12 years and over: 20 mL in dosing cup provided every 6 hours
 children under 12 years: do not use

Other information

- each 20 mL contains: sodium 14 mg
 store at 25°C (77°F); excursions permitted between 15'-30°C (59'-86°F)
 see end flap for expiration date and lot number

Inactive ingredients anhydrous citric acid, FD&C blue #1, FD&C red #40, flavors, glycerin, high fructose corn syrup, polyethylene glycol, propylene glycol, purified water, sodium bezoate, sodium chloride, sodium citrate dihydrate, sucralose, sugar, xanthan gum

Questions or comments? 1-800-426-9391

*This product is not manufactured or distributed by GlaxoSmithKline Consumer Healthcare Holdings (US) LLC, owner of the registered trademark Robitussin® MAXIMUM STRENGTH Nighttime Cough DM. 50844 REV0123A04336

> TAMPER EVIDENT: DO NOT USE IF PRINTED NECK WRAP IS BROKEN OR MISSING

TAMPER EVIDENT: DO NOT USE IF IMPRINTED SAFETY SEAL UNDER CAP IS BROKEN OR MISSING

Distributed by

LNK INTERNATIONAL, INC.

60 Arkay Drive

Hauppauge, NY 11788



QUALITY PLUS

NDC 50844-043-36

Compare to the active ingredients in Robitussin® MAXIMUM STRENGTH Nighttime Cough DM*

MAXIMUM STRENGTH

NIGHTTIME (3) Cough DM

DEXTROMETHORPHAN HBr Cough suppressant

DOXYLAMINE SUCCINATE Antihistamine



Menthol-Berry

- Relieves runny nose & sneezing

Ages 12 Years and Over



4 FL OZ (118 mL)



NDC 50844-043-36

Compare to the active ingredients in Robitussin® MAXIMUM STRENGTH Nighttime Cough DM*


MAXIMUM STRENGTH

NIGHTTIME [3] Cough DM

DEXTROMETHORPHAN HBr Cough suppressant

DOXYLAMINE SUCCINATE

Antihistamine




- Relieves runny nose & sneezing

Menthol-Berry Flavor

Ages 12 Years and Over



4 FL OZ (118 mL)



B-1603-043-36 REV0123A04336



No Print / No Varnish Lot no. & Exp. date

Quality Plus 44-043

NIGHTTIME COUGH DM

dextromethorphan hbr, doxylamine succinate solution

Product Information

HUMAN OTC DRUG Item Code (Source) NDC:50844-043 **Product Type**

Active Ingredient/Active Moiety				
Ingredient Name	Basis of Strength	Strength		
DEXTROMETHORPHAN HYDROBROMIDE (UNII: 9D2RTI9KYH) (DEXTROMETHORPHAN - UNII:7355X3ROTS)	DEXTROMETHORPHAN HYDROBROMIDE	30 mg in 20 mL		
DOXYLAMINE SUCCINATE (UNII: V9BI9B5YI2) (DOXYLAMINE - UNII:95QB77JKPL)	DOXYLAMINE SUCCINATE	12.5 mg in 20 mL		

Ingredient Name ANHYDROUS CITRIC ACID (UNII: XF417D3PSL) FD&C BLUE NO. 1 (UNII: H3R47K3TBD) FD&C RED NO. 40 (UNII: WZB9127XOA) GLYCERIN (UNII: PDC6A3C0OX)				
FD&C BLUE NO. 1 (UNII: H3R47K3TBD) FD&C RED NO. 40 (UNII: WZB9127XOA) GLYCERIN (UNII: PDC6A3C0OX)				
FD&C RED NO. 40 (UNII: WZB9127XOA) GLYCERIN (UNII: PDC6A3C0OX)				
GLYCERIN (UNII: PDC6A3C0OX)				
HIGH FRUCTOSE CORN SYRUP (UNII: XY6UN3QB6S)				
POLYETHYLENE GLYCOL, UNSPECIFIED (UNII: 3WJQ0SDW1A)				
PROPYLENE GLYCOL (UNII: 6DC9Q167V3)				
WATER (UNII: 059QF0KO0R)				
SODIUM BENZOATE (UNII: OJ245FE5EU)				
SODIUM CHLORIDE (UNII: 451W47IQ8X)				
TRISODIUM CITRATE DIHYDRATE (UNII: B22547B95K)				
SUCRALOSE (UNII: 96K6UQ3ZD4)				
SUCROSE (UNII: C151H8M554)				
XANTHAN GUM (UNII: TTV12P4NEE)				

Product Characteristics				
Color	red (maroon)	Score		
Shape		Size		
Flavor	BERRY, MENTHOL	Imprint Code		
Contains				

Packaging					
	#	Item Code	Package Description	Marketing Start Date	Marketing End Date
	1	NDC:50844- 043-36	1 in 1 CARTON	07/16/2021	
	1		118 mL in 1 BOTTLE, PLASTIC; Type 0: Not a Combination Product		

Marketing Information				
Marketing Category	Application Number or Monograph Citation	Marketing Start Date	Marketing End Date	
OTC Monograph Drug	M012	07/16/2021		

Labeler - L.N.K. International, Inc. (038154464)

Establishment				
Name	Address	ID/FEI	Business Operations	
LNK International, Inc.		967626305	manufacture(50844-043) , pack(50844-043)	

Revised: 6/2023 L.N.K. International, Inc.