

**DRAMAMINE ORANGE- dimenhydrinate tablet, chewable**  
**Medtech Products Inc.**

*Disclaimer: Most OTC drugs are not reviewed and approved by FDA, however they may be marketed if they comply with applicable regulations and policies. FDA has not evaluated whether this product complies.*

-----

**Dramamine® Chewable Orange Motion Sickness Relief**

***Drug Facts***

***Active ingredient***

***(in each tablet)***

Dimenhydrinate 50 mg

**▣ *Purpose***

Antiemetic

**▣ *Use***

for prevention and treatment of these symptoms associated with motion sickness:

- nausea
- vomiting
- dizziness

**▣ *Warnings***

**▣ *Do not give to***

children under 2 years of age unless directed by a doctor

**Ask a doctor before use if you have**

- a breathing problem such as emphysema or chronic bronchitis
- glaucoma
- trouble urinating due to an enlarged prostate gland

**Ask a doctor or pharmacist before use if you are taking**

taking sedatives or tranquilizers

**When using this product**

- marked drowsiness may occur
- avoid alcoholic drinks
- alcohol, sedatives, and tranquilizers may increase drowsiness
- be careful when driving a motor vehicle or operating machinery

**If pregnant or breast-feeding,**

ask a doctor before use.

**Keep out of reach of children.**

In case of accidental overdose, get medical help or contact a Poison Control Center (1-800-222-1222) right away.

### ▯ **Directions**

- to prevent motion sickness, the first dose should be taken ½ to 1 hour before starting activity
- to prevent or treat motion sickness, see below:

---

#### **Adults and children 12 years and over**

- take 1 to 2 tablets every 4-6 hours
- do not take more than 8 tablets in 24 hours, or as directed by a doctor

#### **Children 6 to under 12 years**

- give ½ to 1 tablet every 6-8 hours
- do not give more than 3 tablets in 24 hours, or as directed by a doctor

#### **Children 2 to under 6 years**

- give ½ tablet every 6-8 hours
  - do not give more than 1 ½ tablets in 24 hours, or as directed by a doctor
- 

### ▯ **Other Information**

- Phenylketonurics: contains phenylalanine 0.75 mg per tablet
- store at room temperature 20-25°C (68-77°F)
- see side panel for lot number and expiration date
- **Tamper Evident:**▯ Do not use if pouch is torn or open.
- **do not use if carton is opened or if blister is broken or torn**

### ▯ **Inactive ingredients**

anhydrous citric acid, aspartame, FD&C yellow no. 6 aluminum lake, flavors, magnesium stearate, maltodextrin, methacrylic acid copolymer, modified food starch, sorbitol

### ▯ **Questions?**

Call ▯ **1-800-382-7219**

### **Dramamine® Chewable Orange Motion Sickness Relief, 8 ct**



# Dramamine®

## MOTION SICKNESS RELIEF

### DUAL ACTION

1. TREATS SYMPTOMS ON THE SPOT
2. PREVENTS NAUSEA, DIZZINESS & VOMITING



Convenient  
Chewable Tablets

8 ORANGE FLAVORED  
TABLETS (50 mg EACH)

**Dramamin**  
MOTION SICKNESS RELIEF

LOT XXXXXX  
EXP MM/XXXX

# Dramamine®

## MOTION SICKNESS RELIEF



**Drug Facts**

**Active ingredient (in each tablet)** Dimenhydrinate 50 mg. Antiemetic Purpose

**Use** For prevention and treatment of these symptoms associated with motion sickness: nausea, vomiting, dizziness

**Warnings**  
Do not give to children under 2 years of age unless directed by a doctor  
Ask a doctor before use if you have a breathing problem such as emphysema or chronic bronchitis, trouble urinating due to an enlarged prostate gland, glaucoma, or sedatives or tranquilizers may increase drowsiness. When using this product, marked drowsiness may occur. Avoid alcoholic drinks. Be careful when driving a motor vehicle or operating machinery. If pregnant or breast-feeding, ask a doctor before use. Keep out of reach of children. In case of accidental overdose, get medical help or contact a Poison Control Center (1-800-222-1222) right away.

**Directions**  
To prevent motion sickness, the first dose should be taken 1/2 to 1 hour before starting activity.  
To prevent or treat motion sickness, see below:

<b>Adults and children 12 years and over</b>	Take 1 to 2 tablets every 4-6 hours. Do not take more than 8 chewable tablets in 24 hours, or as directed by a doctor.
<b>Children 6 to under 12 years</b>	Give 1/2 to 1 tablet every 6-8 hours. Do not give more than 3 chewable tablets in 24 hours, or as directed by a doctor.
<b>Children 2 to under 6 years</b>	Give 1/2 tablet every 6-8 hours. Do not give more than 1 1/2 chewable tablets in 24 hours, or as directed by a doctor.

**Other information**  
Phenylethanolamine HCl: contains phenylethanolamine HCl 0.4 mg per tablet. Store at room temperature 68-77°F (20-25°C). Do not use if carton is opened or blister unit is broken or torn. See side panel for lot number and expiration date.

**Inactive ingredients** FD & C yellow #6 aluminum lake, flavors, magnesium stearate, maleic acid, polyethylene glycol, polyethylene glycol, sorbitol, FD & C yellow #6 aluminum lake, flavors, magnesium stearate, maleic acid, polyethylene glycol, polyethylene glycol, sorbitol.

**Questions or comments?** Call 1-800-382-7279

©2012 Dist. By: Medtech Products Inc., Irvington, NY 10533  
A Prestige Brands Company. Made in USA  
D/A004101

Dramamine® Chewable Orange Motion Sickness Relief, 4 ct

**#1** Pharmacist Recommended Brand

Dimenhydrinate Tablets/Antiemetic

**Dramamine**  
MOTION SICKNESS RELIEF

LOT: EXP:

**CHEWABLE FORMULA**

**Dramamine**  
MOTION SICKNESS RELIEF



**DUAL ACTION**

1. TREATS SYMPTOMS ON THE SPOT
2. PREVENTS NAUSEA, DIZZINESS & VOMITING

 Actual Size

**4 ORANGE FLAVORED TABLETS (50 mg EACH)**

8 31248 00567 6

©2017 Dr. Williams' Medicine, Inc. New York, NY 10041  
Made in Italy DRAM132

17-10-90-101

**Keep Out of Reach of Children**  
Information on Full Carton

**Dramamine**  
MOTION SICKNESS RELIEF

**Drug Facts**

**Active ingredient (in each tablet):** Dimenhydrinate (50 mg) Antiemetic

**Purpose:** Antiemetic

**Use:** For prevention and treatment of these symptoms associated with motion sickness: nausea, vomiting, dizziness

**Warnings:** Do not give to children under 2 years of age unless directed by a doctor

**Ask a doctor before use if you have:** a breathing problem such as emphysema or chronic bronchitis; glaucoma; trouble urinating due to an enlarged prostate gland

**Ask a doctor or pharmacist before use if you are taking:** sedatives or tranquilizers; alcohol, sedatives, and tranquilizers may increase drowsiness; be careful when driving a motor vehicle or operating machinery

**If pregnant or breast-feeding,** ask a doctor before use.

**Keep out of reach of children.** In case of overdose, get medical help or contact a Poison Control Center (1-800-222-1222) right away.

**Directions:** To prevent motion sickness, the first dose should be taken 1/2 to 1 hour before starting activity. To prevent or treat motion sickness, see below.

**Adults and children:** Take 1 to 2 tablets every 4-6 hours. Do not take more than 6 tablets in 24 hours, or as directed by a doctor.

**Children 6 to 12 years:** Take 1 tablet every 6-8 hours. Do not give more than 3 tablets in 24 hours, or as directed by a doctor.

**Children 2 to 6 years:** Give 1/2 tablet every 6-8 hours. Do not give more than 1 1/2 tablets in 24 hours, or as directed by a doctor.

**Other information:** Phenylethanolamine hydrochloride contains phenylethanolamine hydrochloride. Store at room temperature 20°-25°C (68°-77°F). See side panel for full prescribing information. Contains 4 orange flavored tablets. Do not use if pouch is torn or past expiration date. **Tamper Evident:** Do not use if pouch is torn or past expiration date.

**Inactive ingredients:** anhydrous citric acid, aspartame, FD&C yellow #6 aluminum lake, flavors, magnesium stearate, maltodextrin, methylsilylate, polyethylene glycol, polyethylene glycol, sorbitol.

**Questions? 1-800-382-7219**

**DRAMAMINE ORANGE**  
dimenhydrinate tablet, chewable

**Product Information**

<b>Product Type</b>	HUMAN OTC DRUG	<b>Item Code (Source)</b>	NDC:63029-902
<b>Route of Administration</b>	ORAL		

**Active Ingredient/Active Moiety**

<b>Ingredient Name</b>	<b>Basis of Strength</b>	<b>Strength</b>
DIMENHYDRINATE (UNII: JB937PER5C) (CHLORTHEOPHYLLINE - UNII:GE2UA340FM)	DIMENHYDRINATE	50 mg

**Inactive Ingredients**

<b>Ingredient Name</b>	<b>Strength</b>
ANHYDROUS CITRIC ACID (UNII: XF417D3PSL)	
ASPARTAME (UNII: Z0H242BBR1)	
FD&C YELLOW NO. 6 (UNII: H77VEI93A8)	
ALUMINUM OXIDE (UNII: LM26O6933)	
MAGNESIUM STEARATE (UNII: 70097M6I30)	
MALTODEXTRIN (UNII: 7CVR7L4A2D)	
SORBITOL (UNII: 506T60A25R)	

**Product Characteristics**

<b>Color</b>	orange	<b>Score</b>	2 pieces
<b>Shape</b>	ROUND	<b>Size</b>	12mm
<b>Flavor</b>		<b>Imprint Code</b>	
<b>Contains</b>			

**Packaging**

#	Item Code	Package Description	Marketing Start Date	Marketing End Date
1	NDC:63029-902-01	1 in 1 BOX	01/15/2012	
1		8 in 1 BLISTER PACK; Type 0: Not a Combination Product		
2	NDC:63029-902-02	2 in 1 POUCH; Type 0: Not a Combination Product	01/15/2012	
3	NDC:63029-902-04	1 in 1 BOX	03/30/2013	
3		4 in 1 BLISTER PACK; Type 0: Not a Combination Product		

**Marketing Information**

<b>Marketing Category</b>	<b>Application Number or Monograph Citation</b>	<b>Marketing Start Date</b>	<b>Marketing End Date</b>
OTC monograph final	part336	01/15/2012	

**Labeler** - Medtech Products Inc. (122715688)