

STOOL SOFTENER - docusate sodium capsule, liquid filled
Velocity Pharma

Disclaimer: Most OTC drugs are not reviewed and approved by FDA, however they may be marketed if they comply with applicable regulations and policies. FDA has not evaluated whether this product complies.

Docusate Sodium, USP 100mg

Active Ingredient
(in each softgel)

Docusate Sodium 100 mg

Purpose

Stool softener laxative

Uses

- for temporary relief of occasional constipation and irregularity
- this product generally produces bowel movement in 12 to 72 hours

Warnings

Ask a doctor before use if you have:

- stomach pain, nausea or vomiting
- a sudden change in bowel habits that lasts more than 2 weeks

Ask a doctor or pharmacist before use if you are presently taking mineral oil.

Stop use and ask a doctor if:

- you have rectal bleeding or no bowel movement after using this product. This could be signs of a serious condition.
- you need to use laxative for more than 1 week.

If pregnant or breast-feeding, ask a health professional before use.

Keep out of reach of children.

In case of overdose, get medical help or contact a Poison Control Center right away.

Directions

Take with glass of water

adults and children 12 years and over:

- 1 to 3 liquid gels daily. This dose may be taken as single daily dose or in divided doses.

Children under 12 years of age:

- 1 liquid gel daily

Children under 2 years of age

Ask a doctor

Other Information

- store at 15-30 °C (59-86 °F)
- protect from excessive humidity
- do not use this product if the safety seal under the cap is torn or missing

Inactive Ingredients

FD&C Red #40, FD&C Yellow #6, gelatin, glycerin, hypromellose, polyethylene glycol, propylene glycol, purified water, sorbitol special, titanium dioxide

Questions or Comments

Call 1-855-314-1850

PACKAGE LABEL.PRINCIPAL DISPLAY PANEL

183995
C1/1

639277838957
UPC

Drug Facts (continued)

Stop use and ask a doctor if:

- you have rectal bleeding or no bowel movement after using this product. These could be signs of a serious condition.
- you need to use a laxative for more than 1 week

If pregnant or breast-feeding, ask a health professional before use.

Keep out of reach of children. In case of overdose, get medical help or contact a Poison Control Center right away.

Directions

Take with a glass of water.

Adults and children 12 years and over:

- 1 to 3 liquid gels daily. This dose may be taken as a single daily dose or in divided doses.

Children under 12 years of age:
1 liquid gel daily

Children under 2 years of age:
ask a doctor ▶

Drug Facts (continued)

Other information:

- store at 15-30°C (59-86°F)
- protect from excessive humidity
- do not use this product if the safety seal under the cap is torn or missing

Inactive ingredients:

FD&C Red #40, FD&C Yellow #6, gelatin, glycerin, hypromellose, polyethylene glycol, propylene glycol, purified water, sorbitol special, titanium dioxide

Questions? Call
1-855-314-1850

Distributed by:
Greenbrier International, Inc.
Chesapeake, VA 23320

**This product is not manufactured or distributed by Dulcolax®

25 SOFTGELS

Docosate Sodium, USP 100mg

Gentle

Stool Softener



Compare to the active ingredient in Dulcolax® *

Drug Facts

SEE BOTTLE FOR ADDITIONAL WARNINGS AND PRODUCT INFORMATION.

Active ingredient

(in each softgel)	Purpose
Docosate sodium 100 mg	Stool softener laxative

Uses

- For temporary relief of occasional constipation and irregularity
- his product generally produces bowel movement in 12 to 72 hours

Warnings

Ask a doctor before use if you have:

- stomach pain, nausea or vomiting
- a sudden change in bowel habits that lasts more than 2 weeks

Ask a doctor or pharmacist before use if you are presently taking mineral oil. ▶

Gentle, Dependable
Overnight Relief

NDC 07188-049-05

5

NDC 076168-069-05

25
TABLETS

Docusate Sodium
USP 100mg

Gentle

Stool
Softener



Do not use if imprinted safety seal under cap is broken or missing.

Drug Facts

Active ingredient (in each tablet)	Purpose
Docusate sodium 100 mg	...Stool softener laxative

Uses

- For temporary relief of occasional constipation and irregularity
- This product generally produces bowel movement in 12 to 72 hours

Warnings: KEEP OUTER CARTON FOR COMPLETE WARNINGS AND PRODUCT INFORMATION.

- Ask a doctor or pharmacist before use
- Stop use and ask a doctor
- If pregnant or breast-feeding
- Keep out of reach of children

Directions: Take with a glass of water

Adults and children 12 years and over:	1 to 3 liquid gels daily. This dose may be taken as a single daily dose or in divided doses.
Children under 12 years of age:	1 liquid gel daily
Children under 2 years of age:	ask a doctor

Questions? 1-855-314-1850.

Distributed by:
Greenbrier International, Inc.
Chesapeake, VA 23320

183895

NDC: 076168-069-05 – 25 COUNT

STOOL SOFTENER

docusate sodium capsule, liquid filled

Product Information

Product Type	HUMAN OTC DRUG	Item Code (Source)	NDC:76 168-069
Route of Administration	ORAL		

Active Ingredient/Active Moiety

Ingredient Name	Basis of Strength	Strength
DOCUSATE SODIUM (UNII: F05Q2T2JA0) (DOCUSATE - UNII:M7P27195AG)	DOCUSATE SODIUM	100 mg

Inactive Ingredients

Ingredient Name	Strength
FD&C RED NO. 40 (UNII: WZB9127XOA)	
FD&C YELLOW NO. 6 (UNII: H77VEI93A8)	
GELATIN (UNII: 2G86QN327L)	
HYPROMELLOSES (UNII: 3NXW29V3WO)	
PROPYLENE GLYCOL (UNII: 6DC9Q167V3)	
SORBITOL (UNII: 506T60A25R)	
TITANIUM DIOXIDE (UNII: 15FIX9V2JP)	
WATER (UNII: 059QF0KO0R)	
GLYCERIN (UNII: PDC6A3C0OX)	

Product Characteristics

Color	RED	Score	no score
Shape	OVAL	Size	5mm
Flavor		Imprint Code	V100
Contains			

Packaging

#	Item Code	Package Description	Marketing Start Date	Marketing End Date
1	NDC:76 168-069-05	25 in 1 BOTTLE		
1		1 in 1 CARTON		

Marketing Information

Marketing Category	Application Number or Monograph Citation	Marketing Start Date	Marketing End Date
OTC monograph not final	part334	04/01/2013	

Labeler - Velocity Pharma (962198409)