

## **RYNEX PE- brompheniramine maleate and phenylephrine hcl liquid**

### **EDWARDS PHARMACEUTICALS, INC.**

*Disclaimer: Most OTC drugs are not reviewed and approved by FDA, however they may be marketed if they comply with applicable regulations and policies. FDA has not evaluated whether this product complies.*

-----

## **RYNEX PE**

### **Drug Facts**

<b>Active Ingredients</b> <b>(in each 5 ml teaspoonful)</b>	<b>Purpose</b>
Brompheniramine Maleate 1 mg	Antihistamine
Phenylephrine HCl 2.5 mg	Nasal Decongestant

### **Uses**

temporarily relieves these symptoms due to the common cold, hay fever (allergic rhinitis) or other respiratory allergies:

- runny nose
- sneezing
- itching of the nose or throat
- itchy, watery eyes
- nasal congestion
- reduces swelling of nasal passage

### **Warnings**

Do not exceed recommended dosage.

### **Do not take this product**

- if you are now taking a prescription monoamine oxidase inhibitor (MAOI) (certain drugs for depression, psychiatric, or emotional conditions, or Parkinson's disease), or for 2 weeks after stopping the MAOI drug. If you do not know if your prescription drug contains an MAOI, ask a doctor or pharmacist before taking this product.

### **Ask a doctor before use if you have**

- a breathing problem such as emphysema or chronic bronchitis
- glaucoma
- heart disease
- high blood pressure
- thyroid disease
- diabetes mellitus
- difficulty in urination due to enlargement of the prostate gland

Ask a doctor before use if you are taking sedatives or tranquilizers.

**When using this product**

- excitability may occur, especially in children
- may cause drowsiness
- sedatives and tranquilizers may increase drowsiness effect
- avoid alcoholic beverages
- use caution when driving a motor vehicle or operating machinery

**Stop use and ask doctor if**

- nervousness, dizziness, or sleeplessness occur
- If symptoms do not improve within 7 days or are accompanied by fever, consult a doctor
- new symptoms occur

If pregnant or breast feeding, ask a health professional before use.

Keep out of reach of children.

In case of accidental overdose, seek professional help or contact a Poison Control Center immediately.

**Directions**

Do not exceed recommended dosage.

Adults and children 12 years of age and over:	4 teaspoonfuls (20 mL) every 4 hours, not to exceed 24 teaspoonfuls in 24 hours, or as directed by a doctor.
Children 6 to under 12 years of age:	2 teaspoonfuls(10 mL) every 4 hours, not to exceed 1 teaspoonfuls in 24 hours, or as directed by a doctor.
Children under 6 years of age:	Consult a doctor

**Other information**

Store at 59° - 86° F (15° - 30° C) [see USP for Controlled Room Temperature].

**Inactive ingredients**

Bubblegum Flavor, Citric Acid, FD&C Red #40, Methyl Paraben, Potassium Citrate, Potassium Sorbate, Propyl Paraben, Propylene Glycol, Purified Water, Sorbitol Solution 70%, Sucralose.

**Questions Comments?**

Call 1-800-543-9560

**PRINCIPAL DISPLAY PANEL - 473 mL Bottle Label**

Rynex PE

E

NDC 0485-0202-16

Rynex PE

Antihistamine • Nasal Decongestant

Sugar Free • Alcohol Free

• Gluten Free

Each teaspoonful (5 mL)

for oral administration contains:

Brompheniramine Maleate 1 mg

Phenylephrine HCl 2.5 mg

Bubblegum Flavor

FOR PROFESSIONAL USE ONLY

Tamper evident by foil seal under cap.

Do not use if foil seal is broken or

missing.


Manufactured for:

EDWARDS

Pharmaceuticals, Inc.

Ripley, MS 38663

16oz. (473 mL)



NDC 0485-0202-16

## Rynex PE

Antihistamine • Nasal Decongestant

**Sugar Free • Alcohol Free  
• Gluten Free**

Each teaspoonful (5 mL)  
for oral administration contains:  
Brompheniramine Maleate ..... 1 mg  
Phenylephrine HCl ..... 2.5 mg

**Bubblegum Flavor**

Tamper evident by foil seal under cap.  
Do not use if foil seal is broken or  
missing.

Manufactured for:  
**EDWARDS**  
Pharmaceuticals, Inc.  
Ripley, MS 38663

**16oz. (473 mL)**

### Drug Facts

**Active Ingredients**  
(in each 5 mL teaspoonful)  
Brompheniramine Maleate 1 mg ..... Antihistamine  
Phenylephrine HCl 2.5 mg ..... Nasal Decongestant

**Purpose**

**Uses** temporarily relieves these symptoms due to the common cold, hay fever (allergic rhinitis) or other respiratory allergies: ■ runny nose ■ sneezing ■ itching of the nose or throat ■ itchy, watery eyes ■ nasal congestion ■ reduces swelling of nasal passages

**Warnings**  
Do not exceed recommended dosage.  
Do not take this product  
■ if you are now taking a prescription monoamine oxidase inhibitor (MAOI) (certain drugs for depression, psychiatric, or emotional conditions, or Parkinson's disease), or for 2 weeks after stopping the MAOI drug. If you do not know if your prescription drug contains an MAOI, ask a doctor or pharmacist before taking this product.

**Ask a doctor before use if you have**  
■ a breathing problem such as emphysema or chronic bronchitis ■ glaucoma ■ heart disease ■ high blood pressure ■ thyroid disease ■ diabetes mellitus ■ difficulty in urination due to enlargement of the prostate gland

**Ask a doctor before use if you are taking sedatives or tranquilizers.**

**When using this product**  
■ excitability may occur, especially in children ■ may cause drowsiness ■ sedatives and tranquilizers may increase drowsiness effect ■ avoid alcoholic beverages ■ use caution when driving a motor vehicle or operating machinery

**Stop use and ask doctor if**  
■ nervousness, dizziness, or sleeplessness occur

### Drug Facts (continued)

■ If symptoms do not improve within 7 days or are accompanied by fever, consult a doctor  
■ new symptoms occur

**If pregnant or breastfeeding**, ask a health professional before use.

**Keep out of reach of children.**  
In case of accidental overdose, seek professional help or contact a Poison Control Center immediately.

**Directions**  
Do not exceed recommended dosage.

Adults and children 12 years of age and over:	4 teaspoonfuls (20 mL) every 4 hours, not to exceed 24 teaspoonfuls in 24 hours, or as directed by a doctor.
Children 6 to under 12 years of age:	2 teaspoonfuls (10 mL) every 4 hours, not to exceed 12 teaspoonfuls in 24 hours, or as directed by a doctor.
Children under 6 years of age:	consult a doctor

**Other information**  
Store at 59° - 86° F (15° - 30° C) [see USP for Controlled Room Temperature].

**Inactive ingredients**  
Bubblegum flavor, citric acid, FD&C Red #40, methylparaben, potassium citrate, potassium sorbate, propylene glycol, propylparaben, purified water, sorbitol, sucralose

**Questions? Comments?** Call 1-800-543-9560

Rev. 08/19



N 3 04850 20216 5

## RYNEX PE

brompheniramine maleate and phenylephrine hcl liquid

### Product Information

<b>Product Type</b>	HUMAN OTC DRUG	<b>Item Code (Source)</b>	NDC:0485-0202
<b>Route of Administration</b>	ORAL		

### Active Ingredient/Active Moiety

Ingredient Name	Basis of Strength	Strength
<b>BROMPHENIRAMINE MALEATE</b> (UNII: IXA7C9ZN03) (BROMPHENIRAMINE - UNII:H57G17P2FN)	BROMPHENIRAMINE MALEATE	1 mg in 5 mL
<b>PHENYLEPHRINE HYDROCHLORIDE</b> (UNII: 04JA59TNSJ) (PHENYLEPHRINE - UNII:1WS297W6MV)	PHENYLEPHRINE HYDROCHLORIDE	2.5 mg in 5 mL

### Inactive Ingredients

Ingredient Name	Strength
<b>CITRIC ACID MONOHYDRATE</b> (UNII: 2968PHW8QP)	
<b>FD&amp;C RED NO. 40</b> (UNII: WZB9127XOA)	
<b>METHYLPARABEN</b> (UNII: A2I8C7HI9T)	
<b>POTASSIUM CITRATE</b> (UNII: EE90ONI6FF)	
<b>POTASSIUM SORBATE</b> (UNII: 1VPU26JZZ4)	
<b>PROPYLPARABEN</b> (UNII: Z8IX2SC1OH)	
<b>PROPYLENE GLYCOL</b> (UNII: 6DC9Q167V3)	

<b>WATER</b> (UNII: 059QF0KO0R)				
<b>SORBITOL</b> (UNII: 506T60A25R)				
<b>SUCRALOSE</b> (UNII: 96K6UQ3ZD4)				
<b>Product Characteristics</b>				
<b>Color</b>	pink		<b>Score</b>	
<b>Shape</b>			<b>Size</b>	
<b>Flavor</b>	BUBBLE GUM		<b>Imprint Code</b>	
<b>Contains</b>				
<b>Packaging</b>				
<b>#</b>	<b>Item Code</b>	<b>Package Description</b>	<b>Marketing Start Date</b>	<b>Marketing End Date</b>
1	NDC:0485-0202-16	473 mL in 1 BOTTLE, PLASTIC; Type 0: Not a Combination Product	03/07/2011	
<b>Marketing Information</b>				
<b>Marketing Category</b>	<b>Application Number or Monograph Citation</b>		<b>Marketing Start Date</b>	<b>Marketing End Date</b>
OTC monograph final	part341		03/07/2011	

**Labeler -** EDWARDS PHARMACEUTICALS, INC. (195118880)

Establishment			
Name	Address	ID/FEI	Business Operations
Llorens Pharmaceutical		037342305	manufacture(0485-0202)

Revised: 1/2023

EDWARDS PHARMACEUTICALS, INC.