

ECONAZOLE NITRATE- econazole nitrate cream Bryant Ranch Prepack

Econazole Nitrate Cream, 1%


Rx Only

For Topical Use Only

DESCRIPTION

Econazole Nitrate Cream contains the antifungal agent, econazole nitrate 1%, in a water-miscible base consisting of benzoic acid, butylated hydroxyanisole, mineral oil, peglicol 5 oleate, pegoxol-7 stearate, and purified water. The white to off-white soft cream is for topical use only.

Chemically, econazole nitrate is 1-[2-{(4-chlorophenyl)methoxy}-2-(2,4-dichlorophenyl)ethyl]-1H-imidazole mononitrate. Its structure is as follows:



CLINICAL PHARMACOLOGY

After topical application to the skin of normal subjects, systemic absorption of econazole nitrate is extremely low. Although most of the applied drug remains on the skin surface, drug concentrations were found in the stratum corneum which, by far, exceeded the minimum inhibitory concentration for *dermatophytes*. Inhibitory concentrations were achieved in the epidermis and as deep as the middle region of the dermis. Less than 1% of the applied dose was recovered in the urine and feces.

Microbiology: Econazole nitrate has been shown to be active against most strains of the following microorganisms, both *in vitro* and in clinical infections as described in the INDICATIONS AND USAGE section.

Dermatophytes

Epidermophyton floccosum

Microsporum audouini

Microsporum canis

Microsporum gypseum

Trichophyton

mentagrophytes

Trichophyton rubrum

Trichophyton tonsurans

Yeasts

Candida albicans

Malassezia furfur

Econazole nitrate exhibits broad-spectrum antifungal activity against the following organisms *in vitro*, **but the clinical significance of these data is unknown.**

Dermatophytes

Trichophyton verrucosum

Yeasts

Candida guilliermondii

Candida parapsilosis

Candida tropicalis

INDICATIONS AND USAGE

Econazole Nitrate Cream is indicated for topical application in the treatment of tinea pedis, tinea cruris, and tinea corporis caused by *Trichophyton rubrum*, *Trichophyton mentagrophytes*, *Trichophyton tonsurans*, *Microsporum canis*, *Microsporum audouini*, *Microsporum gypseum*, and *Epidermophyton floccosum*, in the treatment of cutaneous candidiasis, and in the treatment of tinea versicolor.

CONTRAINDICATIONS

Econazole Nitrate Cream is contraindicated in individuals who have shown hypersensitivity to any of its ingredients.

WARNINGS

Econazole Nitrate Cream is not for ophthalmic use.

PRECAUTIONS

General

If a reaction suggesting sensitivity or chemical irritation should occur, use of the medication should be discontinued.

For external use only. Avoid introduction of Econazole Nitrate Cream into the eyes.

Drug Interactions

Warfarin

Concomitant administration of econazole and warfarin has resulted in enhancement of anticoagulation effect. Most cases reported product application with use under occlusion, genital application, or application to large body surface area which may increase the systemic absorption of econazole nitrate. Monitoring of International Normalized Ratio (INR) and/or prothrombin time may be indicated especially for patients who apply econazole to large body surface areas, in the genital area, or under occlusion.

Carcinogenesis, mutagenesis, impairment of fertility

Long-term animal studies to determine carcinogenic potential have not been performed.

Oral administration of econazole nitrate in rats has been reported to produce prolonged gestation. Intravaginal administration in humans has not shown prolonged gestation or other adverse reproductive effects attributable to econazole nitrate therapy.

Pregnancy

Econazole nitrate has not been shown to be teratogenic when administered orally to mice, rabbits or rats. Fetotoxic or embryotoxic effects were observed in Segment I oral studies with rats receiving 10 to 40 times the human dermal dose. Similar effects were observed in Segment II or Segment III studies with mice, rabbits and/or rats receiving oral doses 80 or 40 times the human dermal dose.

Econazole nitrate should be used in the first trimester of pregnancy only when the physician considers it essential to the welfare of the patient. The drug should be used during the second and third trimesters of pregnancy only if clearly needed.

Nursing Mothers

It is not known whether econazole nitrate is excreted in human milk. Following oral administration of econazole nitrate to lactating rats, econazole and/or metabolites were excreted in milk and were found in nursing pups. Also, in lactating rats receiving large oral doses (40 or 80 times the human dermal dose), there was a reduction in post partum viability of pups and survival to weaning; however, at these high doses, maternal toxicity was present and may have been a contributing factor. Caution should be exercised when econazole nitrate is administered to a nursing woman.

Geriatric Use

Clinical studies of Econazole Nitrate Cream did not include sufficient numbers of subjects aged 65 and over to determine whether they respond differently from younger subjects. Other reported clinical experience has not identified differences in responses between the elderly and younger patients. In general, dose selection for an elderly patient should be cautious, usually starting at the low end of the dosing range, reflecting the greater frequency of decreased hepatic, renal, or cardiac function, and of concomitant disease or other drug therapy.

ADVERSE REACTIONS

During clinical trials, approximately 3% of patients treated with econazole nitrate cream, 1% reported side effects thought possibly to be due to the drug, consisting mainly of burning, itching, stinging, and erythema. One case of pruritic rash has also been reported.

OVERDOSE

Overdosage of econazole nitrate in humans has not been reported to date. In mice, rats, guinea pigs and dogs, the oral LD 50 values were found to be 462, 668, 272, and >160 mg/kg, respectively.

DOSAGE AND ADMINISTRATION

Sufficient Econazole Nitrate Cream, 1% should be applied to cover affected areas once daily in patients with tinea pedis, tinea cruris, tinea corporis, and tinea versicolor, and twice daily (morning and evening) in patients with cutaneous candidiasis.

Early relief of symptoms is experienced by the majority of patients and clinical improvement may be seen fairly soon after treatment is begun; however, candidal infections and tinea cruris and corporis should be treated for two weeks and tinea pedis for one month in order to reduce the possibility of recurrence. If a patient shows no clinical improvement after the treatment period, the diagnosis should be redetermined. Patients with tinea versicolor usually exhibit clinical and mycological clearing after two weeks of treatment.

HOW SUPPLIED

Econazole Nitrate Cream, 1% is available as follows:

15 g tube (NDC 72162-1410-4)

30 g tube (NDC 72162-1410-3)

85 g tube (NDC 72162-1410-2)

Store at 20-25°C (68-77°F) [see USP Controlled Room Temperature].

Repackaged/Relabeled by:

Bryant Ranch Prepack, Inc.

Burbank, CA 91504

Econazole Nitrate 1% Cream



Each gram contains: Econazole Nitrate 1%, in a water-miscible base consisting of benzoic acid, butylated hydroxyanisole, mineral oil, peglicol 5 oleate, pegoxol-7 stearate, and purified water.

Usual Dosage: Apply 1 to 2 times a day. PHARMACIST: Dispense the Patient Package Inserts: <https://dailymed.nlm.nih.gov/dailymed/>

Keep out of reach of children. Store at 20° to 25°C (68° to 77°F) [See USP Controlled Room Temperature].

For Topical Use Only. Not for Ophthalmic Use.

Directions for puncturing tube seal: Remove cap. Turn cap upside down and place puncture tip onto tube seal; Screw cap back to close.

NDC 72162-1410-4

Econazole Nitrate Cream

1%



Relabeled by:
Bryant Ranch Prepack, Inc.
Burbank, CA 91504 USA

Rx only
NET WT 15 g

Manufactured by:
Padagis



ECONAZOLE NITRATE

econazole nitrate cream

Product Information

Product Type	HUMAN PRESCRIPTION DRUG	Item Code (Source)	NDC:72162-1410(NDC:45802-466)
Route of Administration	TOPICAL		

Active Ingredient/Active Moiety				
Ingredient Name			Basis of Strength	Strength
ECONAZOLE NITRATE (UNII: H438WYN10E) (ECONAZOLE - UNII:6Z1Y2V4A7M)			ECONAZOLE NITRATE	10 mg in 1 g
Inactive Ingredients				
Ingredient Name				Strength
BENZOIC ACID (UNII: 8SKN0B0MIM)				
BUTYLATED HYDROXYANISOLE (UNII: REK4960K2U)				
MINERAL OIL (UNII: T5L8T28FGP)				
WATER (UNII: 059QF0KO0R)				
PEG-5 OLEATE (UNII: 0240V77G50)				
PEGOXOL 7 STEARATE (UNII: 3EW5AXE5X5)				
Packaging				
#	Item Code	Package Description	Marketing Start Date	Marketing End Date
1	NDC:72162-1410-4	1 in 1 CARTON	08/17/2023	
1		15 g in 1 TUBE; Type 0: Not a Combination Product		
2	NDC:72162-1410-3	1 in 1 CARTON	08/17/2023	
2		30 g in 1 TUBE; Type 0: Not a Combination Product		
3	NDC:72162-1410-2	1 in 1 CARTON	08/17/2023	
3		85 g in 1 TUBE; Type 0: Not a Combination Product		
Marketing Information				
Marketing Category		Application Number or Monograph Citation	Marketing Start Date	Marketing End Date
ANDA		ANDA076479	08/08/2006	

Labeler - Bryant Ranch Prepack (171714327)

Registrant - Bryant Ranch Prepack (171714327)

Establishment

Name	Address	ID/FEI	Business Operations
Bryant Ranch Prepack		171714327	REPACK(72162-1410) , RELABEL(72162-1410)