BASIC CARE CHILDRENS PAIN AND FEVER- acetaminophen suspension Amazon.com Services LLC

Amazon Children's Pain & Fever Drug Facts

Active ingredient (in each 5 mL)

Acetaminophen 160 mg

Purpose

Pain reliever/fever reducer

Uses

temporarily:

- reduces fever
- relieves minor aches and pains due to:
- · the common cold
- flu
- headache
- sore throat
- toothache

Warnings

Liver warning: This product contains acetaminophen. Severe liver damage may occur if your child takes

- more than 5 doses in 24 hours, which is the maximum daily amount
- with other drugs containing acetaminophen

Allergy alert: Acetaminophen may cause severe skin reactions. Symptoms may include:

- skin reddening
- blisters
- rash

If a skin reaction occurs, stop use and seek medical help right away.

Sore throat warning: If sore throat is severe, persists for more than 2 days, is accompanied or followed by fever, headache, rash, nausea, or vomiting, consult a doctor promptly.

Do not use

- with any other drug containing acetaminophen (prescription or nonprescription). If you are not sure whether a drug contains acetaminophen, ask a doctor or pharmacist.
- if your child has ever had an allergic reaction to this product or any of its ingredients

Ask a doctor before use if your child has

liver disease

Ask a doctor or pharmacist before use if your child is

taking the blood thinning drug warfarin

When using this product

do not exceed recommended dose (see overdose warning)

Stop use and ask a doctor if

- pain gets worse or lasts more than 5 days
- fever gets worse or lasts more than 3 days
- new symptoms occur
- redness or swelling is present

These could be signs of a serious condition.

Keep out of reach of children.

Overdose warning: In case of overdose, get medical help or contact a Poison Control Center right away (1-800-222-1222). Quick medical attention is critical even if you do not notice any signs or symptoms.

Directions

- this product does not contain directions or complete warnings for adult use
- do not give more than directed (see overdose warning)
- shake well before using
- mL = milliliter
- find right dose on chart below. If possible, use weight to dose; otherwise, use age.
- remove the child protective cap and squeeze your child's dose into the dosing cup
- repeat dose every 4 hours while symptoms last
- do not give more than 5 times in 24 hours

Weight (lb)	Age (yr)	Dose (mL)*
under 24	nder 24 under 2 years ask	
24-35	2-3 years	5 mL
36-47	4-5 years	7.5 mL
48-59	6-8 years	10 mL
60-71	9-10 years	12.5 mL

I.	-	
72-95	11 years	15 mL

^{*}or as directed by a doctor

Attention: use only enclosed dosing cup specifically designed for use with this product. Do not use any other dosing device.

Other information

- each 5 mL contains: sodium 3 mg
- store at 20-25°C (68-77°F)
- · do not use if printed neckband is broken or missing

Inactive ingredients

anhydrous citric acid, calcium sulfate, carrageenan, FD&C red #40, flavor, glycerin, high fructose corn syrup, hydroxyethyl cellulose, microcrystalline cellulose and carboxymethylcellulose sodium, propylene glycol, purified water, sodium benzoate, sorbitol solution, tribasic sodium phosphate

Questions or comments?

1-800-719-9260

Principal Display Panel

amazon

basic care

For Ages 2 to 11

Compare to Children's Tylenol® active ingredient

Children's Pain & Fever

Acetaminophen 160 mg per 5 mL

Oral Suspension Liquid

Pain Reliever

Fever Reducer

Fast, effective relief of children's pain and fever

Aspirin Free

Ibuprofen Free


Alcohol Free

Paraben Free

Cherry Flavor

Dosage Cup Included

4 FL OZ (118 mL)



amazon

basic@care

For Ages 2 to 11

Children's Pain & Fever



Active ingredient (in each 5 mL)

Purpose

cetaminophen 160 mg......Pain reliever/fever reducer

Uses temporarily:

- USES Temporous

 reduces fever

 relieves minor aches and pains due to:

 the common cold

 fit headache

 over throat

 toothache

Warnings Liver warning: This product contains acetaminophen.

- Severe liver damage may occur if your child takes
 more than 5 doses in 24 hours, which is the maximum
 daily amount
 with other drugs containing acetaminophen
- Allergy alert: Acetaminophen may cause severe skin reactions. Symptoms may include:

 skin redening bisters brash
 If a skin reaction occurs, stop use and seek medical help

right away.

Sore throat warning: If sore throat is severe, persists for more than 2 days, is accompanied or followed by fever, headache, rash, nausea, or vomiting, consult a doctor promptly.

- Do not use
 with any other drug containing acetaminophen
 (prescription or nonprescription). If you are not sure whether a drug contains acetaminophen, ask a doctor or pharmacist.

 if your child has ever had an allergic reaction to this product or any of its ingredients

Ask a doctor before use if your child has liver disease

Ask a doctor or pharmacist before use if your child is taking the blood thinning drug warfarin When using this product do not exceed recommended

lose (see overdose warning) Stop use and ask a doctor if

- pain gets worse or lasts more than 5 days fever gets worse or lasts more than 3 days
- new symptoms occur
- redness or swelling is present These could be signs of a serious condition.

Keep out of reach of children.

Overdose warning: In case of overdose, get medical help or contact a Poison Control Center right away (1-800-222-1222). Quick medical attention is critical even if you do not notice any signs or symptoms.

Drug Facts (continued)

Directions

- this product does not contain directions or complete warnings for adult use
 do not give more than directed (see overdose
- warning)
 shake well before using
 mL = milliliter

- In IL = Infillier
 Initially dose on chart below. If possible, use weight to dose; otherwise, use age.
 In remove the child protective cap and squeeze your child's dose into the dosing cup
 repeat dose every 4 hours while symptoms last
 do not give more than 5 times in 24 hours

٠.	do not give more dian o timeo in E-1 nodro				
	Weight (lb)	Age (yr)	Dose (mL)*		
	under 24	under 2 years	ask a doctor		
	24-35	2-3 years	5 mL		
	36-47	4-5 years	7.5 mL		
	48-59	6-8 years	10 mL		
	60-71	9-10 years	12.5 mL		
	72-95	11 years	15 mL		

*or as directed by a doctor

Attention: use only enclosed dosing cup specifically designed for use with this product. Do not use any other dosing device.

Other information

- each 5 mL contains: sodium 3 mg
 store at 20-25°C (68-77°F)
 do not use if printed neckband is broken or

Inactive ingredients anhydrous citric acid, calcium sulfate, carrageenan, FD&C red #40, flavor, glycerin, high fructose corn syrup, hydroxyethyl cellulicse, microcrystalline cellulose and carboxymethylcellulose sodium, propylene glycol, purified water, sodium benzoate, sorbitol solution,

Questions or comments? 1-800-719-9260

**This product is not manufactured or distributed by Johnson & Johnson Consumer Inc., distributor of Children's Tylenol®.

GLUTEN FREE

amazon

basic @care

For Ages 2 to 11

NDC 72288-175-26

Compare to Children's Tylenol[®]

Children's Pain & Fever

Acetaminophen 160 mg per 5 mL Oral Suspension Liquid

Pain Reliever Fever Reducer

Fast, effective relief of children's pain and fever

Aspirin Free Ibuprofen Free **Alcohol Free** Paraben Free





Dosage Cup Included

4 FL OZ (118 mL)

DIRECTIONS: STEP 1. Shake well





STEP 2. Remove child protective cap



STEP 3. Squeeze vour child's dose into the dosing cup



IMPORTANT:

Please see dosing chart or ask a doctor for your child's correct dose.

DISTRIBUTED BY: AMAZON.COM SERVICES LLC 410 TERRY AVENUE N. SEATTLE, WA 98109 SEAT ILE, WA 98109

2022 AMAZON.COM, INC. OR ITS
AFFILIATES. ALL RIGHTS RESERVED.
BASIC CARE AND ALL RELATED LOGOS
ARE TRADEMARKS OF AMAZON.COM,
INC. OR ITS AFFILIATES.







17526 BN C2

acetaminophen suspension

Product Information

Product Type HUMAN OTC DRUG Item Code (Source) NDC:72288-175

Route of Administration ORAL

Active Ingredient/Active Moiety

Ingredient Name	Basis of Strength	Strength
CETAMINORUEN (UNIV. 20200ITIOD) (ACETAMINORUEN, UNIV. 20200ITIO	ACETAMINODUEN	160

ACETAMINOPHEN (UNII: 36209ITL9D) (ACETAMINOPHEN - UNII: 36209ITL9D) ACETAMINOPHEN 160 mg in 5 mL

Inactive Ingredients

Ingredient Name	Strength
ANHYDROUS CITRIC ACID (UNII: XF417D3PSL)	
CALCIUM SULFATE, UNSPECIFIED FORM (UNII: WATODDB505)	

CALCIUM SULFATE, UNSPECIFIED FORM (UNII: WATUUDBSUS

CARRAGEENAN (UNII: 5C69YCD2YJ)
FD&C RED NO. 40 (UNII: WZB9127XOA)

GLYCERIN (UNII: PDC6A3C0OX)

HIGH FRUCTOSE CORN SYRUP (UNII: XY6UN3QB6S)
MICROCRYSTALLINE CELLULOSE (UNII: OP1R32D61U)

CARBOXYMETHYLCELLULOSE SODIUM, UNSPECIFIED (UNII: K6790BS311)

PROPYLENE GLYCOL (UNII: 6DC9Q167V3)

WATER (UNII: 059QF0KO0R)

SODIUM BENZOATE (UNII: OJ245FE5EU)

SORBITOL (UNII: 506T60A25R)

SODIUM PHOSPHATE, TRIBASIC (UNII: A752Q30A6X)

HYDROXYETHYL CELLULOSE, UNSPECIFIED (UNII: T4V6TWG28D)

Product Characteristics

Color	RED (opaque)	Score	
Shape		Size	
Flavor	CHERRY	Imprint Code	
Contains			

Packaging

#	Item Code	Package Description	Marketing Start Date	Marketing End Date
1	NDC:72288-175- 26	1 in 1 CARTON	03/04/2021	
1		118 mL in 1 BOTTLE; Type 0: Not a Combination Product		

Marketing Information

Marketing Application Number or Monograph Marketing Start Marketing End

Category	Citation	Date	Date
OTC Monograph Drug	M013	03/04/2021	

Labeler - Amazon.com Services LLC (128990418)

Revised: 5/2024 Amazon.com Services LLC