


**NEOMYCIN AND POLYMYXIN B SULFATES AND DEXAMETHASONE- neomycin sulfate, polymyxin b sulfate and dexamethasone suspension**  
**Aidarex Pharmaceuticals LLC**

-----  
**Neomycin and Polymyxin B Sulfates and Dexamethasone Ophthalmic Suspension**  
**Sterile**

**DESCRIPTION**

Neomycin and Polymyxin B Sulfates and Dexamethasone Ophthalmic Suspension is a multiple dose anti-infective steroid combination in sterile suspension form for topical application. The chemical structure for the active ingredient, dexamethasone, is:




MW = 392.45


Established name: dexamethasone

Chemical name: pregna-1, 4-diene-3, 20-dione, 9-fluoro-11, 17, 21-trihydroxy-16-methyl-, (11 $\beta$ , 16 $\alpha$ )-.

The other active ingredients are neomycin sulfate and polymyxin B sulfate. The structural formula for neomycin sulfate is:



The structural formula for polymyxin B sulfate is:



**Each mL contains: Actives:** neomycin sulfate equivalent to neomycin 3.5 mg, polymyxin B sulfate 10,000 units, dexamethasone 0.1%. **Inactives:** hypromellose 2910 0.5%, sodium chloride, polysorbate 20, hydrochloric acid and/or sodium hydroxide (to adjust pH), purified water, benzalkonium chloride 0.004% (preservative).

## CLINICAL PHARMACOLOGY

Corticosteroids suppress the inflammatory response to a variety of agents and they probably delay or slow healing. Since corticosteroids may inhibit the body's defense mechanism against infection, a concomitant antimicrobial drug may be used when this inhibition is considered to be clinically significant in a particular case.

When a decision to administer both a corticosteroid and an antimicrobial is made, the administration of such drugs in combination has the advantage of greater patient compliance and convenience, with the added assurance that the appropriate dosage of both drugs is administered, plus assured compatibility of ingredients when both types of drugs are in the same formulation and, particularly, that the correct volume of drug is delivered and retained.

The relative potency of corticosteroids depends on the molecular structure, concentration and release from the vehicle.

## INDICATIONS AND USAGE

For steroid-responsive inflammatory ocular conditions for which a corticosteroid is indicated and where bacterial infection or a risk of bacterial infection exists.

Ocular corticosteroids are indicated in inflammatory conditions of the papebral and bulbar conjunctiva, cornea, and anterior segment of the globe where the inherent risk of corticosteroids use in certain infective conjunctivitides is accepted to obtain a diminution in edema and inflammation. They are also indicated in chronic anterior uveitis and corneal injury from chemical, radiation or thermal burns; or penetration of foreign bodies.

The use of a combination drug with an anti-infective component is indicated where the risk of infection is high or where there is an expectation that potentially dangerous numbers of bacteria will be present in the eye.

The particular anti-infective drug in this product is active against the following common bacterial eye pathogens: *Staphylococcus aureus*, *Escherichia coli*, *Haemophilus influenzae*, *Klebsiella/Enterobacter* species, *Neisseria* species, and *Pseudomonas aeruginosa*.

This product does not provide adequate coverage against: *Serratia marcescens* and streptococci, including *Streptococcus pneumoniae*.

## **CONTRAINDICATIONS**

Neomycin and Polymyxin B Sulfates and Dexamethasone Ophthalmic Suspension is contraindicated in most viral diseases of the cornea and conjunctiva, including epithelial herpes simplex keratitis (dendritic keratitis), vaccinia, varicella, and also in mycobacterial infection of the eye and fungal diseases of ocular structures. Neomycin and Polymyxin B Sulfates and Dexamethasone Ophthalmic Suspension is also contraindicated in individuals with known or suspected hypersensitivity to any of the ingredients of this preparation and to other corticosteroids.

## **WARNINGS**

Prolonged use of corticosteroids may result in glaucoma with damage to the optic nerve, defects in visual acuity and fields of vision, and in posterior subcapsular cataract formation. Prolonged use may also suppress the host immune response and thus increase the hazard of secondary ocular infections.

Various ocular diseases and long-term use of topical corticosteroids have been known to cause corneal and scleral thinning. Use of topical corticosteroids in the presence of thin corneal or scleral tissue may lead to perforation.

Acute purulent infections of the eye may be masked or activity enhanced by the presence of corticosteroid medication.

If this product is used for 10 days or longer, intraocular pressure should be routinely monitored even though it may be difficult in children and uncooperative patients. Steroids should be used with caution in the presence of glaucoma. Intraocular pressure should be checked frequently.

The use of steroids after cataract surgery may delay healing and increase the incidence of bleb formation.

Use of ocular steroids may prolong the course and may exacerbate the severity of many viral infections of the eye (including herpes simplex). Employment of a corticosteroid medication in the treatment of patients with a history of herpes simplex requires great caution; frequent slit lamp microscopy is recommended.

Neomycin and Polymyxin B Sulfates and Dexamethasone Ophthalmic Suspension is not for injection. It should never be injected subconjunctivally, nor should it be directly introduced into the anterior chamber of the eye.

Products containing neomycin sulfate may cause cutaneous sensitization.

## **PRECAUTIONS**

### **General**

The initial prescription and renewal of the medication order beyond 20 mL of Neomycin and Polymyxin B Sulfates and Dexamethasone Ophthalmic Suspension should be made by a physician only after examination of the patient with the aid of magnification, such as slit lamp biomicroscopy and, where appropriate, fluorescein staining. If signs and symptoms fail to improve after two days, the patient should be re-evaluated.

As fungal infections of the cornea are particularly prone to develop coincidentally with long-term corticosteroid applications, fungal invasion should be suspected in any persistent corneal ulceration where a corticosteroid has been used or is in use. Fungal cultures should be taken when appropriate.

If this product is used for 10 days or longer, intraocular pressure should be monitored (see WARNINGS).

Prolonged use of topical anti-bacterial agents may give rise to overgrowth of nonsusceptible organisms including fungi.

## **Information for Patients**

If inflammation or pain persists longer than 48 hours or becomes aggravated, the patient should be advised to discontinue use of the medication and consult a physician.

This product is sterile when packaged. To prevent contamination, care should be taken to avoid touching the bottle tip to eyelids or to any other surface. The use of this bottle by more than one person may spread infection. Keep bottle tightly closed when not in use. Keep out of reach of children.

## **Carcinogenesis, Mutagenesis, Impairment of Fertility**

Long-term studies in animals to evaluate carcinogenic or mutagenic potential have not been conducted with polymyxin B sulfate. Treatment of cultured human lymphocytes *in vitro* with neomycin increased the frequency of chromosome aberrations at the highest concentration (80 µg/mL) tested. However, the effects of neomycin on carcinogenesis and mutagenesis in humans are unknown.

Polymyxin B has been reported to impair the motility of equine sperm, but its effects on male or female fertility are unknown.

## **Pregnancy**

### **Teratogenic Effects**

#### **Pregnancy Category C**

Dexamethasone has been shown to be teratogenic in mice and rabbits following topical ophthalmic application in multiples of the therapeutic dose.

In the mouse, corticosteroids produce fetal resorptions and a specific abnormality, cleft palate. In the rabbit, corticosteroids have produced fetal resorptions and multiple abnormalities involving the head, ears, limbs, palate, etc.

There are no adequate or well-controlled studies in pregnant women. Neomycin and Polymyxin B Sulfates and Dexamethasone Ophthalmic Suspension should be used during pregnancy only if the potential benefit to the mother justifies the potential risk to the embryo or fetus. Infants born of mothers who have received substantial doses of corticosteroids during pregnancy should be observed carefully for signs of hypoadrenalism.

## **Nursing Mothers**

Systemically administered corticosteroids appear in human milk and could suppress growth, interfere with endogenous corticosteroid production, or cause other untoward effects. It is not known whether topical administration of corticosteroids could result in sufficient systemic absorption to produce detectable quantities in human milk. Because many drugs are excreted in human milk, caution should be exercised when Neomycin and Polymyxin B Sulfates and Dexamethasone Ophthalmic Suspension is administered to a nursing woman.

## **Pediatric Use**

Safety and effectiveness in pediatric patients below the age of 2 years have not been established.

## **Geriatric Use**

No overall differences in safety or effectiveness have been observed between elderly and younger patients.

## **ADVERSE REACTIONS**

Adverse reactions have occurred with corticosteroid/anti-infective combination drugs which can be attributed to the corticosteroid component, the anti-infective component, or the combination. Exact

incidence figures are not available since no denominator of treated patients is available.

Reactions occurring most often from the presence of the anti-infective ingredient are allergic sensitizations. The reactions due to the corticosteroid component are: elevation of intraocular pressure (IOP) with possible development of glaucoma, and infrequent optic nerve damage; posterior subcapsular cataract formation; and delayed wound healing.

Corticosteroid-containing preparations have also been reported to cause perforation of the globe. Keratitis, conjunctivitis, corneal ulcers, and conjunctival hyperemia have occasionally been reported following use of steroids.

## **DOSAGE AND ADMINISTRATION**

One to two drops in the conjunctival sac(s). In severe disease, drops may be used hourly, being tapered to discontinuation as the inflammation subsides. In mild disease, drops may be used up to four to six times daily.

Not more than 20 mL should be prescribed initially and the prescription should not be refilled without further evaluation as outlined in PRECAUTIONS above.

FOR TOPICAL OPHTHALMIC USE ONLY.

## **HOW SUPPLIED**

Neomycin and Polymyxin B Sulfates and Dexamethasone Ophthalmic Suspension is supplied as a sterile ophthalmic suspension in a white, low density polyethylene DROP-TAINER\* dispenser with a pink, polypropylene cap.

5 mL in 8 mL bottle

## **STORAGE**

Store upright at 8° - 27°C (46° - 80° F).

Rx Only

\*DROP-TAINER is a registered trademark of Alcon Manufacturing, Ltd.

Revised: April 2008

Printed in USA

### **Repackaged by:**

Aidarex Pharmaceuticals, LLC

Corona, CA 92880

### **Dist. By:**

**FALCON Pharmaceuticals, Ltd.**

Fort Worth, Texas 76134 USA

Mfd. By:

ALCON LABORATORIES, INC

Fort Worth, Texas 76134 USA

**9003573-0408**

## **PRINCIPAL DISPLAY PANEL**

NDC 33261-0551

Rx Only

Repackaged by:

Aidarex Pharmaceuticals, LLC

Corona, CA 92880

FALCON

PHARMACEUTICALS®

Neomycin and Polymyxin B Sulfates and Dexamethasone Ophthalmic Suspension

5 mL STERILE

AFFILIATE OF

ALCON LABORATORIES, INC.

QUALITY RX

CAUTION: FEDERAL LAW PROHIBITS DISPENSING WITHOUT PRESCRIPTION.  
SEE PACKAGE INSERT: KEEP OUT OF REACH OF CHILDREN. STORE  
AT 20-25°C (68-77°F) [SEE USP CONTROLLED ROOM TEMPERATURE].

**NEO/POLY B SULF./DEXAM., USP**

**5ML**

GENERIC FOR : MAXITROL

NDC: 33261-0551-01

LOT:

Packaged By:

EACH ML CONTAINS THE FOLLOWING  
ACTIVE INGREDIENTS:  
NEOMYCIN SULFATE  
EQUIVALENT TO NEOMYCIN 3.5MG  
POLYMYXIN B SULFATE 10,000U  
DEXAMETHASONE 0.1%  
(BENZALKONIUM CHLORIDE AS A  
PRESERVATIVE) 0.004%

OPHTHALMIC SUSPENSION

PUT \_\_\_ DROP(S) IN EYE(S) \_\_\_ TIMES DAILY  
PONGA \_\_\_ GOT(A)S EN OJO \_\_\_ VEGES AL DIA

MFG: ALCON LABORATORIES, INC.  
FORT WORTH, TX 76134  
INS 61314-0630-06

NEO/POLY B SULF./DEXAM., USP  
LOT: EX:  
NDC: 33261-0551-01  
RX 1001787397 5ML

NEO/POLY B SULF./DEXAM., USP  
LOT: EX:  
NDC: 33261-0551-01  
RX 1001787397 5ML

INS 61314-0630-06  
NEO/POLY B SULF./DEXAM., USP  
NDC: 33261-0551-01  
RX 1001787397 5ML

RX1001787397 EX:

PEEL HERE  
LIFTING TAB  
DO NOT  
CHART  
PEEL HERE  
BILL

**Aidarex**  
PHARMACEUTICALS

33261-0551\_NEO-POLY-B-SULF-DEXAM\_5ML

## NEOMYCIN AND POLYMYXIN B SULFATES AND DEXAMETHASONE

neomycin sulfate, polymyxin b sulfate and dexamethasone suspension

### Product Information

Product Type	HUMAN PRESCRIPTION DRUG	Item Code (Source)	NDC:33261-551(NDC:61314-630)
Route of Administration	OPHTHALMIC		

### Active Ingredient/Active Moiety

Ingredient Name	Basis of Strength	Strength
NEOMYCIN SULFATE (UNII: 057Y626693) (NEOMYCIN - UNII:I16QD7X297)	NEOMYCIN	3.5 mg in 1 mL
POLYMYXIN B SULFATE (UNII: 19371312D4) (POLYMYXIN B - UNII:J2VZ07J96K)	POLYMYXIN B	10000 [USP'U] in 1 mL
DEXAMETHASONE (UNII: 7S5I7G3JQL) (DEXAMETHASONE - UNII:7S5I7G3JQL)	DEXAMETHASONE	1 mg in 1 mL

## Inactive Ingredients

Ingredient Name	Strength
<b>BENZALKONIUM CHLORIDE</b> (UNII: F5UM2KM3W7)	
<b>HYPROMELLOSE 2910 (4000 MPA.S)</b> (UNII: RN3152OP35)	
<b>SODIUM CHLORIDE</b> (UNII: 451W47IQ8X)	
<b>POLYSORBATE 20</b> (UNII: 7T1F30V5YH)	
<b>HYDROCHLORIC ACID</b> (UNII: QTT17582CB)	
<b>SODIUM HYDROXIDE</b> (UNII: 55X04QC32I)	
<b>WATER</b> (UNII: 059QF0KO0R)	

## Packaging

#	Item Code	Package Description	Marketing Start Date	Marketing End Date
1	NDC:33261-551-01	5 mL in 1 BOTTLE, PLASTIC; Type 0: Not a Combination Product	11/13/2017	

## Marketing Information

Marketing Category	Application Number or Monograph Citation	Marketing Start Date	Marketing End Date
ANDA	ANDA062341	11/13/2017	

**Labeler** - Aidarex Pharmaceuticals LLC (801503249)

## Establishment

Name	Address	ID/FEI	Business Operations
Aidarex Pharmaceuticals		801503249	RELABEL(33261-551)

Revised: 8/2018

Aidarex Pharmaceuticals LLC