

IBUPROFEN- ibuprofen tablet
Kinray, Inc.

Drug Facts

Active ingredient (in each tablet)

Ibuprofen 200 mg (NSAID)*

*nonsteroidal anti- inflammatory drug

Purpose

Pain Reliever/ Fever Reducer

Uses

temporarily relieves minor aches and pain due to:

- headache
- toothache
- backache
- menstrual cramps
- the common cold
- muscular aches
- minor pain of arthritis
- temporarily reduces fever

Warnings

Allergy alert:

Ibuprofen may cause a severe allergic reaction, especially in people allergic to Aspirin. Symptoms may include

- hives
- facial swelling
- asthma (wheezing)
- shock
- skin reddening
- rash
- blisters

If an allergic reaction occurs, stop use and seek medical help right away.

Stomach bleeding warning:

This product contains an NSAID, which may cause severe stomach bleeding. The chance is higher if you

- are age 60 or older
- have had stomach ulcers or bleeding problems
- take a blood thinning (anticoagulant) or steroid drug

- take other drugs containing prescription or non prescription NSAIDs (aspirin, ibuprofen, naproxen, or others)
- have 3 or more alcoholic drinks everyday while using this product
- take more or for a longer time than directed

Do not use

- if you have ever had an allergic reaction to any other pain reliever/ fever reducer
- right before or after heart surgery

Ask a doctor before use if

- stomach bleeding warning applies to you
- you have problems or serious side effects from taking pain relievers or fever reducers
- you have a history of stomach problems, such as heartburn
- you have high blood pressure, heart disease, liver cirrhosis, kidney disease or asthma
- you are taking a diuretic

Ask a doctor or pharmacist before use if you are

- under a doctor's care for any serious condition
- taking aspirin for heart attack or stroke, because ibuprofen may decrease this benefit of aspirin
- taking any other drug

when using this product

- take with food or milk if stomach upset occurs
- the risk of heart attack or stroke may increase if you use more than directed or for longer than directed

Stop use and ask a doctor if

- you experience any of the following signs of stomach bleeding:
- feel faint
- vomit blood
- have bloody or black stools
- have stomach pain that does not get better
- pain gets worse or last for more than 10 days
- fever gets worse or last more than 3 days
- redness or swelling is present in the painful area
- any new symptoms appear

If pregnant or breast feeding,

ask a health professional before use. It is especially important not to use ibuprofen during the last 3 months of pregnancy unless definitely directed to do so by a doctor because it may cause problems in the unborn child or complications during delivery.

Keep out of reach of children.

In case of overdose, get medical help or contact a Poison Control Center right away (1-800-222-1222).

Directions

- **do not take more than directed**
- **the smallest effective dose should be used**

adults and children 12 years and older	<ul style="list-style-type: none"> • take 1 tablet every 4 to 6 hours while symptoms persist • if pain or fever does not respond to 1 tablet, 2 tablets may be used • do not exceed 6 tablets in 24 hours, unless directed by a doctor
children under 12 years	<ul style="list-style-type: none"> • consult a doctor

Other information

- **Tamper Evident: do not use if safety seal under cap is broken or missing**
- store at room temperature (20°- 25°C)
- avoid excessive heat above 40°C (104°F)

Inactive Ingredients:

colloidal silicon dioxide, croscarmellose sodium, FD&C Yellow #6 aluminum lake, magnesium stearate, microcrystalline cellulose, polyethylene glycol, polyvinyl alcohol, pregelatinized starch, talc & titanium dioxide

Questions? Adverse drug event call: (866)562-2756 Mon-Fri: 8 AM to 4 PM

*This product is not manufactured or distributed by McNeil Consumer Healthcare, owner of the registered trademark MOTRIN® IB.

NDC 61715-168-50

*Compare to the active ingredient in MOTRIN® IB.

Preferred plus Pharmacy

IBUPROFEN

Tablets USP 200mg

Pain Reliever / Fever Reducer (NSAID)

50 Tablets


Drug Facts (continued)
Keep out of reach of children. In case of overdose, get medical help or contact a Poison Control Center right away (1-800-222-1222).

Directions
 ■ Do not take more than directed ■ The smallest effective dose should be used
 adults and children 12 years of age and over Take 1 tablet every 4 to 6 hours while symptoms persist. If pain or fever does not respond to 1 tablet, 2 tablets may be used. Do not exceed 6 tablets in 24 hours unless directed by a doctor.
 children under 12 years consult a doctor

Other information
 ■ Tamper Evident: do not use if safety seal under cap is broken or missing ■ store at room temperature (20°-25°C) ■ avoid excessive heat above 40°C (104°F)

Inactive ingredients
 colloidal silicon dioxide, croscarmellose sodium, FD&C Yellow #6 aluminum lake, magnesium stearate, microcrystalline cellulose, polyethylene glycol, polyvinyl alcohol, pregelatinized starch, talc & titanium dioxide

Questions? Adverse drug event call: (866) 562-2756 Mon-Fri: 8 AM to 4 PM



All Preferred Plus Pharmacy products are distributed nationwide. This product is not manufactured or distributed by McKel Consumer Healthcare, owner of the registered trademark MOTRIN® IB, NDC 61715-168-50. © 2011 Kinray, Inc. All rights reserved. Preferred Plus, Preferred Plus + and the Preferred Plus logo are trademarks and/or registered trademarks of Kinray Inc. All other marks are property of their respective owners.

0 6 1715 16850 8

Lot No.:
Exp Dt.:


Drug Facts

Active ingredient (in each tablet)	Purpose
Ibuprofen 200mg (NSAID)*	Pain Reliever/ Fever Reducer
*nonsteroidal anti-inflammatory drug	

Uses
 ■ temporarily relieves minor aches and pains due to: ■ headache ■ toothache ■ backache
 ■ menstrual cramps ■ the common cold ■ muscular aches ■ minor pain of arthritis
 ■ temporarily reduces fever

Warnings
Allergy alert: Ibuprofen may cause a severe allergic reaction, especially in people allergic to Aspirin. Symptoms may include ■ hives ■ facial swelling ■ asthma (wheezing) ■ shock ■ skin reddening ■ rash ■ blisters
 If an allergic reaction occurs, stop use and seek medical help right away.
Stomach bleeding warning: This product contains an NSAID, which may cause severe stomach bleeding. The chance is higher if you ■ are age 60 or older ■ have had stomach ulcers or bleeding problems ■ take a blood-thinning (anticoagulant) or steroid drug ■ take other drugs containing prescription or nonprescription NSAIDs (aspirin, ibuprofen, naproxen, or others) ■ have 3 or more alcoholic drinks every day while using this product ■ take more or for a longer time than directed

Do not use ■ if you have ever had an allergic reaction to any other pain reliever/fever reducer



Drug Facts (continued)
 ■ right before or after heart surgery


Ask a doctor before use if ■ stomach bleeding warning applies to you ■ you have problems or serious side effects from taking pain relievers or fever reducers ■ you have a history of stomach problems, such as heartburn ■ you have high blood pressure, heart disease, liver cirrhosis, kidney disease, or asthma ■ you are taking a diuretic

Ask a doctor or pharmacist before use if you are ■ under a doctor's care for any serious condition ■ taking aspirin for heart attack or stroke, because ibuprofen may decrease this benefit of aspirin ■ taking any other drug

When using this product ■ take with food or milk if stomach upset occurs ■ the risk of heart attack or stroke may increase if you use more than directed or for longer than directed

Stop use and ask a doctor if ■ you experience any of the following signs of stomach bleeding:
 ■ feel faint ■ vomit blood ■ have bloody or black stools ■ have stomach pain that does not get better
 ■ pain gets worse or lasts more than 10 days ■ fever gets worse or lasts more than 3 days
 ■ redness or swelling is present in the painful area
 ■ any new symptoms appear

If pregnant or breast-feeding, ask a health professional before use. It is especially important not to use ibuprofen during the last 3 months of pregnancy unless definitely directed to do so by a doctor because it may cause problems in the unborn child or complications during delivery.



IBUPROFEN
 ibuprofen tablet

Product Information			
Product Type	HUMAN OTC DRUG	Item Code (Source)	NDC:61715-168
Route of Administration	ORAL		
Active Ingredient/Active Moiety			
	Ingredient Name	Basis of Strength	Strength

IBUPROFEN (UNII: WK2XYI10QM) (IBUPROFEN - UNII:WK2XYI10QM)	IBUPROFEN	200 mg
--	-----------	--------

Inactive Ingredients

Ingredient Name	Strength
SILICON DIOXIDE (UNII: ETJ7Z6XBU4)	
CROSCARMELOSE SODIUM (UNII: M28OL1HH48)	
FD&C YELLOW NO. 6 (UNII: H77VEI93A8)	
MAGNESIUM STEARATE (UNII: 70097M6I30)	
CELLULOSE, MICROCRYSTALLINE (UNII: OP1R32D61U)	
POLYETHYLENE GLYCOL 3350 (UNII: G2M7P15E5P)	
POLYVINYL ALCOHOL (UNII: 532B59J990)	
STARCH, PREGELATINIZED CORN (UNII: O8232NY3SJ)	
TALC (UNII: 7SEV7J4R1U)	
TITANIUM DIOXIDE (UNII: 15FIX9V2JP)	

Product Characteristics

Color	orange	Score	no score
Shape	ROUND	Size	10mm
Flavor		Imprint Code	115
Contains			

Packaging

#	Item Code	Package Description	Marketing Start Date	Marketing End Date
1	NDC:61715-168-50	1 in 1 CARTON		
1		50 in 1 BOTTLE, PLASTIC; Type 0: Not a Combination Product		
2	NDC:61715-168-51	1 in 1 CARTON		
2		100 in 1 BOTTLE, PLASTIC; Type 0: Not a Combination Product		

Marketing Information

Marketing Category	Application Number or Monograph Citation	Marketing Start Date	Marketing End Date
ANDA	ANDA091237	03/01/2016	

Labeler - Kinray, Inc. (012574513)

Registrant - Pharbest Pharmaceuticals, Inc. (557054835)

Establishment

Name	Address	ID/FEI	Business Operations
Pharbest Pharmaceuticals, Inc.		557054835	repack(61715-168) , relabel(61715-168)

