GOOD NEIGHBOR PHARMACY IBUPROFEN- ibuprofen suspension Amerisource Bergen

Amerisource Bergen Ibuprofen Oral Suspension Drug Facts

Active ingredient (in each 5 mL = 1 teaspoonful)

Ibuprofen 100 mg (NSAID)*

*nonsteroidal anti-inflammatory drug

Purposes

Pain reliever/fever reducer

Uses

temporarily:

- relieves minor aches and pains due to the common cold, flu, sore throat, headache and toothache
- reduces fever

Warnings

Allergy alert: Ibuprofen may cause a severe allergic reaction, especially in people allergic to aspirin. Symptoms may include:

- hives
- facial swelling
- asthma (wheezing)
- shock
- skin reddening
- rash
- blisters

If an allergic reaction occurs, stop use and seek medical help right away.

Stomach bleeding warning: This product contains an NSAID, which may cause severe stomach bleeding. The chance is higher if your child:

- · has had stomach ulcers or bleeding problems
- takes a blood thinning (anticoagulant) or steroid drug
- takes other drugs containing prescription or nonprescription NSAIDs (aspirin, ibuprofen, naproxen, or others)
- takes more or for a longer time than directed

Sore throat warning: Severe or persistent sore throat or sore throat accompanied by high fever, headache, nausea, and vomiting may be serious. Consult doctor promptly. Do not use more than 2 days or administer to children under 3 years of age unless directed by doctor.

Do not use

- if the child has ever had an allergic reaction to any other pain reliever/fever reducer
- right before or after heart surgery

Ask a doctor before use if

- stomach bleeding warning applies to your child
- child has a history of stomach problems, such as heartburn
- child has problems or serious side effects from taking pain relievers or fever reducers
- child has not been drinking fluids
- child has lost a lot of fluid due to vomiting or diarrhea
- child has high blood pressure, heart disease, liver cirrhosis, or kidney disease
- child has asthma
- child is taking a diuretic

Ask a doctor or pharmacist before use if the child is

- under a doctor's care for any serious condition
- taking any other drug

When using this product

- give with food or milk if stomach upset occurs
- the risk of heart attack or stroke may increase if you use more than directed or for longer than directed

Stop use and ask a doctor if

- child experiences any of the following signs of stomach bleeding
- feels faint
- vomits blood
- has bloody or black stools
- has stomach pain that does not get better
- the child does not get any relief within first day (24 hours) of treatment
- fever or pain gets worse or lasts more than 3 days
- redness or swelling is present in the painful area
- any new symptoms appear

Keep out of reach of children.

In case of overdose, get medical help or contact a Poison Control Center right away. (1-800-222-1222)

Directions

- this product does not contain directions or complete warnings for adult use
- do not give more than directed
- shake well before using
- find right dose on chart. If possible, use weight to dose; otherwise use age.
- use only enclosed measuring cup
- if needed, repeat dose every 6-8 hours

- do not use more than 4 times a day
- replace original bottle cap to maintain child resistance
- wash dosage cup after each use

Dosing Chart			
Weight (lbs)	Age(yrs)	Dose (teaspoonful or mL)	
under 2 years		ask a doctor	
24-35 lbs 2-3 years		1 tsp or 5 mL	
36-47 lbs	4-5 years	1 ½ tsp or 7.5 mL	
48-59 lbs	6-8 years	2 tsp or 10 mL	
60-71 lbs 9-10 years		2 ½ tsp or 12.5 mL	
72-95 lbs 11 years		3 tsp or 15 mL	

Other information

- each teaspoon contains: sodium 2 mg
- do not use if printed neckband is broken or missing
- store at 20-25°C (68-77°F)
- do not freeze
- see bottom panel for lot number and expiration date

Inactive ingredients

anhydrous citric acid, artificial mixed berry flavor, D&C yellow #10, FD&C red #40, glycerin, high fructose corn syrup, hypromellose, polysorbate 80, purified water, sodium benzoate, sorbitol solution, xanthan gum

Questions or comments?

1-800-719-9260

Principal Display Panel

Compare to Children's Motrin® active ingredient

FOR AGES 2 TO 11

children's

Ibuprofen Oral Suspension

100 mg per 5 mL (teaspoonful)

pain reliever/fever reducer (NSAID)

Lasts up to 8 hours

berry flavor

Alcohol Free



children's

Ibuprofen

Oral Suspension

100 mg per 5 mL (teaspoonful)

pain reliever/ fever reducer (NSAID)

NDC 24385-905-26

Compare to Children's Motrin® active ingredient



FOR AGES 2 TO 11

children's

lbuprofen

Oral Suspension

100 mg per 5 mL (teaspoonful)

pain reliever/ fever reducer (NSAID)

Lasts up to 8 hours

berry flavor

Alcohol Free

4 fl oz (120 mL)



Important: Read all product information before using. Keep this box for important information. This product is intended for use in children.

Drug Facts

Active ingredient Purposes (in each 5 mL = 1 teaspoonful)

Ibuprofen 100 mg (NSAID)*.....Pain reliever/ fever reducer

'nonsteroidal anti-inflammatory drug

Uses temporarily:

- relieves minor aches and pains due to the common cold, flu, sore throat, headache and toothache
- reduces fever

Warnings

Allergy alert: Ibuprofen may cause a severe allergic reaction, especially in people allergic to aspirin. Symptoms may include:

- facial swelling
- asthma (wheezing)
 shock
- skin reddening
- rash
- blisters

If an allergic reaction occurs, stop use and seek medical help right away.

Stomach bleeding warning: This product contains an NSAID, which may cause severe stomach bleeding. The chance is higher if your child:


- has had stomach ulcers or bleeding problems
- takes a blood thinning (anticoagulant) or steroid drug
- takes other drugs containing prescription or nonprescription NSAIDs (aspirin, ibuprofen, naproxen, or others)
- takes more or for a longer time than directed Sore throat warning: Severe or persistent sore throat or sore throat accompanied by high fever, headache, nausea, and vomiting may be serious. Consult doctor promptly. Do not use more than 2 days or administer to children under 3 years of age unless directed by doctor.

Do not use


- if the child has ever had an allergic reaction to any other pain reliever/fever reducer
- right before or after heart surgery

Ask a doctor before use if

- stomach bleeding warning applies to your child
- child has a history of stomach problems,



Ibuprofen Oral Suspension Carton Image 1





Drug Facts (continued)

- child has problems or serious side effects from taking pain relievers or fever reducers
- child has not been drinking fluids
- child has lost a lot of fluid due to vomiting or diarrhea
- child has high blood pressure, heart disease, liver cirrhosis, or kidney disease
- child has asthma
- child is taking a diuretic

Ask a doctor or pharmacist before use if the child is

- under a doctor's care for any serious condition
- taking any other drug

When using this product

- give with food or milk if stomach upset occurs
- the risk of heart attack or stroke may increase if you use more than directed or for longer than directed

Stop use and ask a doctor if

- child experiences any of the following signs of stomach bleeding
 - feels faint
 - vomits blood
 - has bloody or black stools
- has stomach pain that does not get better
- the child does not get any relief within first day (24 hours) of treatment
- fever or pain gets worse or lasts more than 3 days
- redness or swelling is present in the painful area
- any new symptoms appear

Keep out of reach of children. In case of overdose, get medical help or contact a Poison Control Center right away. (1-800-222-1222)

Directions

- this product does not contain directions or complete warnings for adult use
- do not give more than directed
- shake well before using
- find right dose on chart. If possible, use weight to dose; otherwise use age.
- use only enclosed measuring cup
- if needed, repeat dose every 6-8 hours
- do not use more than 4 times a day
- replace original bottle cap to maintain child resistance
- wash dosage duo after each use

Drug Facts (continued)

Dosing Chart				
Weight (lbs) Age (yrs) Dose (teaspoonful or m				
under 2	2 years	ask a doctor		
24-35 lbs	2-3 years	1 tsp or 5 mL		
36-47 lbs	4-5 years	1 1/2 tsp or 7.5 mL		
48-59 lbs	6-8 years	2 tsp or 10 mL		
60-71 lbs	9-10 years	2 1/2 tsp or 12.5 mL		
72-95 lbs	11 years	3 tsp or 15 mL		

Other information

- each teaspoon contains: sodium 2 mg
- do not use if printed neckband is broken or missing
- store at 20-25°C (68-77°F)
- do not freeze
- see bottom panel for lot number and expiration date

Inactive ingredients anhydrous citric acid, artificial mixed berry flavor, D&C yellow #10, FD&C red #40, glycerin, high fructose corn syrup, hypromellose, polysorbate 80, purified water, sodium benzoate, sorbitol solution, xanthan gum


Questions or comments? 1-800-719-9260

"Good Neighbor Pharmacy® Ibuprofen Oral Suspension 100 mg per 5 mL (teaspoonful) is not manufactured or distributed by McNeil Consumer Healthcare, distributor of Children's Motrin®.

GLUTEN FREE

Distributed By AmerisourceBergen 1300 Morris Drive Chesterbrook, PA 19087 Visit us at www.goodneighborpharmacy.com





Ibuprofen Oral Suspension Carton Image 2

GOOD NEIGHBOR PHARMACY IBUPROFEN

ibuprofen suspension

Product Information				
Product Type	HUMAN OTC DRUG LABEL	Item Code (Source)	NDC:24385-905	
Route of Administration	ORAL	DEA Schedule		

Active Ingredient/Active Moiety			
Ingredient Name	Basis of Strength	Strength	
IBUPRO FEN (IBUPRO FEN)	IBUPRO FEN	100 mg in 5 mL	

Inactive Ingredients			
Ingredient Name	Strength		
ANHYDRO US CITRIC ACID			
GLYCERIN			
HYPROMELLOSES			
POLYSORBATE 80			
WATER			
SO DIUM BENZO ATE			
SORBITOL			
XANTHAN GUM			
HIGH FRUCTO SE CORN SYRUP			
D&C YELLOW NO. 10			
FD&C RED NO. 40			

Product Characteristics			
Color	ORANGE	Score	

Shape		Size	
Flavor	BERRY	Imprint Code	
Contains			

P	Packaging					
#	Item Code	Package Description	Marketing Start Date	Marketing End Date		
1	NDC:24385-905-34	1 in 1 CARTON				
1		240 mL in 1 BOTTLE				
2	NDC:24385-905-26	1 in 1 CARTON				
2		120 mL in 1 BOTTLE				
3	NDC:24385-905-28	1 in 1 CARTON				
3		148 mL in 1 BOTTLE				

Marketing Information				
Marketing Category Application Number or Monograph Citation Marketing Start Date Marketing End				
ANDA	ANDA074937	01/28/1999		

Labeler - Amerisource Bergen (007914906)

Revised: 11/2012 Amerisource Bergen